TABLE OF CONTENTS

CONTENTS	GE
ACKNOWLEDGEMENT	ii
ABSTRACT	iv
ABSTRAK	vi
LIST OF TABLES	xiii
LIST OF FIGURES	xviii
LIST OF PLATES	xxi
LIST OF APPENDICES	xxii
1. INTRODUCTION	1
1.1. BACKGROUND ON CHILLI	1
1.2. THE IMPORTANCE OF CHILLIES	3
1.3. CHILLI INDUSTRY IN MALAYSIA 1.3.1. CONSTRAINTS OF DRIED CHILLI PRODUCTION IN MALAYSIA	4 9
1.4. BASIS OF THE STUDIES	10
2. LITERATURE REVIEW	12
2.1. TAXONOMY AND BASIC BIOLOGY OF CAPSICUM 2.1.1 CLASSIFICATION 2.1.2 BASIC BIOLOGY OF CAPSICUM ANNUUM L 2.1.2.1. Species characteristics 2.1.2.2. Breeding system 2.1.2.3. Chemical composition and nutritive elements in chilli a) Capsaicinoids b) Colour	12 12 13 13 13 14 14 17
c) Vitaminsd) Other nutritive and	18
essential elements	19

2.2.	THE IMPORTANCE OF CHILLI	20
	2.2.1. ECONOMIC AND NUTRITIVE	
	IMPORTANCE	20
	2.2.2. WORLD PRODUCTION OF CHILLI	23
	2.2.3. CHILLI PRODUCTION IN MALAYSIA	26
2.3.	MAJOR PROBLEMS OF CHILLI PRODUCTION	27
	2.3.1. PESTS AND DISEASES	
	2.3.2. INFERIOR VARIETIES	28
	2.3.3. ALLELOPATHIC EFFECTS OF RED	
	CHILLI (CAPSICUM ANNUUM L.)	29
2.4.	FACTORS AFFECTING THE QUALITY OF DRY	29
	2.4.1. GENETIC FACTORS	30
	2.4.2. ENVIRONMENTS	31
	2.4.2.1. Geographical areas	31
		31
	2.4.2.2 Temperature	32
	2.4.2.3 Rainfall (Water availability)	
	2.4.2.4. Growing conditions	
	2.4.2.5. Shading (Light intensity)	33
	2.4.2.6. Disease	. 33
	2.4.3. HARVESTING STAGE OR MATURITY AGE	. 33
	2.4.4. POST HARVEST HANDLING	34
	2.4.4.1. Drying	34
	2.4.4.2. Storage	35
2.5.	GENETIC AND BREEDING STUDIES FOR CHILLI	
	IMPROVEMENT	35
	2.5.1. GERMPLASM EVALUATION	35
	2.5.2. GENETIC STUDIES	36
	2.5.3. CORRELATION STUDIES	39
	2.5.4. GENOTYPE ENVIRONMENTAL	
	INTERACTION STUDIES ON YIELD AND	
	QUALITY OF CHILLI	41
	2.5.5. BREEDING STRATEGIES	42
	2.5.5.1. Heterosis	43
	2.5.5.2. Wide-crosses	. 43
	2.5.5.3. Biotechnology	44
2.6.	GENETIC X EVIRONMENT INTERACTION	46
	2.6.1. INTERPRETATION OF G x E	47
	2.6.2. SIGNIFICANT OF G x E	
	2.6.2 ANALYCIC OF C., E INTERACTION	51

		2.6.3.1. Analysis of variance	52
		2.6.3.2. Comparison of variance components	53
			54
		2.6.3.4. Heritability estimates	56
		2.6.4 STABILITY	59
			59
			60
			61
			62
			63
	2.7.	AGRO-ECOLOGICAL FEATURES OF PENINSULAR	
			64
		2.7.1. SOIL	64
		2.7.2. CLIMATE	64
	2.8.	STATUS OF CHILLI RESEARCH IN MALAYSIA	66
3.	MATE	ERIALS AND METHODS	69
) .	MAIL	ERIALS AND METHODS	0,
	3.1.	PLANT MATERIALS	69
	3.2.	MULTILOCATIONAL TRIALS	72
	3.3.	DATA COLLECTION	75
	3.3.	DATA COLLECTION	,,
	3.4.	METEOROLOGICAL DATA	77
	3.5.	DRYING METHOD	85
	3.6.	STATISTICAL ANALYSIS	86
	3.0.		86
			86
		5.0.2 G 1 B 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	86
			93
			93
			94
		5.0.5 COMMEDITION LENGTHS CHARACTER	94
		5.0.0 CROSS O VERTITIES TO THE	94
		5.0.7 STABLETT THE EVENTER OF THE PERSON	94
		3.6.7.1. Genotype - grouping using mean and coefficient of variance (CV)	95
			95 96
			90 98
		3.1.1.3. Kank - sum index	70
1 .	RESU	ILTS	100
	4.1.	110000000000000000000000000000000000000	100
			100
		4 1 2 ACCAN DEDECORATANCE	10

		 a) Yield, yield component and fruit 	
		characters	102
		b) Agronomic characters	10
		c) Quality factor	108
	4.1.3.	CORRELATION STUDIES	113
1.2.	MULT	I-LOCATION TRIALS	118
	4.2.1.	CHARACTERISTICS OF THE	
		ENVIRONMENT	118
	4.2.2.	DRYING METHOD	120
	4.2.3.	PERFORMANCE OF THE G x E TRIALS	122
		4.2.3.1. ANOVA by environment	122
		4.2.3.2. Mean performance of genotype and	
		environment	126
		a) Yield and yield component	126
		b) Agronomic characters	130
		c) Quality characters	133
		4.2.3.3. Combined analysis of variance	148
		4.2.3.3.1. Combined ANOVA over	1-10
		macro-environments	151
		4.2.3.3.2. Combined ANOVA over locations	131
		and planting-seasons	153
			155
		4.2.3.4.1. Variance components over macro-	15.
			155
		environments	133
		4.2.3.4.2. Variance components over locations	157
		and planting-seasons	13
		4.2.3.5. Genotypic coefficients of variation and	
		heritability estimates	160
		4.2.3.6. Correlation studies among characters	
		in different environments	164
		a) Relationship between yield and other	
		traits	164
		b) Relationship between yield	
		components and other traits	166
		c) Relationship between quality	
		characters and other traits	167
4.3.	FSTIN	MATE OF STABILITY PARAMETERS	180
	4.3.1.	GENOTYPE GROUPING BASED	
	1.5.1.	ON MEAN AND CV OF GENOTYPES	18
		4.3.1.1. Yield and yield components	18
		4.3.1.2. Agronomic characters	183
		4.3.1.3. Quality characters	184
		4.5.1.5. Quality characters	13-
	4.3.2.	REGRESSION AND DEVIATION	198
		4.3.2.1. Yield and yield components	199
		4 3 2 2 Agronomic characters	200

		4.3.2.3. Quality characters	201
		EFFECTS	206
		4.3.3.1. Yield and yield components	206
		4.3.3.2. Agronomic characters	207
		4.3.3.3. Quality characters	208
		4.3.4. RANKING METHODS	221
		4.3.4.1. Yield and yield components	221
		4.3.4.2. Agronomic characters	222
		4.3.4.3. Quality characters	223
5.	DISC	USSION	226
		D.TTD ODLIGTION	230
	5.1.	INTRODUCTION	230
	5.2.	GENETIC X ENVIRONMENTAL STUDIES	231
		5.2.1. YIELD AND COMPONENTS	231
		5.2.2. OTHER AGRONOMIC CHARACTERS	238
		5.2.3. QUALITY DETERMINING CHARACTERS	240
	5.3.	THE RELATIVE IMPORTANCE OF GENOTYPE,	
	0.0.	ENVIRONMENTAL AND GENOTYPE-	
		ENVIRONMENTAL EFFECTS	243
		5.3.1. GENOTYPIC EFFECTS	243
		5.3.2. ENVIRONMENTAL EFFECTS	244
		5.3.3. GENOTYPIC AND ENVIRONMENTAL	
		INTERACTION	245
		INTERACTION	243
	5.4.	HERITABILITY AND GENETIC COEFFICIENT OF	
		VARIATION	246
	5.5.	CORRELATION	247
	3.3.	CONCENTION	
	5.6.	STABILITY ANALYSIS	252
		5.6.1. STABILITY BY CHARACTERS	253
		5.6.2. STABILITY BY VARIETIES	258
		5.6.3. SIMULTANEOUS SELECTION OF	
		VARIETIES	266
		5.6.4. DELINEATING SUITABLE AREAS FOR CHILL	.I
		PRODUCTION	272
	5.7.	CONCLUDING REMARKS	273
6.	SUM	MARY	283
7.	REFE	RENCES	287

LIST OF TABLES

Tables		Pg
1.1.	Chilli consumption in Malaysia (tons)	5
1.2.	Areas of chilli cultivation in Peninsular Malaysia (ha)	6
1.3.	Statistics of chilli imports in Malaysia	8
2.1.	Comparative pungency ratings for chillies from different geographical regions	17
2.2.	Chemical compositions of chilli (Capsicum annuum L.)	20
2.3.	Commercial uses of chilli according to pungency level	22
3.1.	Origins and characteristics of 64 varieties of chilli	70
3.2.	Meteorological data and soil conditions of selected sites for G x E study	74
3.3.	Analysis of variance and expected mean squares for the analysis of Method A: over macro-environments with planting-season and location effects combined	88
3.4.	Analysis of variance and expected mean squares for the analysis of Method B: over planting-season and location effects separated	90
3.5.	Partition of G x E interaction for each genotype into regression and deviation components	97
4.1.	Mean square values (df. within parenthesis) for the various traits of chilli in the preliminary evaluation trials	101
4.2.	Mean values of some chilli characteristics measured in the preliminary trial evaluated in Jalan Kebun	110
4.3.	Simple correlation coefficients among various chilli characteristics measured in the preliminary evaluation	117

4.4.	The air temperature (*C), the relative humidity, evaporation rate, solar radiation, sunshine and the rainfall distribution during the crop cycle	119
4.5.	Mean squares for yield/plant, no. of fruits/plant, mean of fruit weight and dry yield of 22 genotypes for each of the 14 environments	123
4.6.	Mean squares for days to harvest, days to dry, dry weight, conversion rate and plant height of 22 genotypes for each of the 14 environments	124
4.7.	Mean squares for % of bleaching, light transmission and capsaicin content of 22 genotypes for each of the 14 environments	125
4.8.	Mean values of yield (g/plant) for genotypes grown in different environments	137
4.9.	Mean values of number of fruits/plant for genotypes grown in different environments	138
4.10.	Mean values of mean fruit weight (g/fruit) for genotypes grown in different environments	139
4.11.	Mean values of dry yield (g/plant) for genotypes grown in different environments	140
4.12.	Mean values of days to harvest for genotypes grown in different environments	141
4.13.	Mean values of plant height (cm) for genotypes grown in different environments	142
4.14.	Mean values of days to dry for genotypes grown in different environments	143
4.15.	Mean values of conversion rate (solid content as percentage of fresh) for genotypes grown in different environments	144
4.16.	Mean values of percentage of bleaching for genotypes grown in different environments	145

4.17.	grown in different environments	146
4.18.	Mean values of capsaicin for genotypes grown in different environments	147
4.19.	Bartlett's test for homogeneity of error variance over over 14 environments	150
4.20(a).	Mean squares values for agronomic and quality character in a combined analysis over 14 environments	152
4.20(b).	Mean squares values for combined ANOVA over locations and planting-seasons for some chilli traits	154
4.21(a).	Variance components and standard error in parenthesis for some chilli characters over different macro-environments	156
4.21(b).	Variance components and standard error of the estimates in parenthesis for each parameter over locations and planting-seasons	159
4.22(a).	Genotypic and phenotypic variance, genotypic coefficients of variation and heritability estimates for 12 characters from ANOVA over macro-environments	162
4.22(b).	Genotypic and phenotypic variance, genotypic coefficients of variation and heritability estimates for 12 characters from ANOVA over locations and planting-seasons	163
4.23.1.	Correlation coefficients between yield (g/plant) against other traits by environments	168
4.23.2.	Correlation coefficients between mean fruit weight (g/fruit) against other traits by environments	169
4.23.3.	Correlation coefficients between number of fruits/plant against other traits by environments	170
4.23.4.	Correlation coefficients between dry yield and other traits by environments	171
4.23.5.	Correlation coefficients between plant height and other traits by environments	172

4.23.6.	Correlation coefficients between days to harvest and other traits by environments	173
4.23.7.	Correlation coefficients between days to dry and other traits by environments	174
4.23.8.	Correlation coefficients between dry weight against other traits by environments	175
4.23.9.	Correlation coefficients between conversion rates against other traits by environments	176
4.23.10.	Correlation coefficients between percentage of bleaching against other traits by environments	177
4.23.11	Correlation coefficients between light transmission and other traits by environments	178
4.23.12	Correlation coefficients between capsaicin content and other traits by environments	179
4.24.1.	Estimates of regression coefficients and stability variance for yield and yield components in chilli	203
4.24.2	Estimates of regression coefficients and stability variance for some agronomic characters in chilli	204
4.24.3.	Estimates of regression coefficients and stability variance for some quality characters in chilli	205
4.25.1.	Ranking and stability rating (according to Kang 1991a) for yield and yield components of chilli	225
4.25.2.	Ranking and stability rating (according to Kang 1991a) for dry yield of chilli	226
4.26.1.	Ranking and stability rating (according to Kang 1991a) for plant height and days to harvest of chilli	227
4.26.2.	Ranking and stability rating (according to Kang 1991a) For days to dry, dry weight and conversion rates of chilli	228
4.27.	Ranking and stability rating (according to Kang 1991a) for product colour, percentage of bleaching and capsaicin content of chilli	229

5.1.	Stability estimates of 22 varieties of chilli with respect to yields, some agronomic and quality characters	265
5.2.	Stability estimates for yield (g/plant) and overall scores of 22 varieties of hot pepper varieties evaluated at 14 environments	271

LIST OF FIGURES

Figures		Pg
1.1.	Distribution of chilli cultivation in Peninsular Malaysia	7
3.1.	Location of the trial sites	73
3.2.1.	Distribution of temperature, rainfall and sunshine at Cameron Highlands	78
3.2.2.	Distribution of temperature, rainfall and sunshine at Bertam	79
3.2.3.	Distribution of temperature, rainfall and sunshine at Gajah Mati	80
3.2.4.	Distribution of temperature, rainfall and sunshine at Jalan Kebun	81
3.2.5.	Distribution of temperature, rainfall and sunshine at Kuala Linggi	82
3.2.6.	Distribution of temperature, rainfall and sunshine at Telong	83
3.2.7.	Distribution of temperature, rainfall and sunshine at Kundang	84
4.1.1.	Distribution of means of yield per plant and coefficients of variance of 22 varieties of chilli	186
4.1.2.	Distribution of means of mean fruit weight (g/fruit) coefficients of variance of 22 varieties of chilli	187
4.1.3.	Distribution of means of no. of fruits/plant and coefficients of variance of 22 varieties of chilli	188
4.1.4.	Distribution of means of dry yield and coefficients of variance of 22 varieties of chilli	189
4.1.5.	Distribution of means of days to harvest and coefficients	190

LIST OF FIGURES

Figures		Pg
1.1.	Distribution of chilli cultivation in Peninsular Malaysia	7
3.1.	Location of the trial sites	73
3.2.1.	Distribution of temperature, rainfall and sunshine at Cameron Highlands	78
3.2.2.	Distribution of temperature, rainfall and sunshine at Bertam	79
3.2.3.	Distribution of temperature, rainfall and sunshine at Gajah Mati	80
3.2.4.	Distribution of temperature, rainfall and sunshine at Jalan Kebun	81
3.2.5.	Distribution of temperature, rainfall and sunshine at Kuala Linggi	82
3.2.6.	Distribution of temperature, rainfall and sunshine at Telong	83
3.2.7.	Distribution of temperature, rainfall and sunshine at Kundang	84
4.1.1.	Distribution of means of yield per plant and coefficients of variance of 22 varieties of chilli	186
4.1.2.	Distribution of means of mean fruit weight (g/fruit) coefficients of variance of 22 varieties of chilli	187
4.1.3.	Distribution of means of no. of fruits/plant and coefficients of variance of 22 varieties of chilli	188
4.1.4.	Distribution of means of dry yield and coefficients of variance of 22 varieties of chilli	189
4.1.5.	Distribution of means of days to harvest and coefficients	190

4.1.6.	Distribution of means of plant height and coefficients of variance of 22 varieties of chilli	191
4.1.7.	Distribution of means of days to dry and coefficients of variance of 22 varieties of chilli	192
4.1.8.	Distribution of means of dry weight (g) and coefficients of variance of 22 varieties of chilli	193
4.1.9.	Distribution of means of conversion rate (%) and coefficients of variance of 22 varieties of chilli	194
4.1.10.	Distribution of means of percentage of bleaching and coefficients of variance of 22 varieties of chilli	195
4.1.11.	Distribution of means of light transmission and coefficients of variance of 22 varieties of chilli	196
4.1.12.	Distribution of means of capsaicin content (mg/g) and coefficients of variance of 22 varieties of chilli	197
4.2.1.	Plot of regression lines of individual genotypes against environmental index for yield	209
4.2.2.	Plot of regression lines of individual genotypes against environmental index for mean fruit weight	210
4.2.3.	Plot of regression lines of individual genotypes against environmental index for no. of fruits/plant	211
4.2.4.	Plot of regression lines of individual genotypes against environmental index for dry yield	212
4.2.5.	Plot of regression lines of individual genotypes against environmental index for days to harvest	213
4.2.6.	Plot of regression lines of individual genotypes against environmental index for plant height	214
4.2.7.	Plot of regression lines of individual genotypes against environmental index for days to dry	215
4.2.8.	Plot of regression lines of individual genotypes against	216

4.2.9.	Plot of regression lines of individual genotypes against environmental index for conversion rates	217
4.2.10.	Plot of regression lines of individual genotypes against environmental index for percentage of bleaching	218
4.2.11.	Plot of regression lines of individual genotypes against environmental index for light transmission at 490 nm	219
4.2.12.	Plot of regression lines of individual genotypes against environmental index for capsaicin content	220
5.1.	Yield as influenced by climatic factors (rainfall, sunshine, temperature and humidity) during growing period	278

LIST OF PLATES

Plates		pg
1.	Modified conventional dryer. Left: Fresh chilli placed on plastic insulated wire netting	
	ready to be solar dried.	
	Right: Solar dried chilli still inside the drier	
	Bottom: Modified solar dryer: iron angled frame with	
	Plastic coated wire netting as the base.	
	Detachable cover made of UV treated	
	plastic (see inset)	121
2.	Variety Brebes (V44): A medium size plant, above	
	average yielder with good quality attribute and most stable	
	variety with stability in twelve characters	279
3.	Variety purple chilli (V16) selected for characters:	
	days to dry (stable), pungency (stable), fruit number (stable),	
	conversion rate, good colour (stable), high colour	
	retention (stable) on drying and yield (stable).	
	Stability in 9 characters.	280
4.	Variety MC 11 selected for good fruit size(stable), optimum	
	Height (stable), high yield (stable), pungency(stable), conversion	
	rate (stable), no of fruits/plant(stable), days to dry(stable).	
	Stability in 11 characters. MC11 is a double purpose chilli-	
	The glossy bright red (solar dried) and very intent red	
	colour (oven dried) of MC11	281
5.	Variety Ch388 (V57) selected for characters: yield (stable),	
	pungency (stable), fruit number (stable), conversion rate(stable)	
	and long attractive fruits (stable) suitable for fresh consumption	
	and as raw material. Stability in 11 characters	282

LIST OF APPENDICES

Appendices		pg
Appendix I:	Typical growth habit of chilli plant as in IPBGR descriptor list	310
Appendix II:	Temperatures (°C) pattern under solar dryer against air temperatures at various time of the day (31.1.90 - 17.2.90)	311