

ACU 7094



**DIVERSITY AND ABUNDANCE OF PHYTOPLANKTON  
IN TASEK BERA, PAHANG**

TANG SOO HUI

INSTITUTE OF POSTGRADUATE STUDIES AND RESEARCH  
UNIVERSITY OF MALAYA  
KUALA LUMPUR

2001

Perpustakaan Universiti Malaya



A510668932

0K

**DIVERSITY AND ABUNDANCE OF PHYTOPLANKTON  
IN TASEK BERA, PAHANG**

Tang Soo Hui

Dissertation submitted to  
the Institute of Postgraduate Studies and Research,  
University of Malaya, Kuala Lumpur,  
in partial fulfilment of the requirements for the  
Degree of Master of Technology (Environmental Management)

April 2001

## ABSTRACT

Tasek Bera, located in southwest of Pahang state, is Malaysia's largest freshwater swamp and lake system. It is one of the remaining examples of a lowland alluvial riparian swamp system. Tasek Bera was designated as Malaysia's first "Wetlands of International Importance" under the Ramsar Convention in November 1994. The water quality of Tasek Bera was monitored from April to October 1998 to assess environmental conditions affecting phytoplankton assemblages. Measurement of physical and chemical parameters at selected sites ranged as follows : pH (4.45 to 6.73), temperature (25.3 to 33.3 °C), dissolved oxygen (0.40 to 5.80 mgL<sup>-1</sup>), water depth (0.15 to 5.40 m), Secchi depth (0.14 to 2.19 m), conductivity (11.20 to 129.60 μS.cm<sup>-1</sup>), dissolved orthophosphate (0.0000 to 0.0095 mgL<sup>-1</sup>), nitrate (0.000 to 5.600 mgL<sup>-1</sup>) and ammoniacal-N (0.000 to 2.9132 mgL<sup>-1</sup>). In general, most of the sites are relatively undisturbed. However, nitrate, ammoniacal-N, conductivity and turbidity were quite high at some of the sites.

A total of 326 species of phytoplankton, including 42 unidentified species, were found in the study. This consisted of 150 species of Chlorophyta, 90 species of Chrysophyta, 36 species of Euglenophyta, three species of Pyrrhophyta, three species of Xanthophyta and two species of Cyanophyta. Chlorophyll-a concentration ranged from 0.000 to 7.3437 mgm<sup>-3</sup> while primary productivity was measured at a maximum of 0.8573 mgO<sub>2</sub><sup>-1</sup>l<sup>-1</sup>hour<sup>-1</sup>. Cell counts ranged from 1.34 to 400.275 cells.L<sup>-1</sup>.

In the two-way analysis of variance, primary productivity and transparency were significantly different among sampling stations. Chlorophyll-a levels, Shannon-Weiner Diversity Index ( $H'$ ) values, Margalef's Species Index ( $d$ ) values, Species Evenness ( $J$ ) values, phytoplankton cell counts, conductivity and pH were significantly different among sampling occasions. Multiple regression analysis showed that pH and nitrate contributed significantly to chlorophyll-a concentration. Conductivity contributed significantly to Shannon-Weiner Diversity Index ( $H'$ ) values, Margalef's Species Index ( $d$ ) values, Species Evenness ( $J$ ) values and phytoplankton cell counts. Water depth, transparency and temperature contributed significantly to primary productivity. Cluster analysis was used to segregate sampling stations based on biotic parameters as well as environmental parameters.

Pollutant-tolerant species identified were *Trachelomonas* sp., *Gomphonema parvulum*, *Navicula* sp., *Peridinium* sp. and *Euglena acus*. This research looks into the feasibility of using phytoplankton assemblages as indicators of environmental change due to anthropogenic effects, particularly of agricultural activities.

## ABSTRAK

Tasek Bera, yang terletak di barat-daya negeri Pahang, merupakan paya air tawar serta sistem tasik yang terbesar di Malaysia. Ia merupakan satu-satunya daripada contoh paya riparian aluvial tanah rendah yang tinggal. Tasek Bera diisytiharkan sebagai tanah lembab berkepentingan antarabangsa yang pertama di Malaysia di bawah Konvensyen Ramsar dalam November 1994. Kualiti air Tasek Bera dikaji dari April hingga Oktober 1998 untuk menilai bagaimana komposisi fitoplankton dipengaruhi oleh keadaan persekitaran.

Ukuran bagi parameter fizikal dan kimia di stesyen-stesyen tertentu adalah seperti berikut : pH (4.45 to 6.73), suhu (25.3 to 33.3 °C), oksigen terlarut (0.40 to 5.80 mgL<sup>-1</sup>), kedalaman air (0.15 to 5.40 m), kedalaman Secchi (0.14 to 2.19 m), konduktiviti (11.20 to 129.60 μS.cm<sup>-1</sup>), ortofosfat terlarut (0.0000 to 0.0095 mgL<sup>-1</sup>), nitrat (0.000 to 5.600 mgL<sup>-1</sup>) dan ammoniakal-N (0.000 to 2.9132 mgL<sup>-1</sup>). Pada amnya, kebanyakan stesyen kurang mengalami gangguan secara relatif. Namun, nitrat, ammoniakal-N, konduktiviti dan tubiditi di sesetengah stesyen didapati agak tinggi.

Sebanyak 326 spesies phytoplankton, termasuk 42 spesies yang tidak dikenalpasti telah dijumpai dalam kajian ini. Ia merangkumi 150 spesies Chlorophyta, 90 spesies Chrysophyta, 36 spesies Euglenophyta, tiga spesies Pyrrhophyta, tiga spesies Xanthophyta serta dua spesies Cyanophyta. Kepekatan klorofil-a berjulat antara 0.000 hingga 7.3437 mgm<sup>-3</sup> manakala produktiviti primer mempunyai nilai maksimum 0.8573 mgO<sub>2</sub><sup>-1</sup>jam<sup>-1</sup>. Kiraan sel fitoplankton berjulat antara 1.34 hingga 400.275 sel.L<sup>-1</sup>.

Dalam ujian ANOVA dua hala, produktiviti primer dan ketelusan air didapati berbeza secara signifikan di kalangan stesyen-stesyen. Kepekatan klorofil-a, nilai Indeks Diversiti Shannon-Weiner ( $H'$ ), nilai Indeks Spesies Margalef ( $d$ ), nilai Indeks Kesamarataan Spesies ( $J$ ), kiraan sel fitoplankton, konduktiviti dan pH didapati berbeza secara signifikan di antara masa penyampelan. Analisis 'multiple regression' menunjukkan bahawa pH serta nitrat menyumbang secara signifikan terhadap kepekatan klorofil-a. Konduktiviti menyumbang secara signifikan terhadap nilai Indeks Diversiti Shannon-Weiner ( $H'$ ), nilai Indeks Species Margalef ( $d$ ), nilai Indeks Kesamarataan Spesies ( $J$ ) serta kiraan sel fitoplankton. Kedalaman air, ketelusan air serta suhu menyumbang secara signifikan terhadap produktiviti primer. Analisis kluster digunakan untuk mengasingkan stesyen penyampelan berdasarkan parameter biotic dan persekitaran.

Spesies toleran terhadap pencemaran yang telah dikenalpasti termasuklah *Trachelomonas* sp., *Gomphonema parvulum*, *Navicula* sp., *Peridinium* sp. and *Euglena acus*. Kajian untuk menilai kesesuaian menggunakan komposisi fitoplankton sebagai penunjuk kepada perubahan dalam keadaan persekitaran akibat kesan antropogenik, terutamanya aktiviti pertanian.

## ACKNOWLEDGEMENTS

I wish to express my sincere gratitude to my supervisor, Professor Phang Siew Moi for her guidance, suggestions and encouragement. I would like to thank Associate Professor Dr. Abdul Rani Abdullah for his advice and helpful comments.

Special thanks to Wetlands International-Asia Pacific for providing the Masters' student grant that made this project possible. I wish to express immense gratitude to Lead Technical Officer, Crawford Prentice, Technical Officers, Alvin Lopez and TL Murugadas of Wetlands International, who have provided tremendous support in facilitating the field trips. My profound appreciation also goes to a wonderful Semelai friend, Stem, for his patience and cheerful disposition despite the laborious task of navigating around the sites.

I thank Dr. Chu Wan Loy for his guidance in the identification of phytoplankton, En. Wan Yusoff as well as my course mates Kahoru Takushi and Jillian Ooi and my sister, Julianna, for their assistance during the field trips. I also thank Kok Wai Hong for his guidance in laboratory analysis, Associate Professor Chong Ving Ching and A. Baset AI for helping in the statistical analysis. I appreciate the constant support from my laboratory mates, Melor Ismail, Gan Sook Yee, Lim Phaik Eem, Miah Shohidullah, Wong Ching Lee, Edmund Oh, Rohani Ahmad and Quamrul Ahsan, Miss Molly Chuah and her staff in the IPSP library as well as my supervisor at work in World Wide Fund for Nature (WWF) Malaysia, Mr. Balu Perumal, who have been very understanding and helpful.

I wish to dedicate this dissertation to both my parents who have always been there for me during the trying times and challenging moments of the project.

## TABLE OF CONTENTS

	Page
<b>ABSTRACT</b>	I
<b>ABSTRAK</b>	iii
<b>ACKNOWLEDGEMENTS</b>	v
<b>TABLE OF CONTENTS</b>	vi
<b>LIST OF TABLES</b>	xiii
<b>LIST OF FIGURES</b>	xiv
<b>LIST OF ABBREVIATIONS</b>	xxi
<b>SYMBOLS AND UNITS</b>	xxii
<b>1.0 INTRODUCTION</b>	1
1.1 LOCATION OF TASEK BERA	1
1.2 BACKGROUND OF TASEK BERA	1
1.3 TASEK BERA AS A RAMSAR SITE	3
1.4 INTERNATIONAL IMPORTANCE OF TASEK BERA RAMSAR SITE	3
1.5 RESEARCH BY THE INTERNATIONAL BIOLOGICAL PROGRAMME (IBP)	4
1.6 DEVELOPMENT AND HUMAN ACTIVITIES IN THE WATER CATCHMENT AREA	5
1.7 OBJECTIVES OF THE RESEARCH	6
<b>2.0 LITERATURE REVIEW</b>	7
2.1 TASEK BERA ECOSYSTEM AND BIOLOGICAL DIVERSITY	7
2.2 ENVIRONMENTAL CONDITIONS OF TASEK BERA	8
2.2.1 Rainfall	8



2.2.2	Physical properties	9
2.2.2.1	Transparency	9
2.2.2.2	Temperature	10
2.2.2.3	Water level fluctuations	11
2.2.3	Chemical properties	11
2.2.3.1	Dissolved oxygen and carbon dioxide	11
2.2.3.2	pH	12
2.2.3.3	Conductivity	13
2.2.3.4	Phosphate and nitrogen	15
2.2.4	Primary production	17
2.2.4.1	Classification of algal primary producers	17
2.2.4.2	Population density and standing crop	21
2.2.4.3	The chlorophyll content of phytoplankton	22
2.2.4.4	Photosynthetic production	22
2.3	THE DISTRIBUTION PATTERNS OF FRESHWATER PHYTOPLANKTON	23
2.3.1	Spatial distribution	23
2.3.1.1	Vertical distribution	23
2.3.1.2	Horizontal distribution	24
2.3.2	Temporal distribution	25
2.4	FACTORS CONTROLLING THE GROWTH OF PHYTOPLANKTON	25
2.4.1	Physical factors	25
2.4.1.1	Temperature	25
2.4.1.2	Light	26
2.4.1.3	Dissolved oxygen	27
2.4.1.4	Carbon dioxide	27

2.4.1.5 pH	28
2.4.2 Photosynthetic activities	28
2.4.3 Nutrient requirements	31
2.4.3.1 Nutrient uptake and growth rates of phytoplankton	32
2.4.3.2 Phosphorus availability	32
2.4.3.3 Nitrogen availability	33
2.4.3.4 The cycle of phosphorus and nitrogen in freshwater wetlands	35
2.5 PERIODICITY AND SEASONAL CHANGES IN PHYTOPLANKTON COMPOSITION	37
2.5.1 Seasonal succession	38
2.5.2 Adaptive characteristics and class succession patterns	39
2.5.3 Long term compositional response	48
2.6 EFFECTS OF HUMAN ACTIVITIES AND CHANGES OF LAND USE ON THE COMPOSITION OF STREAMS	51
2.6.1 Agriculture and water quality	54
2.6.2 Integrated agricultural practices in palm oil and / or rubber plantations	56
2.7 EUTROPHICATION	61
2.7.1 Natural and anthropogenic eutrophication	61
2.7.2 Eutrophication Indices and their relation to other indices of ecosystem change	63

2.8 USE OF ALGAE FOR MONITORING RIVERS	66
2.8.1 Principles and aims of monitoring	66
2.8.2 Methodological aspects and perspectives of bio-monitoring	69
2.8.3 Problems of sampling river algae for monitoring	75
2.9 MANAGEMENT POTENTIAL OF FRESHWATER WETLANDS	76
<b>3.0 MATERIALS AND METHODS</b>	<b>77</b>
3.1 THE SAMPLING SITES	77
3.2 Field measurements and sampling	77
3.2.1 Physical and chemical parameters	85
3.2.2 Productivity measurements	85
3.2.3 Phytoplankton sampling	86
3.3 LABORATORY MEASUREMENTS	87
3.3.1 Chlorophyll-a measurements	87
3.3.2 Chemical analysis	88
3.3.2.1 Dissolved orthophosphate	88
3.3.2.2 Ammoniacal-nitrogen	89
3.3.2.3 Nitrate	90
3.3.3 Microscopic identification and quantification of phytoplankton	91
3.3.3.1 Microscope calibration	91
3.3.3.2 Phytoplankton counting techniques	92
3.4 DATA ANALYSIS	94
3.4.1 Frequency	94
3.4.2 Diversity studies	95
3.4.3 Statistical analysis	96

<b>4.0 RESULTS</b>	98
4.1 FIELD MEASUREMENTS	98
4.1.1 Physical and chemical parameters	98
4.1.1.1 Water depth	98
4.1.1.2 Transparency	98
4.1.1.3 Conductivity	101
4.1.1.4 pH	101
4.1.1.5 Temperature	101
4.1.1.6 Dissolved oxygen	101
4.1.2 Productivity measurements	105
4.1.3 Rainfall data in the catchment area	114
4.1.4 Fertiliser application data from various plantations in the catchment area	114
4.2 LABORATORY MEASUREMENTS	119
4.2.1 Chlorophyll-a measurements	119
4.2.2 Chemical analysis	126
4.2.2.1 Dissolved orthophosphate	126
4.2.2.2 Ammoniacal-nitrogen	126
4.2.2.3 Nitrate	138
4.2.3 Microscopic identification and quantification of phytoplankton	144
4.2.3.1 Checklist of phytoplankton and abundance	144
4.2.3.2 Frequency (%) of phytoplankton	166
4.2.3.3 Diversity Indices	166
4.2.3.4 Sorensen's Similarity Coefficient	193

4.3 DATA ANALYSIS	196
4.3.1 Statistical analysis	196
4.3.1.1 Two-way ANOVA	196
4.3.3.2 Multiple regression	198
4.3.3.3 Cluster analysis	200
<b>5.0 DISCUSSION</b>	205
5.1 ENVIRONMENTAL QUALITY OF TASEK BERA	205
5.1.1 Water depth and transparency	205
5.1.2 Conductivity	206
5.1.3 pH	207
5.1.4 Temperature	207
5.1.5 Dissolved oxygen	208
5.1.6 Dissolved orthophosphate	208
5.1.7 Ammoniacal-N	210
5.1.8 Nitrate	211
5.1.9 Rainfall in the catchment area	212
5.2 PRIMARY PRODUCTIVITY IN TASEK BERA	212
5.2.1 Chlorophyll-a measurements	212
5.2.2 Photosynthetic productivity	213
5.3 PHYTOPLANKTON DIVERSITY	214
5.3.1 Checklist of phytoplankton and abundance	214
5.3.2 Sorensen's Similarity Coefficient	216
5.3.3 Diversity Indices	217
5.3.4 Dominance	220
5.3.5 Trophic classes	220

5.4 EFFECT OF THE ENVIRONMENTAL PARAMETERS ON PHYTOPLANKTON	226
5.4.1 Statistical analysis	227
5.4.2 Biological indicators	229
5.5 SOURCES OF DEGRADATION AND MANAGEMENT RECOMMENDATIONS	231
5.5.1 Human Induced Factors Influencing Water Quality	233
5.5.1.1 Modification of catchment area	233
5.5.1.2 Pollution (point source)	235
5.5.1.3 Fires	235
5.5.2 Legislation, Administration and Management of Tasek Bera	237
5.5.3 Zonation for Site Management	243
<b>6.0 CONCLUSION</b>	247
<b>REFERENCES</b>	253
<b>APPENDIXES</b>	

<b>List of Tables</b>		Page
Table 1	Number of freshwater algal species identified in Malaysia	20
Table 2	Algal genus pollution index	42
Table 3	Algal species pollution index	43
Table 4	Pollutant-tolerant species of algae. List of the 80 most tolerant species, in order of decreasing emphasis by 165 authorities.	44
Table 5	Characteristics of common major algal associations of the phytoplankton in relation to increasing lake fertility (Wetzel, 1983).	49
Table 6	Diatom species tolerant, indifferent and intolerant to rubber affluent (Nather Khan <i>et al.</i> , 1986)	52
Table 7	Clean water and polluted water species of diatoms from Sungai Bertam, Cameron Highlands (Anton and Wan Othman, 1990)	53
Table 8	Chemical composition and equivalence to fertilisers in 1,000 tonnes of POME	60
Table 9	The saprobic system and physicochemical parameters (Hynes, 1960; 1971)	67
Table 10	Summary of the statistical analysis	97
Table 11	Phytoplankton abundance (cells.L <sup>-1</sup> ) at sampling stations 1 to 12 in Tasek Bera (April – October 1998)	145
Table 12	Frequency (%) of phytoplankton for sampling station 1 to 12 in Tasek Bera (April – October 1998)	167
Table 13	Sorensen's Similarity Coefficient (%) for phytoplankton diversity based on different studies in Tasek Bera.	197

Table 14	Results of the multiple regression analysis showing the important parameters contributing to the biotic parameters observed at Tasek Bera (April – October 1998)	199
Table 15	Distribution (number of species) of phytoplankton by division for Stations 1 to 12.	221
Table 16	Taxonomic composition (number of species) of phytoplankton from April – September 1998	222
Table 17	Trophic classes by chlorophyll-a and algal numbers according to Felfody (1987).	223
Table 18	Trophic class for all stations by chlorophyll-a (Felfody, 1987).	224
Table 19	Trophic class for all stations by algal numbers (Felfody, 1987).	225
Table 20	Tasek Bera Ramsar Site Management Authority Organisation Chart	242

### List of Figures

Figure 1	Location of Tasek Bera	2
Figure 2	Pollution tolerant phytoplankton species : <i>Nitzschia palea</i>	41
Figure 3	Pollution tolerant phytoplankton species : <i>Scenedesmus quadricauda</i>	41
Figure 4	Pollution tolerant phytoplankton species : <i>Trachelomonas</i> sp.	45
Figure 5	Pollution tolerant phytoplankton species : <i>Gomphonema parvulum</i>	45
Figure 6	Pollution tolerant phytoplankton species : <i>Navicula</i> sp.	46
Figure 7	Pollution tolerant phytoplankton species : <i>Peridinium</i> sp.	46



Figure 8	Pollution tolerant phytoplankton species: <i>Euglena acus</i>	47
Figure 9	Location of sampling stations and Felda plantations at Tasek Bera	78
Figure 10	Sampling Station 1 : Tanjung Kuin	79
Figure 11	Sampling Station 2 : Sungai Tasik	79
Figure 12	Sampling Station 3 : Lubuk Pathir	80
Figure 13	Sampling Station 4 : Tanjung Papan	80
Figure 14	Sampling Station 5 : Kampung Benal	81
Figure 15	Sampling Station 6 : Kuala Sungai Tembangan	81
Figure 16	Sampling Station 7 : Lubuk Kuang	82
Figure 17	Sampling Station 8 : Kampung Baapa	82
Figure 18	Sampling Station 9 : Sungai Bera 1	83
Figure 19	Sampling Station 10 : Sungai Bera 2	83
Figure 20	Sampling Station 11 : Paya Kelantong	84
Figure 21	Sampling Station 12 : Sungai Tembangan	84
Figure 22a	Water depth of sampling stations (April – October 1998)	99
Figure 22b	Transparency of sampling stations (April – October 1998)	100
Figure 22c	Conductivity of sampling stations (April – October 1998)	102
Figure 22d	pH of sampling stations (April – October 1998)	103
Figure 22e	Temperature of sampling stations (April – October 1998)	104
Figure 22f	Dissolved oxygen of sampling stations (April – October 1998)	106
Figure 23	Primary productivity by sampling stations (April – October 1998)	107
Figure 24	Primary productivity by month	112

Figure 25a	Rainfall and fertiliser loadings in Tembangau 8 Plantation, Tasik Bera (January – December 1998)	115
Figure 25b	Rainfall and fertiliser loadings in Triang Selatan 2 Plantation, Tasik Bera (January – December 1998)	116
Figure 25c	Rainfall and fertiliser loadings in Triang Selatan 1 Plantation, Tasik Bera (January – December 1998)	117
Figure 25d	Rainfall and fertiliser loadings in Bera Selatan 2 Plantation, Tasik Bera (January – December 1998)	118
Figure 26	Measurement of chlorophyll-a ( $\text{mgm}^{-3}$ ) for sampling stations in Tasik Bera (April – October 1998)	120
Figure 27	Measurement of chlorophyll-a ( $\text{mgm}^{-3}$ ) by month for all stations in Tasik Bera	124
Figure 28	Measurement of dissolved orthophosphate ( $\text{mgL}^{-1}$ ) for sampling stations in Tasik Bera (April – October 1998)	127
Figure 29	Measurement of dissolved orthophosphate by month for all sampling stations in Tasik Bera	130
Figure 30	Measurement of ammoniacal nitrogen ( $\text{mgL}^{-1}$ ) for sampling stations in Tasik Bera (April – October 1998)	133
Figure 31	Measurement of ammoniacal nitrogen by month for all sampling stations in Tasik Bera	136
Figure 32	Measurement of nitrate ( $\text{mgL}^{-1}$ ) for sampling stations in Tasik Bera (April – October 1998)	139
Figure 33	Measurement of nitrate by month for all sampling stations in Tasik Bera	142

Figure 34	Common phytoplankton species found in Tasek Bera : <i>Arthrodesmus octocornis</i>	157
Figure 35	Common phytoplankton species found in Tasek Bera : <i>Cosmarium aspherosporum</i>	157
Figure 36	Common phytoplankton species found in Tasek Bera : <i>Staurastrum tetracerum</i>	158
Figure 37	Common phytoplankton species found in Tasek Bera : <i>Rhizosolenia</i> sp.	158
Figure 38	Common phytoplankton species found in Tasek Bera : <i>Dinobryon sertularia</i>	159
Figure 39	Phytoplankton cell count for sampling stations in Tasek Bera (April – September 1998)	160
Figure 40	Phytoplankton cell count by month from April – September 1998 for all sampling stations in Tasek Bera	164
Figure 41	Shannon-Weiner Diversity Index ( $H'$ ) for phytoplankton in Tasek Bera (April – September 1998)	179
Figure 42	Shannon-Weiner Diversity Index of phytoplankton by in Tasek Bera	183
Figure 43	Margalef's Species Index ( $d$ ) for phytoplankton in Tasek Bera (April – September 98)	185
Figure 44	Margalef's Species Index ( $d$ ) of phytoplankton in Tasek Bera	188
Figure 45	Equitability or Species Evenness ( $J$ ) for phytoplankton in Tasek Bera (April – September 1998)	190
Figure 46	Equitability or Species Evenness ( $J$ ) for phytoplankton in Tasek Bera	194

Figure 47	Cluster analysis based on Sorensen's Index of Similarity for all stations in Tasek Bera (April – September 1998)	201
Figure 48	Cluster analysis based on average biotic parameters for all stations in Tasek Bera (April – September 1998)	202
Figure 49	Cluster analysis based on average environmental parameters for all stations in Tasek Bera (April – October 1998)	203

### List of Appendixes

Appendix 1	Description for the sampling sites
Appendix 2	Measurement of depth of sampling stations (April – October 1998)
Appendix 3	Measurement of transparency of sampling stations (April – October 1998)
Appendix 4	Measurement of conductivity of sampling stations (April – October 1998)
Appendix 5	Measurement of pH of sampling stations (April – October 1998)
Appendix 6	Measurement of temperature of sampling stations (April – October 1998)
Appendix 7	Measurement of dissolved oxygen of sampling stations (April – October 1998)
Appendix 8	Measurement of chlorophyll-a of sampling stations (April – October 1998)
Appendix 9	Measurement of dissolved orthophosphate of sampling stations (April – October 1998)

- Appendix 10 Measurement of ammoniacal nitrogen of sampling stations (April – October 1998)
- Appendix 11 Measurement of nitrate of sampling stations (April – October 1998)
- Appendix 12 Shannon-Weiner Diversity Index ( $H'$ ) for phytoplankton in Tasik Bera (April – September 1998)
- Appendix 13 Margalef's Species Index ( $d$ ) for phytoplankton in Tasik Bera (April – September 1998)
- Appendix 14 Equitability or Species Evenness Index ( $J$ ) for phytoplankton in Tasik Bera (April – September 1998)
- Appendix 15 Sorensen's Quotient of Similarity for sampling stations 1 to 12 in Tasik Bera
- Appendix 16 Phytoplankton Count ( $\text{cells.L}^{-1}$ ) for sampling stations 1 to 12 in Tasik Bera
- Appendix 17 a Productivity of phytoplankton for sampling stations 1 to 12 in Tasik Bera (24 – 26/4/98)
- Appendix 17 b Productivity of phytoplankton for sampling stations 1 to 12 in Tasik Bera (30/5 – 2/6/98)
- Appendix 17 c Productivity of phytoplankton for sampling stations 1 to 12 in Tasik Bera (10 – 13/7/98)
- Appendix 17 d Productivity of phytoplankton for sampling stations 1 to 12 in Tasik Bera (16 – 18/8/98)
- Appendix 17 e Productivity of phytoplankton for sampling stations 1 to 12 in Tasik Bera (19 – 21/9/98)

- Appendix 17 f Productivity of phytoplankton for sampling stations 1 to 12 in Tasik Bera (24 – 26/10/98)
- Appendix 18 Rainfall data (mm) in selected plantations in Tasik Bera (1997 vs 1998)
- Appendix 19 a Fertiliser loading in Tembangau 8 Plantation, Tasik Bera (1998)
- Appendix 19 b Fertiliser loading in Triang Selatan 2 Plantation, Tasik Bera (1998)
- Appendix 19 c Fertiliser loading in Triang Selatan 1 Plantation, Tasik Bera (1998)
- Appendix 19 d Fertiliser loading in Bera Selatan 2 Plantation, Tasik Bera (1998)
- Appendix 20 List of phytoplankton from Tasek Bera (Furtado and Mori, 1982)
- Appendix 21 List of phytoplankton from Tasek Bera (Phang and Murugadas, 1997)
- Appendix 22 List of phytoplankton from Tasek Bera (Williamson, David B., 1998)
- Appendix 23 Results of two-way ANOVA to assess difference in environmental and biotic parameters of all the sampling stations
- Appendix 24 Results of two-way ANOVA to assess difference among sampling occasions (April to October 1998)

## List of Abbreviations

ASEAN – Association of Southeast Asian Nations

BOD<sub>5</sub> – 5-day Biological Oxygen Demand Test

Chll-a – Chlorophyll-a

COD - Chemical Oxygen Demand

CITES – Convention on International Trade in Endangered Species of Wild Fauna and  
Flora

DANCED - Danish Cooperation for Environment and Development

DO – Dissolved oxygen

FASSB – Felda Agricultural Services Sdn. Bhd.

FELDA – Federal Land Development Authority

GPS – Geographical Positioning System

IBP - International Biological Programme

IUCN – The World Conservation Union

LAC – Limits of Acceptable Change

JRP – Jordan Rock Phosphate

MOP – Muriate of Potash

O- PO<sub>4</sub> – Orthophosphate

OFM – Osmotic Flow Modifier

OPL – Oxygen Production Capacity

POME - Palm Oil Mill Effluent

PO<sub>4</sub>-P - Phosphate-phosphorus

SA – Sulphate of Ammonium

UV – Ultra violet

z<sub>eu</sub> - euphotic zone

z<sub>s</sub> - Secchi depth

## Symbols and units

atm – atmospheric pressure

°C - degree Celsius

°Cm<sup>-1</sup> - degree Celsius per metre

cells.L<sup>-1</sup> - cells per litre

d - Margalef's Species Index

H' - Shannon-Weiner Diversity Index

J - Species Evenness Index

km<sup>2</sup> - kilometre square

log<sub>10</sub> - logarithm of base 10

mgO<sub>2</sub>L<sup>-1</sup>hour<sup>-1</sup> - milligram oxygen per litre per hour

mg chl.a.m<sup>-3</sup> - milligram chlorophyll-a per metre cube

mg CL<sup>-1</sup>day<sup>-1</sup> - milligram carbon per litre per day

μL - microlitre

mL - millilitre

L – litre

mM – milimolar

M – Molar

nm - nanometre

μm - micrometre

mm – millimeter

m - metre

μg - microgram

mg – milligram

g - gram

mgL<sup>-1</sup> - milligram per litre

mgm<sup>-3</sup> - milligram per metre cube

mm<sup>2</sup>- milliliter square

mm<sup>3</sup>- millimeter cube

OD<sub>630</sub> - optical density at 630nm

OD<sub>645</sub> - optical density at 645nm

OD<sub>665</sub> - optical density at 665nm



ppm – parts per million

$Q_{10}$  - factor by which the rate increases per 10°C rise in temperature km - kilometre

$\mu\text{g P L}^{-1}$  – microgram phosphorus per litre

$\mu\text{g P}_{\text{tot}} \text{L}^{-1}$  – microgram total phosphorus per litre

$\mu\text{g PO}_4^{3-}\text{P L}^{-1}$  – microgram phosphate-phosphorus per litre

$\mu\text{g N L}^{-1}$  – microgram nitrogen per litre

$\mu\text{g NH}_3\text{-N L}^{-1}$  – microgram ammoniacal nitrogen per litre

$\mu\text{S.cm}^{-1}$  - micro Siemens per centimetre (measurement for conductivity)

$\mu\text{mhos.cm}^{-1}$  - micro Ohms per centimetre (measurement for conductivity)

v/v – volume per volume

w/v – weight per volume