

## LIST OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGMENTS	ix
LIST OF CONTENTS	x
ABBREVIATIONS	xviii
LIST OF FIGURES	xix
LIST OF TABLES	xxii
 <b>CHAPTER ONE : INTRODUCTION</b>	
1.1 : Introduction	1
1.2 : <i>Yarrowia lipolytica</i>	8
1.2.1 : Morphology and life cycle	8
1.2.2 : Biochemical properties	10
1.2.3 : Genetics	13
 1.3 : Genetic application in obtaining high producers of citric acid	14
1.3.1 : Mutation	15
1.3.2 : Cloning in yeast	17
1.3.2.1 : Genetic marker for yeast transformation	17
1.3.2.2 : Selection of enhanced producers of citric acid	18
1.3.2.3 : Yeast cloning vectors	19
1.3.2.4 : Transformation of yeast cells	22
 1.4 : Palm oil- its composition	24
1.4.1 : NBD palm olein	27
1.4.2 : NBD palm olein as the carbon source	29
 1.5 : Citric acid fermentation	31
1.5.1 : Effects of fermentation medium on citric acid production	31
1.5.2 : pH	35
1.5.3 : Aeration and agitation	36
1.5.4 : Temperature and duration of incubations	36
 1.6 : Citric acid production from lipids as the carbon source	37
1.6.1 : Utilisation of lipids	37
1.6.2 : Tricarboxylic acid cycle and citric acid formation	42

1.7	:	The objectives of the project	45
-----	---	-------------------------------	----

## CHAPTER TWO : MATERIALS AND METHODS

2.1	:	<i>Yarrowia lipolytica</i> strains	46
2.2	:	<i>Escherichia coli</i> strains	47
2.3	:	Plasmids	47
2.4	:	Materials	47
2.5	:	Storage and maintenance of bacterial and yeast strains	49
2.6	:	Sterilisation	50
2.7	:	Storage	51
2.8	:	Culture media of <i>Yarrowia lipolytica</i>	51
2.8.1	:	Yeast extract peptone glucose (YEPA) medium	51
2.8.2	:	Yeast extract potassium acetate (YEPA) medium	51
2.8.3	:	Minimal solid medium	51
2.8.4	:	Minimal medium with 0.1 % bromocresol green	52
2.8.5	:	Tricarboxylic acid medium	52
2.8.6	:	Fermentation medium	53
2.8.7	:	Seed medium	54
2.8.8	:	Yeast extract-malt extract medium (YM)	54
2.8.9	:	Restrictive growth medium (RG)	55
2.9	:	Culture media of <i>Escherichia coli</i>	55
2.9.1	:	Luria-Bertani medium (LB)	55
2.9.2	:	LB with ampicillin	55
2.9.3	:	LB with tetracycline	56
2.9.4	:	M9 minimal medium	56
2.9.5	:	M9 minimal medium with 0.01% bromocresol green	56
2.10	:	Solutions	57
2.10.1	:	80% (v/v) Glycerol stock solution	57
2.10.2	:	0.1 M Phosphate buffer, pH 7.0	57

2.10.3 :	0.05 M Phosphate buffer, pH 7.0	57
2.10.4 :	20% (v/v) Sodium thiosulphate solution	57
2.10.5 :	N-methyl-N'-nitro-N-nitrosoguanidine (NTG) stock solution	58
2.10.6 :	3:2 (v/v) Dioxane:ethyl acetate solution	58
2.10.7 :	1 M Sodium hydroxide	58
2.10.8 :	10 M Sodium hydroxide	58
2.10.9 :	0.01 M Calcium chloride	59
2.10.10:	0.1 M Citric acid	59
2.10.11:	1 % (w/v) Citric acid	59
2.10.12:	1 % (w/v) threo-D <sub>s</sub> [+] isocitric acid	59
2.10.13:	0.5 % (w/v) <i>di</i> -Ammonium orthophosphate	59
2.11 :	Lipase assay solutions	60
2.11.1 :	2 % (v/v) Polyvinyl alcohol (PVA) solution	60
2.11.2 :	Foamy emulsion	60
2.11.3 :	1:2 (v/v) Acetone: ethanol	60
2.11.4 :	0.05 M NaOH	61
2.11.5 :	0.1 M NaOH	61
2.11.6 :	5% (w/v) Phenophthalen	61
2.12 :	Protein determination solutions	62
2.12.1 :	Bovine serum albumin (BSA) stock solution	62
2.12.2 :	Solution A	62
2.12.3 :	1% (w/v) Sodium citrate	62
2.12.4 :	Solution B	62
2.12.5 :	Solution C	62
2.12.6 :	Solution D	63
2.12.7 :	0.1 M Tris-HCL, pH 7.8	63
2.13 :	Citrate synthase assay solutions	63
2.13.1 :	3 M Tris-base, pH 8.0	63
2.13.2 :	20 mM tris-HCL	63
2.13.3 :	0.05 M 5, 5'-dithio-bis(2-nitrobenzoic acid) (DTNB)	63
2.13.4 :	0.2 M Oxaloacetate	64
2.13.5 :	100 mM Acetyl-CoA	64
2.14 :	Aconitate hydratase assay solutions	64
2.14.1 :	1 M Potassium phosphate	64
2.14.2 :	1 M <i>di</i> -Potassium phosphate	64
2.14.3 :	0.5 M Phosphate buffer, pH 6.0	65
2.14.4 :	1 mM cis-Aconitic acid	65

2.15 :	Isocitrate lyase assay solutions	65
2.15.1 :	0.5 M Phosphate buffer, pH 6.8	65
2.15.2 :	15 mM Magnesium chloride	65
2.15.3 :	10 mM Phenylhydrazine HCL	65
2.15.4 :	6 mM Cysteine HCL	66
2.15.5 :	0.5 mM Isocitric acid	66
2.16 :	Malate synthase assay solutions	66
2.16.1 :	18 mM Tris-HCL	66
2.16.2 :	15 mM Magnesium chloride	66
2.16.3 :	0.2 $\mu$ M Acetyl-CoA	66
2.16.4 :	1 mM Glyoxylate	67
2.17 :	Isocitrate dehydrogenase assay solutions	67
2.17.1 :	0.5 M Phosphate buffer	67
2.17.2 :	0.1 M Magnesium chloride	67
2.17.3 :	25 mM Diphosphopyridine nucleotide (DPN)	67
2.17.4 :	5 mM Isocitric acid	67
2.18 :	Electrophoresis gel and solutions	68
2.18.1 :	0.7 % Agarose gel	68
2.18.2 :	10X strength TBE buffer	68
2.18.3 :	6X strength Bromocresol blue (BPB) loading dye	68
2.18.4 :	Ethidium bromide	69
2.19 :	Solutions for the isolation of the chromosomal DNA of <i>Yarrowia lipolytica</i>	69
2.19.1 :	0.89 % (w/v) Saline solution	69
2.19.2 :	Solution A (0.2 M Tris-base, pH 10.3)	69
2.19.3 :	Solution B	70
2.19.4 :	0.8 M Sodium chloride	70
2.19.5 :	0.4 M Ethylenediaminetetra-acetic acid, disodium salt (EDTA), pH 8.0	70
2.19.6 :	20 % (w/v) Sodium dodecyl sulphate (SDS)	70
2.19.7 :	0.2 M Tris-HCL, pH 8.5	71
2.19.8 :	Solution C	71
2.19.9 :	1 M Tris-HCL, pH 8.0	71
2.19.10 :	Neutral phenol	71
2.19.11 :	Isoamyl-chloroform (1:24, v/v)	72
2.19.12 :	Buffered Phenol-chloroform (1:1, v/v)	72
2.19.13 :	20 mM Tris-HCL, pH 7.5	72
2.19.14 :	30 mM Sodium chloride	72
2.19.15 :	RNase A (10 mg/ml)	72

2.19.16:	0.02 M Tris-HCL, pH 8.0	73
2.19.17:	0.002 M EDTA disodium salt	73
2.19.18:	Tris-HCL-EDTA (TE) buffer, pH 8.0	73
2.19.19:	3 M Sodium acetate	73
2.20 :	Solutions for plasmid DNA isolation from <i>Escherichia coli</i>	74
2.20.1 :	1 M Glucose	74
2.20.2 :	1M Tris-HCL, pH 8.0	74
2.20.3 :	0.25 M EDTA disodium salt, pH 8.0	74
2.20.4 :	Solution I	75
2.20.5 :	0.4 M Sodium hydroxide	75
2.20.6 :	2 % (w/v) Sodium dodecyl sulphate	75
2.20.7 :	Solution II	75
2.20.8 :	5 M Potassium acetate	76
2.20.9 :	Solution III	76
2.21 :	Solutions for the transformation of <i>Yarrowia lipolytica</i>	76
2.21.1 :	0.1 M Citric acid	76
2.21.2 :	0.1 M Lithium acetate	76
2.21.3 :	Carrier DNA for transformation of <i>Yarrowia lipolytica</i>	77
2.21.4 :	40% PEG 4000 in 0.1 M lithium acetate	77
2.22 :	Citric acid fermentation	77
2.22.1 :	Seed culture preparation	77
2.22.2 :	Fermentation	78
2.22.3 :	Determination of optical density	78
2.22.4 :	Measuring of the pH	78
2.22.5 :	Dry weight	78
2.23 :	Analysis of citric acid	79
2.23.1 :	Samples preparation	79
2.23.2 :	Analysis of citric acid by High Performance Liquid Chromatography (HPLC)	79
2.24 :	Induction of the mutants with high citric acid production	80
2.24.1 :	Culture preparation for mutagenesis	80
2.24.2 :	Mutagenesis of <i>Yarrowia lipolytica</i> M240 (ATCC 8661) with NTG	80
2.24.3 :	Selection of the mutants	82

2.24.4 :	Screening for citric acid over producer	82
2.25 :	Lipase activity	83
2.25.1 :	Fermentation	83
2.25.2 :	Samples preparation for lipase activity	83
2.25.3 :	Lipase activity	83
2.26 :	Tricarboxylic acid cycle enzyme activities	84
2.26.1 :	Fermentation	84
2.26.2 :	Sample preparation for enzyme activities	84
2.26.3 :	Protein preparation for standard curve	85
2.26.4 :	Protein determination	85
2.27 :	Citrate synthase activity	85
2.28 :	Aconitate hydratase activity	86
2.29 :	Isocitrate lyase activity	87
2.30 :	Malate synthase activity	88
2.31 :	Isocitrate dehydrogenase activity	89
2.32 :	Genetic analysis	89
2.32.1 :	Mutagenesis	89
2.32.2 :	Screening of the auxotrophic colonies	90
2.32.3 :	Characterisation of auxotrophic mutants	90
2.32.4 :	Genetic crossing by mass mating	91
2.33 :	Isolation of high molecular weight DNA of <i>Yarrowia lipolytica</i>	92
2.33.1 :	Culture preparation	92
2.33.2 :	Isolation of chromosomal DNA	92
2.34 :	Transformation of <i>Escherichia coli</i>	93
2.34.1 :	Preparation of competent cells	93
2.34.2 :	Transformation experiments	94
2.35 :	Small scale plasmid isolation	94
2.36 :	Large scale preparation of plasmid DNA	95
2.36.1 :	Elution and purification of plasmid from the agarose gel using the GENECLEAN II	96

2.37	:	Large scale preparation of pure covalently closed circular (ccc) DNA from <i>Escherichia coli</i>	97
2.38	:	Electrophoresis of DNA	97
2.39	:	Determination of DNA concentrations	98
2.40	:	Lithium acetate method for the transformation of <i>Yarrowia lipolytica</i>	98
2.40.1	:	Preparation of recipient cells	98
2.40.2	:	Transformation of the yeast cells	99
2.41	:	Digestion of high molecular weight DNA of <i>Yarrowia lipolytica</i> by <i>Sau</i> 3AI and gel electrophoresis	99
2.42	:	Restriction endonuclease digestion of plasmid DNA	100
2.43	:	Ligations of fragments DNA with compatible ends	100
2.44	:	Production of citric acid in continous culture	101

### CHAPTER THREE : RESULTS

3.1	:	Isolation of strains with high level of citric acid production	103
3.2	:	Growth and production of citric acid on Glucose and NBD palm olein	110
3.2.1	:	On glucose	110
3.2.2	:	On 2% NBD palm olein	112
3.3.3	:	On 4% NBD palm olein	114
3.3	:	Lipase production	116
3.3.1	:	On 2 % NBD palm olein	116
3.3.2	:	On 4 % NBD palm olein	117
3.4	:	Activity of key enzymes in TCA cycle	117
3.4.1	:	On 2 % glucose	120
3.4.2	:	On 2 % NBD palm olein	122
3.4.3	:	On 4 % NBD palm olein	124
3.5	:	Involvement of glyoxylate cycle	127
3.5.1	:	On 2 % glucose	127
3.5.2	:	On 2 % NBD palm olein	127

3.5.3 :	On 4 % NBD palm olein	130
3.6 :	Production of citric acid by <i>Yarrowia lipolytica</i> F21A at fermenter level	133
3.7 :	Genetic analysis of strain F21A	135
3.7.1 :	Screening of auxotrophic mutants	136
3.7.2 :	Induction of sporulation in auxotrophic derivatives of strain M240 and F21A	137
3.7.3 :	Mating of auxotrophic derivative of M240 and F21A	137
3.8 :	Improved citric acid production in <i>Yarrowia lipolytica</i> via DNA recombinant techniques	138
3.8.1 :	Assessment of vectors to be used as cloning vehicles	138
3.8.1.1 :	pINA62	139
3.8.1.2 :	Digestion pINA62 with <i>Pst</i> I	141
3.8.1.3 :	pINA214 and pINA230	145
3.8.1.4 :	Digestion pINA230 with <i>Bam</i> HI	151
3.8.2 :	Digestion of chromosomal DNA of <i>Yarrowia lipolytica</i> with <i>Sau</i> 3AI	152
3.8.3 :	Cloning of <i>Sau</i> 3AI-generated DNA fragments of <i>Yarrowia lipolytica</i> with vector DNA of <i>Bam</i> HI-digested of pINA230	152
<b>CHAPTER FOUR :</b>	<b>DISCUSSION</b>	
4.1	Isolation of mutants with altered ability to produce citric acid	156
4.2 :	Growth and citric acid production of glucose and NBD palm olein	161
4.3	Production of citric acid by <i>Yarrowia lipolytica</i> at fermenter level	171
4.4	Mating type determination and genetic analysis	175
4.5	Improved citric acid production in <i>Yarrowia lipolytica</i> via DNA recombinant techniques	177
REFERENCES		190
APPENDIX		210



## ABBREVIATIONS

ACH	Aconitate hydratase
BPB	Bromophenol blue
BSA	Bovine serum albumin
CIAP	Calf intestinal alkaline phosphatase
DTNB	5,5'-dithio-bis(2-nitrobenzoic acid)
EDTA	Ethylenediaminetetra-acetic acid
EMS	Ethylmethane sulphonate
HPLC	High performance liquid chromatography
ICDH	Isocitrate dehydrogenase
MFA	Monofluoroacetate
MFC	Monofluorocitrate
MgSO <sub>4</sub> ·7H <sub>2</sub> O	Magnesium sulphate 7-hydrate
μM	Micromolar
NBD	Neutralised bleached deodorised
NTG	N-methyl-N'-nitro-N-nitrosoguanidine
PEG	Polyethylene glycol
PVA	Polyvinyl alcohol
SDS	Sodium dodecyl sulphate
TBE	Tris-Boric-EDTA
TE	Tris-HCL-EDTA
TPN	Triphosphopyridine nucleotide
μg	Microgram
YEPA	Yeast extract peptone potassium acetate
YEPD	Yeast extract peptone glucose

## LIST OF FIGURES

Figure	Page
1      Growth of <i>Yarrowia lipolytica</i> on YEPD medium	3
2      Oil palm tree ( <i>Elaeis guineensis</i> )	5
3      Life cycle of <i>Yarrowia (Saccharomycopsis) lipolytica</i>	9
4      The lipase reaction	11
5      The net lipase reaction	11
6      Structure of alkylating agents	16
7      Plasmid pINA62 with <i>LEU2</i> marker from <i>Yarrowia lipolytica</i>	20
8      Plasmid transformation of yeast using either spheroplasts or lithium-treated cells	23
9      A. Fruit bunch of <i>Elaeis guineensis</i> hybrid <i>Tenera</i> B. Ripe palm oil fruit	25
10     Production of various refined fractions from crude palm oil	28
11     NBD palm olein at 28°C	30
12     Possible roles of peroxisomes in connection with mitochondria and microsomes in alkane assimilation yeasts	40
13     A. Oxidation pathway of alkanes to the corresponding fatty acids B. Fatty acid $\beta$ -oxidation system in yeast peroxisomes	41
14     The citric acid cycle and glyoxylate cycles	43
15     HPLC of standard mixture of citric acid and isocitric acid	81
16     The anatomy of fermenter	102
17     Growth and citric acid production of <i>Yarrowia lipolytica</i> (F21) utilising 2% NBD palm olein as the carbon source	105

18	Acid forming capacity as shown by size of halo formed on minimal medium incorporated with bromocresol green	106
19	Two different morphological colonies <i>Yarrowia lipolytica</i> grown on YEPD plate medium after 48 hours incubation	109
20	Growth and citric acid production by different strains of <i>Yarrowia lipolytica</i> utilising 2% glucose as the carbon source	111
21	Growth and citric acid production by different strains of <i>Yarrowia lipolytica</i> utilising 2% NBD palm olein as the carbon source	113
22	Growth and citric acid production by different strains of <i>Yarrowia lipolytica</i> utilising 4% NBD palm olein as the carbon source	115
23	Growth and extracellular lipases activity by different strains of <i>Yarrowia lipolytica</i> utilising 2% NBD palm olein as the carbon source	118
24	Growth and extracellular lipases activity by different strains of <i>Yarrowia lipolytica</i> 4% NBD palm olein as the carbon source	119
25	The TCA cycle enzymes activity of by different strains <i>Yarrowia lipolytica</i> utilising 2% glucose as the carbon source	121
26	The TCA cycle enzymes activity by different strains of <i>Yarrowia lipolytica</i> utilising 2% NBD palm olein as the carbon source	123
27	The TCA cycle enzymes activity by different strains of <i>Yarrowia lipolytica</i> utilising 4% NBD palm olein as the carbon source	125
28	The glyoxylate cycle enzymes activity by different strains of <i>Yarrowia lipolytica</i> utilising 2% glucose as the carbon source	128
29	The glyoxylate cycle enzymes activity by different strains of <i>Yarrowia lipolytica</i> utilising 2% NBD palm olein as the carbon source	129
30	The glyoxylate cycle enzymes activity by different strains of <i>Yarrowia lipolytica</i> utilising 4% NBD palm olein as the carbon source	131

31	Growth and production of citric acid on 2% NBD palm olein at fermenter level	134
32	Growth and production of citric acid on 4% NBD palm olein at fermenter level	134
33	Growth of <i>Escherichia coli</i> (HB101) transformant on M9 medium without leucine	140
34	Agarose (0.7%) gel electrophoresis of plasmid pINA62 extracted from <i>Escherichia coli</i> transformant	142
35	Agarose (0.7%) gel electrophoresis of chromosomal DNA and plasmid pINA62 extracted from <i>Escherichia coli</i> JM109 transformant	143
36	Genetic map of plasmid pINA214	146
37	Genetic map of plasmid pINA230	147
38	Growth of <i>Yarrowia lipolytica</i> strains M240, F21A and non-reversable <i>leu</i> strain M24062 on minimal medium after 48 hours incubation	149
39	Agarose (0.7%) gel electrophoresis of plasmid pINA230 extracted from <i>Escherichia coli</i> JM109 transformant	153
40	Agarose (0.7%) gel electrophoresis of chromosomal DNA <i>Yarrowia lipolytica</i> digested with <i>Sau</i> 3AI	154
41	A method for cloning segments of foreign DNA into <i>Yarrowia lipolytica</i>	181
42	A. A cleavage of two strands of DNA results by restriction endonuclease <i>Bam</i> HI B. Ligation of identical cohesive <i>Bam</i> HI end with T <sub>4</sub> DNA ligase	185
43	Ligation of a fragment with compatible cohesive ends into plasmid vector treated with calf intestinal alkaline phosphatase	188

## LIST OF TABLES

Table		Page
1	Oil palm: Area, production and yield	6
2	World's requirement and Malaysian production	6
3	Composition of palm oil	26
4	Fatty acid composition of palm oil	27
5	Fatty acid components of NBD palm olein	29
6	Composition of citric acid producing media by <i>Yarrowia lipolytica</i> utilising coconut and olive oil as the carbon source	32
7	<i>Yarrowia lipolytica</i> strains used in this study	46
8	Enzymes used in this study	49
9	Amino acids and nucleic acids used in this study	52
10	Substituted substrates in place of sodium acetate in the tricarboxylic acid medium	53
11	Components in 1 ml reaction mixture of citrate synthase activities	85
12	Components in 1 ml reaction mixture of aconitate hydratase activities	87
13	Components in 1 ml reaction mixture of isocitrate lyase activities	87
14	Components in 1 ml reaction mixture of malate synthase activities	88
15	Components in 1 ml reaction mixture of isocitrate dehydrogenase activities	89
16	Minimal medium supplemented with appropriate amino acids or nucleic acid bases	91
17	Effects of sodium monofluoroacetate (MFA) on growth of <i>Yarrowia lipolytica</i>	107

18	Utilisation by different strains <i>Yarrowia lipolytica</i> of carbon sources related to TCA cycle after 24 hours incubation	108
19	Morphology of colonies and cells of different strains of <i>Yarrowia. lipolytica</i> grown on YEPD plate medium after 48 hours incubation	108
20	The maximum activity of TCA cycle enzymes assayed during growth of different strains of <i>Yarrowia lipolytica</i> on glucose, 2% and 4% NBD palm olein as the carbon source	126
21	The maximum activity attained by enzymes of the glyoxylate cycle during growth of by different strains of <i>Yarrowia lipolytica</i> on glucose, 2% and 4% NBD palm olein as the carbon source	132
22	Auxotrophic mutants of <i>Yarrowia lipolytica</i> derived from M240 (A) and its mutants F21A (B)	136
23	Transformants obtained after transformation of <i>Yarrowia lipolytica</i> with pINA62	144
24	Transformants population of <i>Escherichia coli</i> (HB101) after transformation with pINA230 ( <i>Bam</i> HI-digested) ligated of <i>Sau</i> 3AI-generated fragments of tetracycline in <i>Yarrowia lipolytica</i>	155
25	The utilisation of carbon sources by <i>Yarrowia lipolytica</i> strains ATCC 201111, K-20 and S-22 (M243) on medium containing di- or tricarboxylic acids	157