

6: References

- Agudelo, P., Robbins, R. T., Stewart, J. M. and Murphy, J. B. (2004). Glycoproteins in the gelatinous matrix of *Rotylenchulus reniformis*. *Nematropica*, Vol. 34, No. 2: 229-234.
- Baldwin, J. G. (1978). The morphology of nematodes. *Nematology*, circular No.41.
- Belin, C., Schmitt, C., Demangeat, G., Komar, V., Pinck, L. and Fuchs, M. (2001). Involvement of RNA2-encoded proteins in the specific transmission of grapevine fanleaf virus by its nematode vector *Xiphinema index*. *Virology*, Vol. 291, No. 1: 161-171.
- Berry, S. D., Fargette, M., Spaul, V. W., Morand, S. and Cadet, P. (2008). Detection and quantification of root-knot nematode (*Meloidogyne javanica*), lesion nematode (*Pratylenchus zaeae*) and dagger nematode (*Xiphinema elongatum*) parasites of sugarcane using real-time PCR. *Molecular and Cellular Probes*, Vol. 22, No. 3: 168-176.
- Beyrem, H., Mahmoudi, E., Essid, N., Hedfi, A., Boufahja, F. and Aïssa, P. (2007). Individual and combined effects of cadmium and diesel on a nematode community in a laboratory microcosm experiment. *Ecotoxicology and Environmental Safety*, Vol. 68, No. 3: 412-418.
- Bird, D. M., Opperman, C. H. and Davies, K. G. (2003). Interactions between bacteria and plant-parasitic nematodes: now and then. *International Journal for Parasitology*, Vol. 33, No. 11: 1269-1276.
- Bridge, J., Fogain, R. and Speijer, P. (1997). The root lesion nematodes of banana: *Pratylenchus coffeae* (Zimmermann, 1898) Filip. & Schu. Stek., 1941 *Pratylenchus goodeyi* Sher & Allen, 1953. *Musa pest fact sheet*, No. 2. Montpellier: International Network for the Improvement of Banana and Plantain.
- Bridge, J. and Starr, J. L. (2007). *Plant nematodes of agricultural importance: a color handbook*. London: Manson Publishing Ltd. Pp. 152.
- Brown, T. A. (2001). *Gene cloning and DNA analysis: an introduction (4th ed.)*. Manchester: Blackwell Science Ltd. Pp. 363.
- Cadwell, R. C. and Joyce, G. F. (1994). Mutagenic PCR. *PCR Methods and Applications*, Vol. 3, No. 6: S136-S140.
- Campbell, C. S., Wojciechowski, M. F., Baldwin, B. G., Alice, L. A. and Donoghue, M. J. (1997). Persistent nuclear ribosomal DNA sequence polymorphism in the *Amelanchier* agamic complex (Rosaceae). *Molecular Biology and Evolution*, Vol. 14, No. 1: 81-90.

Chromas software version 2.33. Technelysium Pty Ltd. Available at <http://www.technelysium.com.au/chromas.html>.

- Demangeat, G., Komar, V., Cornuet, P., Esmenjaud, D. and Fuchs, M. (2004). Sensitive and reliable detection of grapevine fanleaf virus in a single *Xiphinema index* nematode vector. *Journal of Virological Methods*, Vol. 122, No. 1: 79-86.
- Fallas, G. A., Hahn, M. L., Fargette, M., Burrows, P. R. and Sarah, J. L. (1996). Molecular and biochemical diversity among isolates of *Radopholus* spp. from different areas of the world. *Journal of Nematology*, Vol. 28, No. 4: 422-430.
- Flemming, A. J., Shen, Z. Z., Cunha, A., Emmons, S. W. and Leroi, A. M. (2000). Somatic polyploidization and cellular proliferation drive body size evolution in nematodes. *Proceedings of the National Academy of Sciences, USA*. Vol. 97, No. 10: 5285-5290.
- Food and Agriculture Organization of the United Nations, 2009.
<http://faostat.fao.org/site/339/default.aspx>. Accessed on 2 July 2009.
- Hamilton, K., Barfoot, J., Crawford, K. E., Simpson, C. G., Beaumont, P. C. and Bownes, M. (2006). Amplification of chloroplast DNA using the polymerase chain reaction (PCR): a practical activity for secondary school students. *JBE*, Vol. 40, No. 4: 172-177.
- Holterman, M., Wurff, A. V. D., Elsen, S. V. D., Megen, H. V., Bongers, T., Holovachov, O., Bakker, J. and Helder, J. (2006). Phylum-wide analysis of SSU rDNA reveals deep phylogenetic relationships among nematodes and accelerated evolution toward crown clades. *Molecular Biology and Evolution*, Vol. 23, No. 9: 1792-1800.
- Hooper, D. J. (1974). *Commonwealth institute of helminthology, description of plant-parasitic nematode*. England: Commonwealth Agriculture Bureau. Set4, No. 50.
- Hugall, A., Stanton, J. and Moritz, C. (1999). Reticulate evolution and the origins of ribosomal internal transcribed spacer diversity in apomictic Meloidogyne. *Molecular Biology and Evolution*, Vol. 16, No. 2: 157-164.
- Inserra, R. N., Dunn, R. A., Mcsorley, R., Langdon, K. R. and Richmer, A. Y. (1989). Weed hosts of *Rotylenchulus reniformis* in ornamental nurseries of southern Florida. *Nematology*, circular No. 171.
- Kaplan, D. T., Thomas, W. K., Frisse, L. M., Sarah, J. L., Stanton, J. M., Speijer, P. R., Marin, D. H. and Opperman, C. H. (2000). Phylogenetic analysis of geographically diverse *Radopholus similis* via rDNA sequence reveals a monomorphic motif. *Journal of Nematology*, Vol. 32, No. 2: 134-142.

- Kleynhans, K. P. N. (1999). *Collecting and preserving nematodes: a manual for nematology*. South Africa: Plant Protection Research Institute. Pp. 54.
- Kumari, S., Kundu, J. K. and Polak, J. (2004). Identification of *Xiphinema vuittenezi* by polymerase chain reaction. *Plant Protect. Sci.*, Vol. 40, No. 1: 1-4.
- Madani, M., Subbotin, S. A. and Moens, M. (2005). Quantitative detection of the potato cyst nematode, *Globodera pallida*, and the beet cyst nematode, *Heterodera schachtii*, using Real-Time PCR with SYBR green I dye. *Molecular and Cellular Probes*, Vol. 19, No. 2: 81-86.
- MEGA4 software version 4.0. Available at <http://www.megasoftware.net/mega.html>.
- Mes, T. H. M. and Cornelissen, A. W. C. A. (2004). Non-homogenized ITS regions in the parasitic nematode *Cooperia oncophora*. *Parasitology*, Vol. 129, No. 2: 213-222.
- Musabyimana, T. and Saxena, R. C. (1999). Efficacy of neem seed derivatives against nematodes. *Phytoparasitica*, Vol. 27, No.1: 43-49.
- Nadler, S. A., Hoberg, E. P., Hudspeth, D. S. S. and Rickard, L. G. (2000). Relationships of *Nematodirus* species and *Nematodirus battus* isolates (Nematoda: Trichostrongyloidea) based on nuclear ribosomal DNA sequences. *Journal of Parasitology*, Vol. 86, No. 3: 588-601.
- Newton, L. A., Chilton, N. B., Beveridge, I. and Gasser, R. B. (1998). Genetic evidence indicating that *Cooperia surnabada* and *Cooperia oncophora* are one species. *International Journal for Parasitology*, Vol. 28, No. 2: 331-336.
- Nickle, W. R. (1970). A Taxonomic Review of the Genera of the Aphelenchoidea (Fuchs, 1937) Thorne, 1949 (Nematoda: Tylenchida). *Journal of Nematology*, Vol. 2, No. 4: 375-392.
- O'Bannon, J. H. and Inserra, R. N. (1989). *Helicotylenchus* species as crop damaging parasitic nematodes. *Nematology*, circular No. 165.
- Piotte, C., Arthaud, L., Abad, P. and Rosso M. N. (1999). Molecular cloning of an acetylcholinesterase gene from the plant parasitic nematodes, *Meloidogyne incognita* and *Meloidogyne javanica*. *Molecular and Biochemical Parasitology*, Vol. 99, No. 2: 247-256.
- Powers, T. O., Todd, T. C., Burnell, A. M., Murray, P. C. B., Fleming, C. C., Szalanski, A. L., Adams, B. A. and Harris, T. S. (1997). The rDNA internal transcribed spacer region as a taxonomic marker for nematodes. *Journal of Nematology*, Vol. 29, No. 4: 441-450.

- Raut, S. P. and Ranade, S. (2004). *Diseases of Fruits and Vegetables: Volume II*. Dordrecht: Kluwer Academic. Pp.704.
- Sarah, J. L., Pinochet, J. and Stanton, J. (1996). The burrowing nematode of bananas, *Radopholus similis* Cobb, 1913. *Musa pest fact sheet*, No. 1. Montpellier: International Network for the Improvement of Banana and Plantain.
- Scholl, E. H., Thorne, J. L., McCarter, J. P. and Bird, D. M. (2003). Horizontally transferred genes in plant-parasitic nematodes: a high-throughput genomic approach. *Genome Biology*, Vol. 4, No. 6: R39.
- Speiger, P. R. and Waele, D. D. (1997). Screening of *Musa* germplasm for resistance and tolerant to nematodes. *INIBAP technical guidelines*. Montpellier: International Network for the Improvement of Banana and Plantain. Pp 47.
- Subbotin, S. A., Ragsdale, E. J., Mullens, T., Roberts, P. A., Mundo-Ocampo, M. and Baldwin, J. G. (2008). A phylogenetic framework for root lesion nematodes of the genus *Pratylenchus* (Nematoda): evidence from 18S and D2–D3 expansion segments of 28S ribosomal RNA genes and morphological characters. *Molecular Phylogenetics and Evolution*, Vol. 48, No. 2: 491-505.
- Tesarova, B., Zouhar, M. and Rysanek, P. (2003). Development of PCR for specific determination of Root-knot nematode *Meloidogyne incognita*. *Plant Protect. Sci.*, Vol. 39, No.1: 23-28.
- Tylka, G. L. and Jaslavich, C. A. (2001). Free-living and plant-parasitic nematodes (Roundworms). *The Plant Health Instructor*. DOI: 10.1094/PHI-K-2001-0409-01.
- Wang, K. H. (2007). Reniform nematode, *Rotylenchulus reniformis* linfor and oliveira (Nematoda: Tylenchida: Tylenchoidea: Hoplolaimidae: Rotylenchulinae). *EENY-210*. Entomology and Nematology Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Available at <http://creatures.ifas.ufl.edu>.
- Wang, X., Li, H., Hu, Y., Fu, P. and Xu, J. (2007). Molecular cloning and analysis of a new venom allergen-like protein gene from the root-knot nematode *Meloidogyne incognita*. *Experimental Parasitology*, Vol. 117, No. 2: 133-140.
- Ward, J. G., Blomberg, P., Hoffman, N. and Yao, M. C. (1997). The intranuclear organization of normal, hemizygous and excision-deficient rRNA genes during developmental amplification in *Tetrahymena thermophila*. *Chromosoma*, Vol. 106, No. 4: 233-242.
- Wrona, A. F., Blasingame, D., Goodell, P., Gazaway, W. S., Mueller, J., Overstreet, C. and Wheeler, T. (1996). Timely soil sampling a nematode's Nemesis. *Cotton Physiology Today*, Vol. 7, No. 5: 21-24.