

## CONTENTS

ABSTRACT

ABSTRAK

ACKNOWLEDGEMENT

ABBREVIATION

CONTENTS

LIST OF FIGURES

LIST OF TABLES

1.0 : INTRODUCTION	1
1.1 : Salmonellosis	1
1.1.1 : Classification and Characteristics	1
1.1.2 : Disease Syndromes	2
1.1.3 : Diagnosis, Treatment and Prevention	4
1.2 : Laboratory Diagnosis of Typhoid Fever	6
1.2.1 : Isolation and Culture	6
1.2.2 : Serology	7
1.2.3 : Antigen Detection	8
1.2.4 : DNA Detection	9
1.3 : Polymerase Chain Reaction (PCR)	12
1.3.1 : PCR Optimization	12
1.3.1.1 : <i>Taq</i> DNA polymerase	15

1.3.1.2 : Primer	16
1.3.1.3 : Magnesium chloride	17
1.3.1.4 : Deoxynucleotide triphosphate (dNTP)	18
1.3.1.5 : Buffers	18
1.3.1.6 : Template DNA	19
1.3.2 : Limitations of PCR	19
1.3.2.1 : Error rate of <i>Taq</i> polymerase	19
1.3.2.2 : Contamination	20
1.3.3 : <i>S. typhi</i> Detection by PCR	20
1.4 : The 16S rRNA	24
1.4.1 : The Use of The 16S rRNA Gene in Probe Construction	25
1.5 : Hybridization	29
1.5.1 : Dot blots (Slot-blot)	29
1.5.2 : Southern Blotting	30
1.6 : DNA Sequencing	30
1.7 : Objectives of This Study	32
2.0 : MATERIALS AND METHODS	33
2.1 : Bacterial Strains	33
2.2 : Materials	34
2.3 : Media, Buffers and Other Solutions	36
2.3.1 : Luria- Bertani (LB) Medium (Miller, 1972)	36
2.3.2 : Common Solutions for DNA Extraction	36

2.3.3 : Solutions for Agarose Gel Electrophoresis	37
2.3.3.1 : 10X Tris-borate EDTA (TBE) buffer, pH 8.3	37
2.3.3.2 : 6X Bromophenol blue (BPB) loading dye	37
2.3.4 : Common Reagents for Slot-blot and Southern Hybridization	38
2.3.4.1 : 0.2M NaOH	38
2.3.4.2 : 50X Tris EDTA (TE) buffer	38
2.3.4.3 : 20X SSC	38
2.3.4.4 : 2X SSC	39
2.3.4.5 : Depurination solution	39
2.3.4.6 : 1M Tris-HCl, pH 7.5	39
2.3.4.7 : Denaturation solution	39
2.3.4.8 : Neutralization solution for slot blots (3M sodium acetate)	40
2.3.4.9 : Neutralization solution for southern blots	40
2.3.4.10 : 15X SSPE	40
2.3.4.11 : 10 % SDS	41
2.3.4.12 : 50% PEG 8000	41
2.3.4.13 : Prehybridization buffer	41
2.3.4.14 : 2X SSC + 0.1% SDS	42
2.3.4.15 : 0.2X SSC + 0.1% SDS	42
2.3.4.16 : 0.5M EDTA, pH 8.0	42
2.3.4.17 : Deprobing solution	43
2.3.5 : Solutions for Southern Blotting	43

2.3.5.1 : Solutions for labeling DNA probes	43
2.3.5.2 : Solutions for hybridization and blot washing	43
2.3.5.3 : Hybridization buffer	44
2.3.5.4 : Primary wash buffer (without urea)	44
2.3.5.5 : Secondary wash buffer	44
2.3.5.6 : Signal generation and detection	45
2.3.6 : Solutions for DNA Sequencing	45
2.3.6.1 : 10mM deoxyribonucleoside 5'-triphosphate (dATP, dCTP, dGTP, dTTP)	45
2.3.6.2 : 10mM dideoxyribonucleoside 5'-triphosphate (ddATP, ddCTP, ddGTP, ddTTP)	45
2.3.6.3 : Solutions for 8% sequencing gels	46
2.3.6.4 : 8% Sequencing gel solutions for a gel of 30.0X 40.0X 0.04 cm <sup>3</sup>	47
2.4 : Sterilization Techniques	48
2.4.1 : Moist Heat	48
2.4.2 : Dry Heat	48
2.5 : Methods	48
2.5.1 : Preparation of Bacterial Genomic DNA	48
2.5.2 : DNA Extraction from Serum Samples	50
2.5.3 : Determination of DNA Concentrations	51
2.5.4 : Oligonucleotide Primers for PCR Amplification	51
2.5.5 : PCR Amplification of rRNA Gene Regions and Analysis of PCR Products	52
2.5.6 : Electrophoresis of DNA	53

2.5.7 : Purification of PCR Products	54
2.5.8 : Slot-Blot Hybridization	55
2.5.8.1 : Preparation of hybridization membrane	55
2.5.8.2 : Labeling of hybridization probes	57
2.5.8.3 : Hybridization experiments	58
2.5.8.4 : Washing the hybridization membranes	59
2.5.8.5 : Developing the X-ray film	59
2.5.8.6 : Deprobing the hybridization membranes	60
2.5.9 : Southern Hybridization	60
2.5.9.1 : Southern Transfer of DNA from agarose gel to nylon membrane by capillary transfer	60
2.5.9.2 : Processing the gel	61
2.5.9.3 : Capillary blotting	61
2.5.9.4 : Processing the blot	62
2.5.9.5 : DNA labeling	62
2.5.9.6 : Prehybridization	64
2.5.9.7 : Hybridization	64
2.5.9.8 : Post-hybridization membrane washing	65
2.5.9.9 : Signal generation/detection	65
2.5.10 : DNA Sequencing	66
2.5.10.1 : Autoradiography	68
3.0 : RESULTS	70
3.1 : Isolation of Genomic DNA from <i>Salmonella</i> Species	70

3.2 : Determination of DNA Concentration	70
3.3 : Amplification of rRNA Gene Regions	70
3.4 : Optimization of PCR Conditions	77
3.5 : PCR Optimization and Specificity of Primers	78
3.5.1 : Effect of Different Annealing Temperatures on PCR	78
3.5.2 : Effect of Different MgCl <sub>2</sub> Concentrations on PCR	81
3.5.3 : Effect of Different DNA Concentrations on PCR	88
3.6 : Amplification of DNA from Clinical (Serum) Samples	90
3.7 : Southern Hybridization with the PCR-amplified V2 Region of <i>S. typhi</i> as the Probe	93
3.8 : Slot-Blot Hybridization with the PCR-amplified V2 Region as the Probe	96
3.9 : DNA Sequencing Results of the PCR-amplified V2 Region of <i>S. typhi</i> (1106) and <i>S. typhimurium</i>	98
4.0 : DISCUSSION	104
4.1 : Amplification of rRNA Gene Regions	105
4.2: PCR Optimization	106
4.3 : PCR Amplification for Clinical Samples	112
4.4 : Southern and Slot-blot Hybridization Using PCR-amplified V2 Region as Probe	113
4.5 : DNA Sequencing of the PCR-amplified Product of the V2 Region	115
4.6 : Suggestions for Further Work	118
5.0 : CONCLUSION	119
REFERENCES	121