

**ANTI-TUMOUR PROMOTER ACTIVITY  
AND PHYTOCHEMICAL STUDIES OF  
THREE SELECTED *Diospyros sp***

by

**ZURAIDAH HUSIN**

**A dissertation submitted in partial fulfilment  
of the requirement for the Degree of Masters of  
Biotechnology**

**Institute of Postgraduate Studies and Research  
University of Malaya  
Kuala Lumpur**

Perpustakaan Universiti Malaya



A508674662

**June, 1998**

Dimikrofisikan pada..... 14.08.2000  
No. Mikrofis..... 1H809  
Jumlah Mikrofis..... 2

HAMSIAH BT. MOHAMAD ZAHARI  
UPR UNIT REPROGRAFI  
PERPUSTAKAAN UTAMA  
UNIVERSITI MALAYA

## **ABSTRACT**

Three selected taxonomically identified *Diospyros* species namely *Diospyros graciliflora* Hiern., *D. discolor* Willd. and *D.lanceifolia* Roxburgh. were studied for their anti-tumour promoter activity. For *D. graciliflora* and *D. discolor*, the plant parts used were stem and stem bark, whereas in the case of *D. lanceifolia* only the stem was used. The dried powdered samples were extracted with petroleum ether (PE), chloroform and methanol respectively to yield all together 15 crude extracts. These crude extracts were preliminary screened for bioactive compounds using brine shrimp (*Artemia salina* Leach.) lethality procedure. The PE extracts of *D. discolor* and *D. lanceifolia* stem and the PE extract of *D. discolor* stem bark exhibited the LC<sub>50</sub> values of 122.4, 196.5 and 146.7 ppm respectively. The *D. graciliflora* stem exhibited a lower LC<sub>50</sub> value of 301.7 ppm. The other extracts were not active against brine shrimp with LC<sub>50</sub> values of above 3 500 ppm.

The three crude extracts positive for brine shrimp lethality (LC<sub>50</sub> < 200 ppm) and one with low activity (LC<sub>50</sub> 307.1 ppm) were further tested for their anti-tumour promoter activities by observing their inhibitory effect on 12-O-tetradecanoyl-phorbol-13-acetate (TPA) induced Epstein-Barr virus early antigen (EBV EA) in Raji cells. The crude PE extract of *D. discolor* stem exhibited inhibitory effect with IC<sub>50</sub> of 16.43 µg/ml. The other crude extracts, did not exhibit any EBV EA inhibitory effects.

Phytochemical studies were also carried out on these four crude extracts. The results of phytochemical analysis showed that the major secondary metabolites were terpenes. Separation of compounds from the crude PE extract of *D. discolor* stem was then conducted using column chromatography and preparative TLC method. From this procedure, three pure compounds were yielded and labelled as Dd P2, Dd P3 and Dd P6. Dd P2, Dd P3 and Dd P6 were identified as lupeol, betulin and betulinic acid respectively, identified by comparison with authentic compounds and spectroscopic analysis. Experiments for anti-tumour promoter activities were further conducted on these pure compounds. Only lupeol which was labelled as Dd P2 exhibited inhibitory activity against EBV EA induction with IC<sub>50</sub> value of 0.009 µg/ml.

Five plants from the family of Euphorbiaceae which were known for their tumour promoter activity were also collected from University of Malaya Botanical Garden, Rimba Ilmu. They were *Euphorbia hirta* L., *E. tirucalli* L., *E. splendens*, *Jatropha podagraria* Hook. and *Pedilanthus tithymaloides* (L.) Poitt.. These plants were extracted with PE in order to obtain natural phorbol esters. Experiments for the anti-tumour promoter activity were then conducted on lupeol, betulin and betulinic acid with the phorbol esters from the Euphorbiaceae plants, used in place of TPA, to induce EBV EA in Raji cells. Of these three compounds, only lupeol showed strong inhibitory activity while betulin and betulinic acid were ineffective as anti-tumour promoter in the same treatment system. The IC<sub>50</sub> values of *E. hirta*, *E. tirucalli*, *E. splendens*, *J. podagraria* and *P. tithymaloides* induced EBV EA treated with lupeol were 0.012, 0.014, 0.013, 0.016 and 0.015 µg/ml respectively.

In conclusion, PE extract of *D. discolor* stem was found to be effective as an anti-tumour promoter. Three triterpenes namely lupeol, betulin and betulinic acid were isolated from the PE extract of *D. discolor* stem. Lupeol demonstrated the strongest anti-tumour promoter activity whereas betulin and betulinic acid were not active against EBV EA induction in Raji cells. Natural phorbol esters obtained from the Euphorbiaceae plants could be used in place of TPA as after induction with the phorbol esters, the equivalent values of inhibitory action by the PE extracts was obtained in Raji cells.

## **Abstrak**

Kajian bagi aktiviti anti-penggalak tumor telah dilakukan ke atas tiga spesies terpilih *Diospyros* yang telah dikenalpasti ciri-ciri taksonomi mereka, iaitu *Diospyros graciliflora* Hiern., *Diospyros discolor* Willd. dan *Diospyros lanceifolia* Roxburgh. Bahagian batang dan kulit batang bagi spesies *D. graciliflora* dan *D. discolor* telah digunakan untuk kajian, manakala bagi spesies *D. lanceifolia* pula, hanya bahagian batang yang digunakan. Sampel-sampel dalam bentuk serbuk kering telah diekstrak menggunakan larutan petroleum eter (PE), diikuti dengan klorofom dan akhir sekali dengan metanol bagi menghasilkan sejumlah 15 ekstrak mentah. Ekstrak-ekstrak mentah ini telah digunakan di dalam ujian awal bagi mengesan kehadiran sebatian-sebatian bioaktif dengan menggunakan kaedah kematian anak udang (*Artemia salina* Leach). Ekstrak mentah PE bagi bahagian batang *D. discolor* dan *D. lanceifolia*, juga kulit batang *D. discolor* masing-masing memberikan nilai LC<sub>50</sub> 122.4, 196.5 dan 146.7 ppm. Ekstrak batang *D. graciliflora* pula, menunjukkan nilai LC<sub>50</sub> yang lebih rendah iaitu 301.7 ppm. Ekstrak-ekstrak mentah lain adalah tidak aktif terhadap anak udang dengan nilai LC<sub>50</sub> melebihi 3 500 ppm.

Ujian bagi menentukan aktiviti anti-penggalak tumor telah dijalankan ke atas tiga ekstrak mentah yang positif terhadap ujian kematian anak udang (LC<sub>50</sub> < 200 ppm) dan satu ekstrak beraktiviti rendah (LC<sub>50</sub> 301.7 ppm) dengan menggunakan kaedah perencatan ke atas antigen awal virus Epstein-Barr yang digalakkkan oleh 12-O-tetradecanoyl-phorbol-

13-acetate (TPA) di dalam sel Raji. Ekstrak mentah PE bagi bahagian batang *D. discolor* menunjukkan kesan perencatan dengan  $IC_{50}$  bernilai 16.43  $\mu\text{g}/\text{ml}$ . Ekstrak-ekstrak mentah lain tidak memberikan sebarang kesan perencatan.

Kajian fitokimia telah juga dijalankan ke atas ke empat-empat ekstrak mentah tersebut. Keputusan analisis fitokimia menunjukkan bahawa sebatian kimia utama adalah daripada kelas terpen. Kaedah kromatografi turus dan persediaan kromatografi lapisan nipis telah digunakan untuk mengasingkan sebatian-sebatian di dalam ekstrak mentah PE bagi bahagian batang *D. discolor*. Tiga sebatian tulen telah diperoleh dan dilabelkan sebagai Dd P2, Dd P3 dan Dd P6. Pengecaman sebatian tersebut telah dilakukan melalui perbandingan dengan sampel autentik. Dd P2 telah dikenalpasti sebagai lupeol, manakala Dd P3 adalah betulin, Dd P6 pula adalah asid betulinik. Ujian bagi aktiviti anti-penggalak tumor juga dijalankan ke atas tiga-tiga sebatian tulen ini. Hanya lupeol yang dilabelkan sebagai Dd P2 menunjukkan aktiviti perencatan dengan nilai  $IC_{50}$  0.009  $\mu\text{g}/\text{ml}$ .

Lima jenis tumbuhan daripada family Euphorbiaceae yang telah diketahui menunjukkan aktiviti penggalak tumor telah juga dikutip dari Rimba Ilmu, University Malaya. Tumbuhan tersebut adalah *Euphorbia hirta* L., *E. tirucalli* L., *E. splendens*, *Jatropha podagraria* Hook. dan *Pedilanthus tithymaloides* (L.) Poitt. Tumbuhan tersebut telah diekstrak dengan menggunakan larutan PE untuk menghasilkan phorbol ester asli. Ujian bagi aktiviti anti-penggalak tumor telah dijalankan ke atas tiga-tiga sebatian tulen (lupeol, betulin dan asid betulinik) yang ditindakbalaskan bersama-sama phorbol ester

asli dari ke lima-lima tumbuhan tersebut yang bertindak sebagai pengganti kepada TPA. Keputusan ujikaji menunjukkan hanya lupeol memberikan kesan perencatan manakala betulin dan asid betulinik tidak menunjukkan sebarang aktiviti anti-penggalak tumor. Nilai IC<sub>50</sub> bagi *E. hirta*, *E. tirucalli*, *E. splendes*, *J. podagraria* dan *P. tithymaloides* adalah 0.012, 0.014, 0.013, 0.016 dan 0.015 µg/ml.

Kesimpulannya, didapati bahawa ekstrak mentah PE bagi bahagian batang *D. discolor* adalah efektif sebagai anti-penggalak tumor. Tiga jenis triterpene iaitu lupeol, betulin dan asid betulinik berjaya diasangkan daripada ekstrak PE batang *D. discolor*. Lupeol menunjukkan aktiviti anti-penggalak tumor paling berkesan manakala betulin dan asid betulinik adalah tidak aktif. Phorbol ester asli yang diperolehi daripada tumbuhan Euphorbiaceae boleh digunakan sebagai pengganti kepada TPA kerana selepas induksi dengan phorbol ester tersebut, kesan perencatan yang setara dengan ekstrak-ekstrak mentah PE diperoleh dalam sel Raji.

## **Acknowledgements**

I would like to express my sincere gratitude and heartfelt thanks to my supervisors, Dr. Norhanom Abdul Wahab and Prof. Dr. Muhammad B. Zakaria for their invaluable guidance, advice and constructive criticism throughout this study.

I would also like to thank my dear colleagues who have provided valuable encouragement and support in their own way throughout the success of this study.

I would like to acknowledge University of Malaya for awarding me with a postgraduate fellowship during the tenure of this study.

Last but not least, I wish to thank my husband, Abdul Karim and both my children, Abdullah Munzir and Ainnur Munirah for their neverending love and support.

## Table of Contents

	Page
Abstract	ii
Abstrak	v
Acknowledgements	viii
Table of Contents	ix
List of Figures	xiii
List of Tables	xvi
<b>CHAPTER 1 : INTRODUCTION</b>	<b>1</b>
<b>CHAPTER 2 : LITERATURE REVIEW</b>	<b>4</b>
2.1 Medicinal Plants	4
2.1.1 Anti-Tumour Plants	5
2.1.2 Anti HIV Plants	6
2.1.3 Anti-Malarial Plants	6
2.1.4 Anti-Spasmodic Plants	7
2.1.5 Anti-Asthmatic and Anti-Tussive Plants	8
2.1.6 Anti-Fertility Plants	9
2.2 The Ebenaceae	13
2.3 The Genus <i>Diospyros</i> Linn.	14

2.3.1	<i>Diospyros graciliflora</i> Hiern.	16
2.3.2	<i>Diospyros discolor</i> Willd.	17
2.3.3	<i>Diospyros lanceifolia</i> Roxburgh.	18
2.4	Anti-Tumour Promoter Activity	19
2.4.1	Anti-Tumour Promoter Activity in Herbage Vegetables and Spices Tree Plants	22
2.4.2	Anti-Tumour Promoter Activity of Medicinal Plants	23
2.5	Bioassays	48
2.5.1	Brine Shrimp Lethality Bioassay	48
2.5.2	Anti-Tumour Promoter Activity Bioassay	49
<b>CHAPTER 3 : MATERIAL AND METHODS</b>		51
3.1	Plant Materials	51
3.2	Extraction of Plant Materials	52
3.3	Phytochemical Analysis	52
3.3.1	Chromatographic Methods	53
i	Thin Layer Chromatography (TLC)	53
ii	Column Chromatography	53
3.3.2	Preparation of Reagents for Detection of Compounds	54
i	10% Vanillin in Concentrated Sulphuric Acid	54
ii	Dragendorff's Reagents	54
3.3.3	Detection of Compounds	54
i	Alkaloids	54

ii Terpenoids	54
<b>3.4 Bioassays</b>	<b>55</b>
<b>3.4.1 Bioassay for Bioactive Compounds : Brine Shrimp Lethality Bioassay</b>	<b>57</b>
<b>3.4.2 Anti-Tumour Promoter Activity Bioassay : Inhibition of Epstein Barr Virus (EBV) Early Antigen (EA) Expression in Raji Cells</b>	<b>57</b>
i Preparation of Plant Extracts	57
ii Cell Line and Tissue Culture Medium	57
iii Inhibition of EBV EA Induction in Raji Cells	57
iv Reading of Slides	58
<b>3.4.3 Cytotoxicity</b>	<b>59</b>
<b>CHAPTER 4 : RESULTS</b>	<b>61</b>
<b>4.1 Bioassays</b>	<b>61</b>
<b>4.1.1 Brine Shrimp Lethality Bioassays</b>	<b>61</b>
<b>4.1.2 Anti-Tumour Promoter Activity Bioassays</b>	<b>63</b>
<b>4.1.3 Cytotoxicity of <i>Diospyros sp</i> in Raji Cells</b>	<b>72</b>
<b>4.2 Phytochemical Screening of Secondary Metabolites of <i>Diospyros graciliflora</i>, <i>Diospyros discolor</i> and <i>Diospyros lanceifolia</i></b>	<b>75</b>
<b>4.3 Phytochemical Investigation of PE Extract <i>D. discolor</i> Stem</b>	<b>89</b>
<b>4.3.1 Isolation of Compound Labelled Dd P2</b>	<b>90</b>

4.3.2 Isolation of Compound Labelled Dd P3	91
4.3.3 Isolation of Compound Labelled Dd P6	92
<b>CHAPTER 5 : DISCUSSIONS</b>	<b>93</b>
5.1 Bioactivity of <i>Diospyros sp</i>	93
5.1.1 Brine Shrimp Lethality Bioassay	93
5.1.2 Anti-Tumour Promoter Activity Bioassay	95
5.2 Phytochemical Screening of <i>Diospyros sp</i>	99
5.2.1 TLC Screening of <i>D. graciliflora</i> Stem	99
5.2.2 TLC Screening of <i>D. discolor</i> Stem	99
5.2.3 TLC Screening of <i>D. discolor</i> Stem Bark	100
5.2.4 TLC Screening of <i>D. lanceifolia</i> Stem	100
5.3 Structure Elucidation of Isolated Compounds of <i>D. discolor</i>	102
5.3.1 Isolated Compound Dd P2	102
5.3.2 Isolated Compound Dd P3	106
5.3.3 Isolated Compound Dd P6	109
5.4 Conclusion	112
<b>BIBLIOGRAPHY</b>	<b>114</b>
<b>APPENDIX</b>	<b>129</b>

## List of Figures

- Figure 2.1 Chemical structures of some active principles isolated from medicinal plants
- Figure 2.2 Three-stage carcinogenesis model (Berenbelum, 1975)
- Figure 2.3 Chemical structures of the euglobals isolated from the leaves and flower buds of *E. globulus*
- Figure 2.4 Chemical structures of the euglobals isolated from the leaves of *E. grandis*
- Figure 2.5 Chemical structures of the euglobal T1 isolated from the juvenile leaves of *E. tereticornis*
- Figure 2.6 Chemical structures of phloroglucinol derivatives from chloroform extract of *M. japonicus*
- Figure 2.7 Chemical structures of oleane- and lupane-type triterpenes isolated from the bark of *E. polyandra*
- Figure 2.8 Chemical structures of saponin isolated from the knots of *W. brachybotrys*
- Figure 2.9 Chemical structures of ursolic and oleanic acid from *G. hederaceae*
- Figure 2.10 Chemical structure of acycloartenoid triterpenes isolated from the leaves of *L. domesticum*
- Figure 2.11 Chemical structures of derivatives of 3-oxo-24-cycloarten-21-oic acid

- Figure 2.12 Chemical structures of bisbenzylisoquinoline alkaloids
- Figure 2.13 Chemical structures of the neolignans isolated from the bark of *M. officinalis*
- Figure 2.14 Chemical structures of some anthraquinones isolated from the plants of family Annonaceae
- Figure 2.15 Chemical structures of some naphtoquinones isolated from the plants of family Annonaceae
- Figure 2.16 Chemical structures of some azaanthraquinones isolated from the plants of family Annonaceae
- Figure 2.17 Chemical structures of angular-type furanocoumarins isolated from the fruits of *A. endulis*
- Figure 2.18 Chemical structures of  $\alpha$ -monolinolein isolated from the seed of *C. lachryma-jobi* var.*mayuen*
- Figure 3.1 Set-up of a tank for brine shrimp lethality bioassay
- Figure 4.1 Dose response curve of crude PE extract of *Diospyros sp* stem and stem bark on EBV EA expression induced by TPA in Raji cells
- Figure 4.2 Dose response curve of lupeol, betulin and betulinic acid on EBV EA expression induced by TPA in Raji cells
- Figure 4.3 Dose response curve of lupeol, betulin and betulinic acid on EBV EA expression induced by PE extract of *E. hirta* in Raji cells
- Figure 4.4 Dose response curve of lupeol, betulin and betulinic acid on EBV EA expression induced by PE extract of *E. tirucalli* in Raji cells
- Figure 4.5 Dose response curve of lupeol, betulin and betulinic acid on EBV

- EA expression induced by PE extract of *E. splendes* in Raji cells  
Figure 4.6 Dose response curve of lupeol, betulin and betulinic acid on EBV  
EA expression induced by PE extract of *J. podagraria* in Raji cells  
Figure 4.7 Dose response curve of lupeol, betulin and betulinic acid on EBV  
EA expression induced by PE extract of *P. tithymaloides* in Raji cells  
Figure 4.8 Schematic isolation of *D. discolor*  
Figure 5.1 Fragmentation process of compound labelled as Dd P2  
Figure 5.2 Structure of lupeol \*  
Figure 5.3 Structure of betulin  
Figure 5.4 Structure of betulinic acid

## List of Tables

- Table 2.1 Some medicinal plants used as an active principles in traditional medicine
- Table 4.1 LC<sub>50</sub> values for brine shrimp lethality bioassays of various crude extracts of *Diospyros sp*
- Table 4.2 Dry weight of crude extracts of *Diospyros sp* stem and stem bark
- Table 4.3 TLC analysis of PE extract of stem of *D. graciliflora* stem
- Table 4.4 TLC analysis of chloroform extract of *D. graciliflora* stem
- Table 4.5 TLC analysis of methanol extract of *D. graciliflora* stem
- Table 4.6 TLC analysis of PE extract of *D. discolor* stem
- Table 4.7 TLC analysis of chloroform extract of *D. discolor* stem
- Table 4.8 TLC analysis of methanol extract of *D. discolor* stem
- Table 4.9 TLC analysis of PE extract of *D. discolor* stem bark
- Table 4.10 TLC analysis of chloroform extract of *D. discolor* stem bark
- Table 4.11 TLC analysis of methanol extract of *D. discolor* stem bark
- Table 4.12 TLC analysis of PE extract of *D. lanceifolia* stem
- Table 4.13 TLC analysis of chloroform extract of *D. lanceifolia* stem
- Table 4.14 TLC analysis of methanol extract of *D. lanceifolia* stem