

REFERENCES

REFERENCES

- Abe, H. & Doi, Y. (1996). Enzymatic and environmental degradation of racemic poly(3-hydroxybutyric acid)s with different stereoregularities. *Macromol.*, 29, 8683-8688.
- Abe, H., Doi, Y., Aoki, H., Akehata, T., Hori, Y. & Yamaguchi, A. (1995a). Physical properties and enzymatic degradability of copolymers of (R)-3-hydroxybutyric and 6-hydroxyhexanoic acids. *Macromol.*, 28, 7630-7637.
- Abe, H., Matsubara, I. & Doi, Y. (1995b). Physical properties and enzymatic degradability of polymer blends of bacterial poly[(R)-3-hydroxybutyrate] and poly[(R,S)-3-hydroxybutyrate] stereoisomers. *Macromol.*, 28, 844-853.
- Abe, H., Matsubara, I., Doi, Y., Hori, Y. & Yamaguchi, A. (1994). Physical properties and enzymatic degradability of poly(3-hydroxybutyrate) stereoisomers with different stereoregularities. *Macromol.*, 27, 6018-6025.
- Agamuthu, P. (1997, July). *Solid waste management*. Paper presented for the lecture of HXGD6103-Pollution control and waste treatment technology, University of Malaya, Kuala Lumpur, Malaysia.
- American Public Health Association (APHA). (1989). *Standard methods for the examination of water and waste water*. Washington, DC.
- American Society for Testing and Materials (ASTM). (1992). D5209-91: Standard test method for determining the aerobic biodegradation of plastic materials in the presence of municipal sewage sludge. In *Annual book of ASTM standards* (pp.815-818). Philadelphia.
- Berger, K. G. (1983). Palm oil. In H. T. Chan (Ed), *Handbook of tropical foods* (pp. 433-468). New York: Dekker.
- Braunegg, G., Sonnleitner, B. & Lafferty, R. M. (1978). A rapid gas chromatography method for the determination of poly- β -hydroxybutyric acid in microbial biomass. *Eur. J. Appl. Microbiol. Biotechnol.*, 6, 29-37.
- Briese, B-H., Schmidt, B., Jendrossek, D. (1994). *Pseudomonas lemoignei* has fine poly(hydroxyalkanoic acid) (PHA) depolymerase genes: a comparative study of bacterial and eukaryotic depolymerases. *J. Environ. Polym. Degrad.*, 2, 75-87.
- Brock, T. D., Madigan, M. T., Martinko, J. M. & Parker, J. (1994). *Biology of microorganisms* (pp. 327-329, 7th ed.). Englewood Cliffs, N.J.: Prentice-Hall.

- Byrom, D. (1994). Polyhydroxyalkanoates. In D. P. Mobley (Ed.), *Plastics from microbes* (pp. 5-33). Munich: Hanser Publishers.
- Chang, H. N., Kim, Y. C., Lee, S. Y. and Kim, B. S. (1994). Current status of biodegradable plastics in Korea: research, commercial production and government policy. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp. 286-297). Amsterdam: Elsevier.
- Chiellini, E. (1997, Nov). An Overview of environmentally degradable polymers. *UNIDO-ICS International workshop on environmentally degradable polymers* (pp. 25-93). Pune, India.
- Cox, M.K. (1994). Properties and applications of polyhydroxyalkanoates. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp. 120-135). Amsterdam: Elsevier.
- Doi, Y. (1990). *Microbiol polyester* (pp. 11-31, 89-98, 135-150). New York: UCH Publish.
- Doi, Y. (1995). Microbial synthesis, physical properties and biodegradability of polyhydroxyalkanoates. *Macromol. Symp.*, 98, 585-599.
- Doi, Y., Kanesawa, Y., Kunioka, M., Saito, T. (1990). Biodegradation of microbial copolymers: poly (3-hydroxybutyrate-*co*-3-hydroxyvalerate) and poly (3-hydroxybutyrate-*co*-4-hydroxybutyrate). *Macromols.*, 23, 26-31.
- Doi, Y., Kasuya, K., Abe, H., Koyama, N., Ishiwatari, S., Takagi, K. & Yoshida, Y. (1996). Evaluation of biodegradabilities of biosynthetic and chemosynthetic polyesters in river water. *Polym. Degrad. Stab.*, 51, 281-286.
- Doi, Y., Mukai, K., Kasulya, K. & Yamada, K. (1994). Biodegradation of biosynthetic and chemosynthetic polyhydroxyalkanoates. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp. 39-51). Amsterdam: Elsevier.
- Fukui, T. & Doi, Y. (1997). Cloning and analysis of the poly(3-hydroxybutyrate-*co*-3-hydroxyhexanoate) biosynthesis genes of *Aeromonas caviae*. *J. Bacteriol.*, 179, 4821-4830.
- Furniss, B. S., Hannaford, A. J., Smith, P.W.G. & Tatchell, A.R. (Eds.) (1989). *Vogel's textbook of practical organic chemistry* (5th ed.). London: Longman.
- Grushko, Y. A. M. (1992). *Handbook of dangerous properties of inorganic and organic substances in industrial wastes* (pp. 217, 220-225). Boca Raton: CRC Press.
- Henry, J. (1997). Implications of product-based environmental management standards on industry. Paper presented at *APEC seminar on environmental management standards and their implications on global trade*. Singapore.

- Huang, S. J., Ho, L. H., Huang, M. T., Keonig, M. F. & Cameron, J. A. (1994). Similarities and differences between biodegradation and non-enzymatic degradation. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp.3-10). Amsterdam: Elsevier.
- Imam, S. H., Gordon, S. H., Shogren, R. L., Tsteson, T. R., Govind, N. S. & Greene, R. V. (1999). Degradation of starch-poly(β -hydroxybutyrate-co- β -hydroxyvalerate) bioplastic in tropical coastal waters. *Appl. Environ. Microbiol.*, Feb, 431-437.
- Jabatan Kimia. (n.d.). *Petunjuk makmal SC341, SC342* [Laboratory manual] (pp.17-19). Universiti of Malaya at Kuala Lumpur.
- Jendrossek, D., Schirmer, A. & Schlegel, H. G. (1996). Biodegradation of polyhydroxyalkanoate acids. *Appl. Microbiol. Biotechnol.*, 46, 451-463.
- Kanesawa, Y., Tanabashi, N., Doi, Y. & Saito, T. (1994). Enzymatic degradation of microbial poly (3-hydroxyalkanoates). *Polym. Deg. Stab.*, 45, 179-185.
- Kimura, M., Toyota, K., Iwatsuki, M. & Sawada, H. (1994). Effects of soil conditions on biodegradation of plastics and responsible microorganisms. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp.92-108). Amsterdam: Elsevier.
- Kitano, M. & Yakabe, Y. (1994). Strategy for biodegradability testing in OECD . In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp.217-227). Amsterdam: Elsevier.
- Koyama, N. & Doi, Y. (1995). Morphology and biodegradability of a binary blend of poly[(R)-3-hydroxybutyric acid] & poly[(R,S)-lactic acid]. *Can. J. Microbiol.*, 41, 316-322.
- Lageveen, R. G., Huisman, G. W., Preusting, H., Ketalaar, P., Eggink, G. & Witholt, B. (1988). Formation of polyesters by *Pseudomonas oleovorans*. *Appl. Environ. Microbiol.*, 54, 2924-2932.
- Marchessault, R. H., Monasterios, C. J., Jesudason, J. J., Ramsay, B., Saracovan, I., Ramsay, J. & Saito, T. (1994). Chemical, enzymatic and microbial degradation of bacterial and synthetic poly- β -hydroxyalkanoates. *Polym. Deg. Stab.*, 45, 187-196.
- Mergaert, J., Anderson, C., Wouters, A., Swings, J. & Kersters, K. (1992). Biodegradation of polyhydroxyalkanoates. *FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Rev.*, 317-322.
- Mergaert, J., Webb, A., Anderson, C., Wouters, A. & Swings, J. (1994). Microbial degradation of poly (3-hydroxybutyrate) & poly (3-hydroxybutyrate-co-3-hydroxyvalerate) in compost. *J. Environ. Polym. Degrad.*, 2, 177-183.

- Mergaert, J., Wouters, A., Anderson, C. & Swings, J. (1995). In situ biodegradation of poly (3-hydroxybutyrate) and poly (3-hydroxybutyrate-co-3-hydroxyvalerate) in natural waters. *Can. J. Microbiol.*, 41, 154-159.
- Merrick, J. M. & Doudoroff, M. (1964). Depolymerisation of poly- β -hydroxybutyrate by an intracellular enzyme syntase. *J. Bacteriol.*, 88, 60-71.
- Minear, R. A. & Keith, L. H. (Eds.) (1984). *Water analysis : Vol. 3. Organic species* (pp. 43-45). London: Academic Press.
- Molitoris, H. P., Moss, S. T., Koning, G. J. M. & Jendrossek, D. (1996). Scanning electron microscopy of polyhydroxyalkanoate degradation by bacteria. *Appl. Microbiol. Biotechnol.*, 46, 570-579.
- Mukai, K., Yamada, K. & Doi, Y. (1994). Efficient hydrolysis of polyhydroxyalkanoates by *Pseudomonas stutzeri* YM1414 isolated from lake water. *Polym. Deg. Stab.*, 43, 319-327.
- Muller, R. J., Aufusta, J., Walter, T. & Widdecke, H. (1994). The development and modification of some special test methods and the progress in standardisation of test methods in Germany. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp. 237-249). Amsterdam: Elsevier.
- Narayan, R. (1997, Nov). Standards development for biodegradable plastics. In *UNIDO-ICS international workshop on environmentally degradable polymers* (pp. 314-316). Pune, India.
- Nebel, B. J. & Wright, R. T. (1996). *Environmental science – the way the world works*. (pp. 349 & 545, 5th ed., International edition). U.S.A.: Prentice Hall.
- Pagga, U., Beimborn, D. B., Boelens, J. & Wilde, B. De. (1995). Determination of the aerobic biodegradability of polymeric material in laboratory controlled composting test. *Chemosphere*, 31, 11/12, 4475-4487.
- Pirt, S. J. (1975). *Principle of microbe & cell cultivation*. Great Britain: William Clowes & Son.
- Ramsay, B. A., Saracovan, I., Ramsay, J. A. & Marchessault, R. H. (1994). A method for the isolation of microorganisms producing extracellular long-side-chain poly (β -hydroxyalkanoate) depolymerase. *J. Environ. Polym. Deg.*, 2, 1-7.
- Reade, E. (Ed.) (1991). *Microbiological Technique* (pp. 124-125). Melbourne: School of Microbiology, University of Melbourne.

- Scott, G. (1994). Environmental biodegradation of hydrocarbon polymers: initiation and control. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp.79-91). Amsterdam: Elsevier.
- Shimadzu Corporation. (no date). *Instruction manual – total organic carbon analyzer model TOC-5000A*. Japan.
- Shimamura, E., Scandola, M. & Doi, Y. (1994). Microbial synthesis and characterization of poly (3-hydroxybutyrate-co-3-hydroxypropionate). *Macromols.*, 27, 4429.
- Smet, M. J. de, Eggink, G., Witholt, B., Kingma, J. & Wynberg, H. (1983). Characterization of intracellular inclusions formed by *Pseudomonas oleovorans* during growth on octane. *J. Bacteriol.*, 154, 870-878.
- Steinbuchel, A. & Valentin, H. E. (1995). Diversity of bacterial polyhydroxyalkanoic acids. *FEMS Microbiol. Lett.*, 128, 219-228.
- Streitwieser, A. & Heathcock, C. H. Jr (1976). *Introduction to organic chemistry* (pp. 340-341). New York: Macmillan Publishing Co., Inc.
- Sudesh, K. (1994). Production and characterization of polyhydroxyalkanoates (PHAs) by *Pseudomonas putida* PGA1 from palm oil & palm kernel oil derivatives. Master of Biotechnology dissertation, University of Malaya, Kuala Lumpur, Malaysia.
- Swift, G. (1994). Expectations for biodegradation testing methods. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp.228-236). Amsterdam: Elsevier.
- Tan, I. K. P., Kumar, K. S., Theanmalar, M., Gan, S. N. & Gordon III, B. (1997). Saponified palm kernel oil and its major free fatty acids as carbon substrates for the production of polyhydroxyalkanoates in *Pseudomonas putida* PGA1. *Appl. Microbiol. Biotechnol.*, 47, 207-211.
- Theanmalar, M. (1996). Studies of some cultivation conditions on polyhydroxyalkanoates (PHA) accumulation in *Pseudomonas putida* grown with palm kernel oil derivatives. Master of Biotechnology dissertation, University of Malaya, Kuala Lumpur, Malaysia.
- Tosi, C. & Ciampelli, F. (1973). Applications of infrared spectroscopy to ethylene-propylene copolymers. *Adv. Polym. Sci.*, 12, 87-129.
- Wade, L. G. (Ed.). (1987). *Organic chemistry*. Englewood Cliffs, N.J.: Prentice Hall.
- Wool, R. P. (1994). Perspective on standard test methods for biodegradable plastics. In Y. Doi and K. Fukuda (Eds.), *Biodegradable plastics and polymers* (pp.250-260). Amsterdam: Elsevier.