

BIBLIOGRAPHY

- Akopyanz, N. S., Eaton, K. A. and Berg, D. E. (1995). Adaptive mutation and cocolonization during *Helicobacter pylori* infection of gnotobiotic piglets. *Infect. Immun.* **63**, 116-121.
- Atlas, R. M. and Bej, A. K. (1994). Polymerase chain reaction. In Gerhardt, P. and Murray, R. G. E. (eds.), *Methods for General and Molecular Bacteriology*. Washington, D. C.: ASM Press (pp. 419-432).
- Bej, A. K., Mahbubani, M. H., Miller, R., DiCesare, J. L., Hoff, L. and Atlas, R. M. (1990). Multiplex PCR amplification and immobilized capture probes for detection of bacterial pathogens and indicators in water. *Mol. Cell. Probes* **4**, 353-365.
- Bell, D. A. and DeMarini, D. M. (1991). Excessive cycling converts PCR products to random-length higher molecular weight fragments. *Nucleic Acids Res.* **19(18)**, 5079.
- Birch, D. E., Kolmodin, L., Laird, W. J., McKinney, N., Wong, J., Young, K. K. Y., Zangenberg, G. A. and Zoccoli, M. A. (1996). Simplified hot start PCR. *Nature* **381**, 445-446.
- Bopp, C. A., Brenner, F. W., Wells, J. G. and Strockbine, N. A. (1999). *Escherichia*, *Shigella*, and *Salmonella*. In Murray, P. R., Baron, E. J., Pfaller, M. A., Tenover, F. C. and Tenover, R. H. (eds.), *Manual of Clinical Microbiology*, 7th edn. Washington, D. C.: ASM Press. (pp. 459-474).
- Broude, N. E., Zhang, L., Woodward, K., Englert, D. and Cantor, C. R. (2001). Multiplex allele-specific target amplification based on PCR suppression. *Proc. Natl. Acad. Sci. USA* **98(1)**, 206-211.
- Bubert, A., Hein, I., Rauch, M., Lehner, A., Yoon, B., Goebel, W. and Wagner, M. (1999). Detection and differentiation of *Listeria* spp. by a single reaction based on multiplex PCR. *Appl. Environ. Microbiol.* **65(10)**, 4688-4692.
- Caeiro, J.-P., Estrada-Garcia, M. T., Jiang, Z. D., Mathewson, J. J., Adachi, J. H., Steffen, R. and DuPont, H. L. (1999). Improved detection of enterotoxigenic *Escherichia coli* among patients with travelers' diarrhea, by use of the polymerase chain reaction. *J. Infect. Dis.* **180**, 2053-2055.
- Cantey, J. R. (1985). Shiga toxin - An expanding role in the pathogenesis of infectious diseases. *J. Infect. Dis.* **151**, 766-771.

Chamberlain, J. S., Gibbs, R. A., Ranier, J. E., Nguyen, P. N. and Caskey, C. T. (1988). Deletion screening of the Duchenne muscular dystrophy locus via multiplex DNA amplification. *Nucleic Acids Res.* **16**. 11141-11156.

Chamberlain, J. S., Gibbs, R. A., Ranier, J. E. and Caskey, C. T. (1990). Multiplex PCR for the diagnosis of Duchenne muscular dystrophy. In Innis, M. A., Gelfand, D. H., Sninsky, J. J. and White, T. J. (eds.), *PCR protocols: A guide to methods and applications*. San Diego: Academic Press. (pp. 272-281).

Chiu, C. H. and Ou, J. T. (1996). Rapid identification of *Salmonella* serovars in feces by specific detection of virulence genes, *invA* and *spvC*, by an enrichment broth culture-multiplex PCR combination assay. *J. Clin. Microbiol.* **34**(10), 2619-2622.

Chizhikov, V., Rasooly, A., Chumakov, K. and Levy, D. D. (2001). Microarray analysis of microbial virulence factors. *Appl. Environ. Microbiol.* **67**(7), 3258-3263.

Cimino, G. D., Metchette, K., Isaacs, S. T. and Zhu, Y. S. (1990). More false-positive problems. *Nature* **345**. 773-774.

Cohen, D., Ashkenazi, S., Green, M. S. *et al.* (1997). Double-blind vaccine-controlled randomized efficacy trial of an investigational *Shigella sonnei* conjugate vaccine in young adults. *Lancet* **349**. 155-159.

Dale, D. C. and Mata, L. J. (1968). Studies of diarrheal disease in Central America. XI. Intestinal bacterial flora in malnourished children with shigellosis. *Am. J. Trop. Med. Hyg.* **17**(3), 397-403.

Dorman, C. J. and Porter, M. E. (1998). The *Shigella* virulence gene cascade: A paradigm of bacterial gene control mechanisms. *Mol. Microbiol.* **29**(3), 677-684.

DuPont, H. L., Levine, M. M., Hornick, R. B. and Formal, S. B. (1989). Inoculum size in shigellosis and implications for expected mode of transmission. *J. Infect. Dis.* **159**(6), 1126-1128.

Dutta, S., Chatterjee, A., Dutta, P., Rajendran, K., Roy, S., Pramanik, K. C. and Bhattacharya, S. K. (2001). Sensitivity and performance characteristics of a direct PCR with stool samples in comparison to conventional techniques for diagnosis of *Shigella* and enteroinvasive *Escherichia coli* infection in children with acute diarrhoea in Calcutta, India. *J. Med. Microbiol.* **50**. 667-674.

Dytham, C. *Choosing and using statistics: A biologist's guide*. London: Blackwell Science Ltd. (p. 153).

Elnifro, E. M., Ashshi, A. M., Cooper, R. J. and Klapper, P. E. (2000). Multiplex PCR: Optimization and application in diagnostic virology. *Clin. Microbiol. Rev.* **13**(4), 559-570.

Erllich, H. A., Gelfand, D. and Sninsky, J. J. (1991). Recent advances in the polymerase chain reaction. *Science* **252**, 1643-1651.

Farnleitner, A. H., Kreuzinger, N., Kavka, G. G., Grillenberger, S., Rath, J. and Mach, R. L. (2000). Simultaneous detection and differentiation of *Escherichia coli* populations from environmental freshwaters by means of sequence variations in a fragment of the β -D-glucuronidase gene. *Appl. Environ. Microbiol.* **66**(4), 1340-1346.

Faruque, S. M., Khan, R., Kamruzzaman, M., Yamasaki, S., Ahmad, Q. S., Azim, T., Nair, G. B., Takeda, Y. and Sack, D. A. (2002). Isolation of *Shigella dysenteriae* type 1 and *S. flexneri* strains from surface waters in Bangladesh: Comparative molecular analysis of environmental *Shigella* isolates versus clinical strains. *Appl. Environ. Microbiol.* **68**(8), 3908-3913.

Fasano, A., Noriega, F. R., Liao, F. M., Wang, W. and Levine, M. M. (1997). Effect of *Shigella* enterotoxin 1 (ShET1) on rabbit intestine *in vitro* and *in vivo*. *Gut*. **40**(4), 505-511.

Fasano, A., Noriega, F. R., Maneval Jr, D. R., Chanasongram, S., Russel, R., Guandalini, S. and Levine, M. M. (1995). *Shigella* enterotoxin 1: An enterotoxin of *Shigella flexneri* 2a active in rabbit small intestine *in vivo* and *in vitro*. *J. Clin. Invest.* **95**, 2853-2861.

Fernandez-Prada, C. M., Hoover, D. L., Tall, B. D., Hartman, A. B., Kopelowitz, J. and Venkatesan, M. M. (2000). *Shigella flexneri* IpaH_{7.8} facilitates escape of virulent bacteria from the endocytic vacuoles of mouse and human macrophages. *Infect. Immun.* **68**(6), 3608-3619.

Forbes, K. J., Fang, Z. and Pennington, T. H. (1995). Allelic variation in the *Helicobacter pylori* flagellin genes *flaA* and *flaB*: Its consequences for strain typing schemes and population structure. *Epidemiol. Infect.* **114**, 257-266.

Frankel, G., Giron, J. A., Valmassoi, J. and Schoolnik, G. K. (1989). Multi-gene amplification: Simultaneous detection of three virulence genes in diarrhoeal stool. *Mol. Microbiol.* **3**(12), 1729-1734.

Frankel, G., Riley, L., Giron, J. A., Valmassoi, J., Friedmann, A., Strockbine, N., Falkow, S. and Schoolnik, G. K. (1990). Detection of *Shigella* in feces using DNA amplification. *J. Infect. Dis.* **161**, 1252-1256.

- Gaudio, P. A., Sethabutr, O., Echeverria, P. and Hoge, C. W. (1997). Utility of a polymerase chain reaction diagnostic system in a study of the epidemiology of shigellosis among dysentery patients, family contacts and well controls living in a shigellosis-endemic area. *J. Infect. Dis.* **176**, 1013-1018.
- Gelfand, D. H. and White, T. J. (1990). Thermostable DNA polymerases. In Innis, M. A., Gelfand, D. H., Sninsky, J. J. and White, T. J. (eds.), *PCR protocols: A guide to methods and applications*. San Diego: Academic Press. (pp. 129-141).
- Grattard, F., Berthelot, P., Reyrolle, M., Ros, A. Etienne, J. and Pozzetto, B. (1996). Molecular typing of nosocomial strain of *Legionella pneumophila* by AP-PCR. *J. Clin. Microbiol.* **34**, 1595-1598.
- Hacker, J., Blum-Oehler, G., Muhldorfer, I. and Tschape, H. (1997). Pathogenicity islands of virulent bacteria: Structure, function and impact on microbial evolution. *Mol. Microbiol.* **23**(6), 1089-1097.
- Hale, T. L. (1991). Genetic basis of virulence in *Shigella* species. *Microbiol. Rev.* **55**(2), 206-224.
- Harris, E., Kropp, G., Belli, A., Rodriguez, B. and Agabian, N. (1998). Single-step multiplex PCR assay for characterization of new world *Leishmania* complexes. *J. Clin. Microbiol.* **36**(7), 1989-1995.
- Hendolin, P. H., Markkanen, A., Ylikoski, J. and Wahlfors, J. J. (1997). Use of multiplex PCR for simultaneous detection of four bacterial species in middle ear effusions. *J. Clin. Microbiol.* **35**(11), 2854-2858.
- Henegariu, O., Heerema, N. A., Diouhy, S. R., Vance, G. H. and Vogt, P. H. (1997). Multiplex PCR: Critical parameters and step-by-step protocol. *BioTechniques* **23**, 504-511.
- Hirose, K., Itoh, K.-I., Nakajima, H., Kurazono, T., Yamaguchi, M., Moriya, K., Ezaki, T., Kawamura, Y., Tamura, K. and Watanabe, H. (2002). Selective amplification of *tyv* (*rfbE*), *prt* (*rfbS*), *viaB* and *fliC* genes by multiplex PCR for identification of *Salmonella enterica* serovars typhi and paratyphi A. *J. Clin. Microbiol.* **40**(2), 633-636.
- Hogrefe, H. H., Hansen, C. J., Scott, B. R. and Nielson, K. B. (2002). Archaeal dUTPase enhances PCR amplifications with archaeal DNA polymerases by preventing dUTP incorporation. *Proc. Natl. Acad. Sci. USA* **99**(2), 596-601.

Holland, J. L., Louie, L., Simor, A. E. and Louie, M. (2000). PCR detection of *Escherichia coli* O157:H7 directly from stools: Evaluation of commercial extraction methods for purifying fecal DNA. *J. Clin. Microbiol.* **38(11)**, 4108-4113.

Houng, H. S. H., Sethabutr, O. and Echeverria, P. (1997). A simple polymerase chain reaction to detect and differentiate *Shigella* and enteroinvasive *Escherichia coli* in human feces. *Diagn. Microbiol. Infect. Dis.* **28**, 19-25.

Hu, Y., Zhang, Q. and Meitzler, J. C. (1999). Rapid and sensitive detection of *Escherichia coli* O157:H7 in bovine faeces by a multiplex PCR. *J. Appl. Microbiol.* **87(6)**, 867-876.

Innis, M. A. and Gelfand, D. H. (1990). Optimization of PCRs. In Innis, M. A., Gelfand, D. H., Sninsky, J. J. and White, T. J. (eds.), *PCR protocols: A guide to methods and applications*. San Diego: Academic Press. (pp. 3-12).

Islam, D. and Lindberg, A. A. (1992). Detection of *Shigella dysenteriae* type 1 and *S. flexneri* in feces by immunomagnetic isolation and polymerase chain reaction. *J. Clin. Microbiol.* **30(11)**, 2801-2806.

Islam, M. S., Hossain, M. S., Hasan, M. K., Rahman, M. M., Fuchs, G., Mahalanabis, D., Baqui, A. H. and Albert, M. J. (1998). Detection of shigellae from stools of dysentery patients by culture and polymerase chain reaction techniques. *J. Diarrhoeal Dis. Res.* **16(4)**, 248-251.

Iverson, A. J. and Taylor, G. R. (1991). PCR in genetic diagnosis. In McPherson, M.J. (ed.), *PCR: A practical approach*. Oxford: Oxford University Press. (pp. 15-28).

Jegathesan, M. (1984). Serotype prevalence and antibiotic susceptibility of *Shigella* strains isolated in Malaysia during 1980 and 1981. *J. Diarrhoeal. Dis. Res.* **2(2)**, 102-104.

Jiang, Q., Hiratsuka, K. and Taylor, D. E. (1996). Variability of gene order in different *Helicobacter pylori* strains contributes to genome diversity. *Mol. Microbiol.* **20(4)**, 833-842.

Jin, Q., Yuan, Z., Xu, J. *et al.* (2002). Genome sequence of *Shigella flexneri* 2a: Insights into pathogenicity through comparison with genomes of *Escherichia coli* K12 and O157. *Nucleic Acids Res.* **30(20)**, 4432-4441.

Keusch, G. T. (2002). *Shigella*. In Sussman, M. (ed.), *Molecular Medical Microbiology*. London: Academic Press. (pp. 1279-1290).

- Khan, A. M., Rabbani, G. H., Faruque, A. S. G. and Fuchs, G. J. (1999). WHO-ORS in treatment of shigellosis. *J. Diarrhoeal Dis. Res.* **17(2)**, 88-89.
- Khan, W. A., Seas, C., Dhar, U., Salam, M. A. and Bennish, M. L. (1997). Treatment of shigellosis: V. Comparison of azithromycin and ciprofloxacin. *Ann. Intern. Med.* **126**, 697-703.
- Kocher, T. D. and Wilson, A. C. (1991). DNA amplification by the polymerase chain reaction. In Brown, T. A. (ed.), *Essential Molecular Biology*. Vol. II. Oxford: Oxford University Press. (p. 199).
- Kong, R. Y. C., Lee, S. K. Y., Law, T. W. F., Law, S. H. W. and Wu, R. S. S. (2002). Rapid detection of six types of bacterial pathogens in marine waters by multiplex PCR. *Water Res.* **36**, 2802-2812.
- Kongmuang, U., Luk, J. M. C. and Lindberg, A. A. (1994). Comparison of three stool-processing methods for detection of *Salmonella* serogroups B, C2 and D by PCR. *J. Clin. Microbiol.* **32(12)**, 3072-3074.
- Kotloff, K. L., Noriega, F. R., Samandari, T., Sztein, M. B., Losonsky, G. A., Nataro, J. P., Picking, W. D., Barry, E. M. and Levine, M. M. (2000). *Shigella flexneri* 2a strain CVD1207, with specific deletions in *virG*, *sen*, *set* and *guaBA*, is highly attenuated in humans. *Infect. Immun.* **68(3)**, 1034-1039.
- Kotloff, K. L., Winickoff, J. P., Ivanoff, B., Clemens, J. D., Swerdlow, D. L., Sansonetti, P. J., Adak, G. K. and Levine, M. M. (1999). Global burden of *Shigella* infections: Implications for vaccine development and implementation of control strategies. *Bull. WHO* **77(8)**, 651-666.
- Kramvis, A., Bukofzer, S. and Kew, M. C. (1996). Comparison of hepatitis B virus DNA extractions from serum by the QIAamp Blood Kit, GeneReleaser and the phenol-chloroform method. *J. Clin. Microbiol.* **34(11)**, 2731-2733.
- Kwok, S. and Higuchi, R. (1989). Avoiding false positives with PCR. *Nature*. **339**, 237-238.
- Lampel, K. A., Jagow, J. A., Trucksess, M. and Hill, W. E. (1990). Polymerase chain reaction for detection of invasive *Shigella flexneri* in food. *Appl. Environ. Microbiol.* **56(6)**, 1536-1540.

- Lee, S. E., Kim, S. Y., Kim, S. J., Kim, H. S., Shin, J. H., Choi, S. H., Chung, S. S. and Rhee, J. H. (1998). Direct identification of *Vibrio vulnificus* in clinical specimens by nested PCR. *J. Clin. Microbiol.* **36**(10), 2887-2892.
- Levine, M. M. (2000). Immunization against bacterial diseases of the intestine. *J. Pediatr. Gastroenterol. Nutr.* **31**(4), 336-355.
- Lichtsteiner, S. and Tjian, R. (1993). Cloning and properties of the *Caenorhabditis elegans* TATA-box-binding protein. *Proc. Natl. Acad. Sci. USA* **90**, 9673-9677.
- Lievano, F. A., Reynold, M. A., Waring, A. L., Ackelsberg, J., Bisgard, K. M., Sanden, G. N., Guris, D., Golaz, A., Bopp, D. J., Limberger, R. J. and Smith, P. F. (2002). Issues associated with and recommendations for using PCR to detect outbreaks of pertussis. *J. Clin. Microbiol.* **40**(8), 2801-2805.
- Lin, M. H., Chen, T. C., Kuo, T. T., Tseng, C. C. and Tseng, C. P. (2000). Real-time PCR for quantitative detection of *Toxoplasma gondii*. *J. Clin. Microbiol.* **38**(11), 4121-4125.
- Lim, Y. S. and Tay, L. (1991). Serotype distribution and antimicrobial resistance of *Shigella* isolates in Singapore. *J. Diarrhoeal Dis. Res.* **9**(4), 328-331.
- Lindqvist, R. (1999). Detection of *Shigella* spp. in food with a nested PCR method- and performance compared with a conventional culture method. *J. Appl. Microbiol.* **86**(6), 971-978.
- Luo, G. and Mitchell, T. G. (2002). Rapid identification of pathogenic fungi directly from cultures by using multiplex PCR. *J. Clin. Microbiol.* **40**(8), 2860-2865.
- Lüscher, D. and Altwegg, M. (1994). Detection of shigellae, enteroinvasive and enterotoxigenic *Escherichia coli* using the polymerase chain reaction (PCR) in patients returning from tropical countries. *Mol. Cell. Probes* **8**, 285-290.
- Markoulatos, P., Sifakas, N. and Moncany, M. (2002). Multiplex polymerase chain reaction: A practical approach. *J. Clin. Lab. Anal.* **16**, 47-51.
- Maurelli, A. T., Blackmon, B. and Curtiss III, R. (1984). Temperature-dependent expression of virulence genes in *Shigella* spp. *Infect. Immun.* **43**, 195-201.

- Maurelli, A. T., Fernández, R. E., Bloch, C. A., Rode, C. K. and Fassano, A. (1998). "Black holes" and bacterial pathogenicity: A large genomic deletion that enhances the virulence of *Shigella* spp. and enteroinvasive *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **95**, 3943-3948.
- McIngvale, S. C., Elhanafi, D. and Drake, M. A. (2002). Optimization of reverse transcriptase PCR to detect viable Shiga-toxin-producing *Escherichia coli*. *Appl. Environ. Microbiol.* **68**(2), 799-806.
- Mecenas, J. and Strauss, E. J. (1996). Molecular mechanisms of bacterial virulence: Type III secretion and pathogenicity islands. *Emerg. Infect. Dis.* **2**(4), 271-288.
- Meisenholder, G. (1999). New lids on the block. *The Scientist*. **13**(23), 17.
- Ménard, R., Prévost, M.-C., Gounon, P., Sansonetti, P. and Dehio, C. (1996). The secreted Ipa complex of *Shigella flexneri* promotes entry into mammalian cells. *Proc. Natl. Acad. Sci. USA* **93**, 1254-1258.
- Monteiro, L., Bonnemaïson, D., Vekris, A., Petry, K. G., Bonnet, J., Vidal, R., Cabrita, J. and Megraud, F. (1997). Complex polysaccharides as PCR inhibitors in feces: *Helicobacter pylori* model. *J. Clin. Microbiol.* **35**(4), 995-998.
- Morello, J. A., Mizer, H. E., Wilson, M. E. and Granato, P. A. (1998). *Microbiology in Patient Care*, 6th edn. Boston: WCB McGraw-Hill. (pp. 336-359).
- Moss, J. E., Cardozo, T. J., Zychlinsky, a. and Groisman, E. A. (1999). The *selC*-associated SHI-2 pathogenicity island of *Shigella flexneri*. *Mol. Microbiol.* **33**(1), 74-83.
- Mutter, G. L. and Boyton, K. A. (1995). PCR bias in amplification of androgen receptor alleles, a trinucleotide repeat marker used in clonality studies. *Nucl. Acid Res.* **23**, 1411-1418.
- Nataro, J. P., Seriwatana, J., Fasano, A., Maneval, D. R., Guers, L., D., Noriega, F., Dubovsky, F., Levine, M. M. and Morris Jr, J. G. (1995). Identification and cloning of a novel plasmid-encoded enterotoxin of enteroinvasive *Escherichia coli* and *Shigella* strains. *Infect. Immun.* **63**(12), 4721-4728.
- Niyogi, S. K., Dutta, P. and Mitra, U. (2000). Trends in antimicrobial resistance of *Shigella* spp. isolated from children with acute diarrhea. *Indian Pediatrics* **37**, 296-307.

- Noriega, F. R. (1995). ShMu, a protein of *Shigella flexneri* 2a with hemagglutinin and mucinase activities is encoded by an ORF (*she*) that forms an antisense gene pair with the operon encoding *Shigella* enterotoxin 1 (ShET1). GenBank accession number U35656.
- Noriega, F. R., Liao, F. M., Formal, S. B., Fasano, A. and Levine, M. M. (1995). Prevalence of *Shigella* enterotoxin 1 among *Shigella* clinical isolates of diverse serotypes. *J. Infect. Dis.* **172**, 1408-1410.
- O'Sullivan, B., Delpuch, V., Pontivivo, G., Karagiannis, T., Marriott, D., Harkness, J. and McAnulty, J. M. (2002). Shigellosis linked to sex venues, Australia. *Emerg. Infect. Dis.* **8(8)**, 862-864.
- Ochman, H. and Moran, N. A. (2001). Genes lost and genes found: Evolution of bacterial pathogenesis and symbiosis. *Science*. **292**, 1096-1098.
- Oyofe, B. A., Subekti, D., Tjaniadi, P. et al. (2002). Enteropathogens associated with acute diarrhea in community and hospital patients in Jakarta, Indonesia. *FEMS Immunol. Med. Microbiol.* **34**, 139-146.
- Parsot, C. and Sansonetti, P. J. (1996). Invasion and the pathogenesis of *Shigella* infections. *Curr. Topics Microbiol. Immun.* **209**, 25-42.
- Pass, M. A., Odedra, R. and Batt, R. M. (2000). Multiplex PCRs for identification of *Escherichia coli* virulence genes. *J. Clin. Microbiol.* **38(5)**, 2001-2004.
- Peng, X., Luo, W., Zhang, J., Wang, S. and Lin, S. Rapid detection of *Shigella* species in environmental sewage by an immunocapture PCR with universal primers. *Appl. Environ. Microbiol.* **68(5)**, 2580-2583.
- Penn, C. (1991). *Handling laboratory microorganisms*. Philadelphia: Open University Press. (pp. 88-99).
- Prats, G., Mirelis, B., Llovet, T., Munoz, C., Miro, E. and Navarro, F. (2000). Antibiotic resistance trends in enteropathogenic bacteria isolated in 1985-1987 and 1995-1998 in Barcelona. *Antimicrob. Agents Chemother.* **44(5)**, 1140-1145.
- Pupo, G. M., Lan, R. and Reeves, P. R. (2000). Multiple independent origins of *Shigella* clones of *Escherichia coli* and convergent evolution of many of their characteristics. *Proc. Natl. Acad. Sci. USA* **97(19)**, 10567-10572.

- Rahman, I., Shahamat, M., Chowdhury, M. A. R. and Colwell, R. R. (1996). Potential virulence of viable but nonculturable *Shigella dysenteriae* type 1. *Appl. Environ. Microbiol.* **62**(1), 115-120.
- Rajakumar, K., Sasakawa, C. and Adler, B. (1997). Use of a novel approach, termed island probing, identifies the *Shigella flexneri* she pathogenicity island which encoded a homolog of the immunoglobulin A protease-like family of proteins. *Infect. Immun.* **65**(11), 4606-4614.
- Rhee, S. J., Wilson, K. T., Gobert, A. P., Nataro, J. P. and Fasano, A. (2001). The enterotoxigenic activity of *Shigella* enterotoxin 1 (ShET1) is mediated by inducible nitric oxide synthase activity. *J. Pediatr. Gastroenterol. Nutr.* **33**(3), 400-416.
- Rolfs, A., Schuller, I., Finckh, U. and Weber-Rolfs, I. (1992). *PCR: Clinical diagnostics and research*. Berlin: Springer-Verlag. (p. 52).
- Rossen, L., Norskov, P., Holmstrom, K. and Rasmussen, D. F. (1992). Inhibition of PCR by components of food samples, microbial diagnostic assays and DNA-extraction solutions. *Int. J. Food Microbiol.* **17**, 37-45.
- Ruth, L. (1999). *Feces, Norwalk virus, rotavirus, urine*. In *Alkami Quick Guide™ for PCR*. Vol. 1. USA: Alkami Biosystems, Inc. (pp. 82-85).
- Saiki, R. K. (1989). Optimization of the polymerase chain reaction. In Erlich, H. A., Gibbs, R. and Kazanian Jr, H. H. (eds.), *Current Communications in Molecular Biology: Polymerase Chain Reaction*. Cold Spring Harbor: CSH Laboratory Press. (pp. 25-30).
- Saiki, R. K., Scharf, S., Faloona, F., Mullis, K. B., Horn, G. T., Erlich, H. A. and Arnheim, N. (1985). Enzymatic amplification of β -globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science*. **230**, 1350-1354.
- Sails, A. D., Bolton, F. J., Fox, A. J., Wareing, D. R. A. and Greenway, D. L. A. (2002). Detection of *Campylobacter jejuni* and *C. coli* in environmental waters by PCR enzyme-linked immunosorbent assay. *Appl. Environ. Microbiol.* **68**(3), 1319-1324.
- Saito, H. and Miura, K. I. (1963). Preparation of transforming deoxyribonucleic acid by phenol treatment. *Biochim. Biophys. Acta* **72**, 619-629.
- Sakai, T., Sasakawa, C. and Yoshikawa, M. (1988). Expression of four virulence antigens of *Shigella flexneri* is positively regulated at the transcriptional level by the 30kDa *virF* protein. *Mol. Microbiol.* **2**, 589-597.

- Salam, M. A., Seas, C., Khan, W. A. and Bennis, M. L. (1995). Treatment of shigellosis: IV. Cefixime is ineffective in shigellosis in adults. *Ann. Intern. Med.* **123**, 505-508.
- Salyers, A. A. and Whitt, D. D. (1994). *Bacterial pathogenesis: A molecular approach*. Washington, D. C.: ASM Press. (pp. 73-180).
- Sambrook, J. and Russell, D. (2001). *Molecular cloning: A laboratory manual*, 3rd edn. New York: Cold Spring Harbor Laboratory.
- Sandvig, K. (2001). Shiga toxins. *Toxicon*. **39**, 1629-1635.
- Sansonetti, P. J., Kopecko, D. J. and Formal, S. B. (1982). Involvement of a plasmid in the invasive ability of *Shigella flexneri*. *Infect. Immun.* **35**(3), 852-860.
- Sarkar, G. and Sommer, S. S. (1990). Shedding light on PCR contamination. *Nature* **343**, 27.
- Sasakawa, C., Kamata, K., Sakai, T., Murayama, S. Y., Makino, S. and Yoshikawa, M. (1986). Molecular alteration of the 140-megadalton plasmid associated with loss of virulence and Congo red binding activity in *Shigella flexneri*. *Infect. Immun.* **51**(2), 470-475.
- Schuch, R. and Maurelli, A. T. (1997). Virulent plasmid instability in *Shigella flexneri* 2a is induced by virulent gene expression. *Infect. Immun.* **65**(9), 3686-3692.
- Sethabutr, O., Venkatesan, M., Murphy, G. S. Eampokalap, B., Hoge, C. W. and Echeverria, P. (1993). Detection of shigellae and enteroinvasive *Escherichia coli* by amplification of the invasive plasmid antigen H DNA sequence in patients with dysentery. *J. Infect. Dis.* **167**, 458-461.
- Stacy-Phipps, S., Mecca, J. J. and Weiss, J. B. (1995). Multiplex PCR assay and simple preparation method for stool specimens to detect enterotoxigenic *Escherichia coli* DNA during course of infection. *J. Clin. Microbiol.* **33**(5), 1054-1059.
- Stone, G. G., Oberst, R. D., Hays, M. P., McVey, S. and Chengappa, M. M. (1994). Detection of *Salmonella* serovars from clinical samples by enrichment broth cultivation-PCR procedure. *J. Clin. Microbiol.* **32**(7), 1742-1749.

- Talukder, K. A., Islam, M. A., Dutta, D. K., Hassan, F., Safa, A., Nair, G. B. and Sack, D. A. (2002). Phenotypic and genotypic characterization of serologically atypical strains of *Shigella flexneri* type 4 isolated in Dhaka, Bangladesh. *J. Clin. Microbiol.* **40**(7), 2490-2497.
- Tamano, K., Katayama, E., Toyotome, T. and Sasakawa, C. (2002). *Shigella* Spa32 is an essential secretory protein for functional type III secretion machinery and uniformity of its needle length. *J. Bacteriol.* **184**(5), 1244-1252.
- Tellier, R., Bukh, J., Emerson, S. U. and Purcell, R. H. (1996). Amplification of the full-length hepatitis A virus genome by long reverse transcription-PCR and transcription of infectious RNA directly from the amplicon. *Proc. Natl. Acad. Sci. USA* **93**(9), 4370-4373.
- Theron, J., Morar, D., DuPreez, M., Brozel, V. S. and Venter, S. N. (2001). A sensitive seminested PCR method for the detection of *Shigella* in spiked environmental water samples. *Water Res.* **35**(4), 869-874.
- Tobe, T., Nagai, S., Okada, N., Adler, B., Yoshikawa, M. and Sasakawa, C. (1991). Temperature-regulated expression of invasive genes in *Shigella flexneri* is controlled through the transcriptional activation of the *virB* gene on the large plasmid. *Mol. Microbiol.* **5**, 887-893.
- Toye, B., Woods, W., Bobrowska, M. and Ramotar, K. (1998). Inhibition of PCR in genital and urine specimens submitted for *Chlamydia trachomatis* testing. *J. Clin. Microbiol.* **36**(8), 2356-2358.
- Vantarakis, A., Komninou, G., Venieri, D. and Papapetropoulou, M. (2000). Development of a multiplex PCR detection of *Salmonella* spp. and *Shigella* spp. in mussels. *Lett. Appl. Microbiol.* **31**(2), 105-109.
- Vargas, M., Gascon, J., DeAnta, M. T. J. and Vila, J. (1999). Prevalence of *Shigella* enterotoxins 1 and 2 among *Shigella* strains isolated from patients with traveler's diarrhea. *J. Clin. Microbiol.* **37**, 3608-3611.
- Venkatesan, M. M., Buysse, J. M. and Kopecko, D. J. (1989). Use of *Shigella flexneri* *ipaC* and *ipaH* gene sequences for the general identification of *Shigella* spp. and enteroinvasive *Escherichia coli*. *J. Clin. Microbiol.* **27**(12), 2687-2691.
- Venkatesan, M. M., Goldberg, M. B., Rose, D. J., Grotbeck, E. J., Burland, V. and Blattner, F. R. (2001). Complete DNA sequence and analysis of the large virulence plasmid of *Shigella flexneri*. *Infect. Immun.* **69**(5), 3271-3285.

Wang, G. Z., Miyashita, N. T. and Tsunewaki, K. (1997). Plasmon analyses of *Triticum* (wheat) and *Aegilops*: PCR-single-strand conformational polymorphism (PCR-SSCP) analyses of organellar DNAs. *Proc. Natl. Acad. Sci. USA* **94**, 14570-14577.

Wang, L., Rothmund, D., Curd, H. and Reeves, P. R. (2000). Sequence diversity of the *Escherichia coli* H7 *fliC* genes: Implications for a DNA-based typing scheme for *E. coli* O157:H7. *J. Clin. Microbiol.* **38**(5), 1786-1790.

Watanabe, H. and Nakamura, A. (1985). Large plasmids associated with virulence in *Shigella* species have a common function necessary for epithelial cell penetration. *Infect. Immun.* **48**(4), 260-262.

Weekly Epidemiological Record (WER). (1998). Outbreak news. *WER* **43**, 335.

Weekly Epidemiological Record (WER). (2000). Outbreak news. *WER* **75**, 29.

Welsh, J. and McClelland, M. (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res.* **18**, 7213-7218.

White, T. J., Madej, R. and Persing, D. H. (1992). The PCR: Clinical applications. *Adv. Clin. Chem.* **29**, 161-196.

Wilde, J., Eiden, J. and Yolken, R. (1990). Removal of inhibitory substances from human fecal specimens for detection of group A rotaviruses by reverse transcriptase and polymerase chain reaction. *J. Clin. Microbiol.* **28**, 1300-1307.

Yavzori, M., Cohen, D. and Orr, N. (2002). Prevalence of the genes for *Shigella* enterotoxins 1 and 2 among clinical isolates of *Shigella* in Israel. *Epidemiol. Infect.* **128**(3), 533-535.

Yavzori, M., Cohen, D., Wasserlauf, R., Ambar, R., Rechavi, G. and Ashkenazi, S. (1994). Identification of *Shigella* species in stool specimens by DNA amplification of different loci of the *Shigella* virulent plasmid. *Eur. J. Clin. Microbiol. Infect. Dis.* **13**(3), 232-237.