

## *REFERENCES*

Aaronson,S. 1970. Experimental Ecology. Academic Press Inc. (London) Ltd.  
pp.88.

Abdul Majid,M.I., Hori,K, Akiyama,M. and Doi,Y. 1994. Production of poly(3-hydroxybutyrate) from plant oils by *Alcaligenes* sp. In: Doi,Y. and Fukuda,K. (Eds). Biodegradable Plastics and Polymers. Elsvier Science B.V. pg 417-424.

Anderson,A.J. and Dawes,E.A. 1990. Occurrence, metabolism, metabolic role and industrial uses of bacterial polyhydroxyalkanoates. *Microbiol Rev.* 54:450-470.

Ashby,R.D., Foglia,T.A. 1998. Polyhydroxyalkanoates biosynthesis from triglyceride substrates. *Appl. Microbiol. Biotechnol.* 49:431-437.

Akiyama,M., Taima,Y. and Doi,Y. 1992. Production of poly(3-hydroxyalkanoates) by a bacterium of genus *Alcaligenes* utilising long chain fatty acids. *Appl. Microbiol. Biotechnol.* 37:698-701.

Brandl,H., Bachofen,R., Mayer,J. and Wintermantel,E. 1995. Degradation and application of polyhydroxyalkanoates. *Can. J. Micobiol.* 41(Supp 1):143-153.

Burdon,K.L. 1946. Fatty Acid material in bacteria and fungi revealed by staining dried, fixed slide preparations. *J. Bacteriol.* 52: 665-678.

Chesbro,W., Arbige,M. and Robin,E. 1990. When nutrient limitation places bacteria in the domain of slow growth: metabolic, morphologic and cell cycle behaviour. *FEMS Microbiol. Ecol.* 74:103-120.

Cocks,L.V. and Rede,C.V. 1966. Laboratory Handbook for Oil and Fat Analysis. Academic Press London & New York. pp.113-117.

Demain,A.L. and Soloman,N.A. 1986. Manual of Industrial Microbiology and Biotechnology. American Society for Microbiology Washington,D.C. pp.3-23.

de Smet,M.J., Eggink,G., Witholt,B., Kingma,J. and Wynberg,H. 1983. Characterisation of intracellular inclusion formed by *Pseudomonas oleovorans* during growth on octane. *J. Bacteriol.* 154:870-878.

Doi,Y. 1990. In Microbial Polyester. Research Laboratory of Resources Utilisation Tokyo Institute of Technology, Yokohama, Japan. VCH Publish, New York.

Doi,Y., Kawaguchi,Y., Nakamura,Y. and Kunioka,M. 1989. Nuclear Magnetic Resonance studies of Poly(3-hydroxybutyrate) and polyphosphate metabolism in *Alcaligenes eutrophus*. *Appl. Env. Microbiol.* 55:2932-2938.

Doi,Y., Kunioka,M., Nakamura,Y. and Soga,K. 1986.  $^1\text{H}$  and  $^{13}\text{H}$  NMR Analysis of poly( $\beta$ -hydroxybutyrate) isolated from *Bacillus megaterium*. *Macromol.* 19:1274-1276.

Doi,Y., Kunioka,M., Tamaki,A., Nakamura,Y. and Soga,K. 1988. Nuclear Magnetic Resonance studies on unusual bacterial copolyester of 3-hydroxybutyrate and 4-hydroxybutyrate. *Macromol.* 21:2722-2727.

Doi,Y., Segawa,A., Kagawaguchi,Y. and Kunioka,M. 1990. Cyclic nature of poly(3-hydroxyalkanoate) metabolism in *Alcaligenes eutrophus*. *FEMS Microbiol. Lett.* 67:165-170.

Eggink,G., de Waard,P. and Huijbert,G.N.M. 1995. Formation of novel polyhydroxyalkanoates from long-chain fatty acids. *Can. J. Microbiol.* 43(Suppl.1):14-21.

Eggink,G., van der Wal, H., Huijbert,G.N.M. and de Waard,P. 1993. Oleic acid as a substrate for poly-3-hydroxyalkanoate formation in *Alcaligenes eutrophus* and *Pseudomonas putida*. *Inds. Crops Products.* 1:157-163.

Fay,J.P. and Farias,R.N. 1975. The inhibitory action of fatty acids on the growth of *Escherichia coli*. *J. Gen. Microbiol.* 91:233-240.

Freese,E., Sheu,C.W. and Galliers,E. 1973. Function of lipophilic acids as antimicrobial food additives. *Nature*. 241:321-325.

Griffin,G.J.L. 1994. Chemistry and Technology of Biodegradable Polymers. Blackie Academic and Professional, U.K. pp.53.

Gross,R.A., DeMello,C. and Lenz,R.W. 1989. Biosynthesis and characterisation of poly( $\beta$ -hydroxyalkanoates) produced by *Pseudomonas oleovorans*. *Macromol.* 22:1106-1115.

Harder, W. and Dijkhuizen, L. 1983. Physiological responses to nutrient limitation. *Ann. Rev. Microbiol.* 37:1-23.

Havolson, H.O. 1957. Rapid and simultaneous sporulation. *J. Appl. Bacteriol.* 20(3):305-314.

Hrabak, O. 1992. Industrial production of poly- $\beta$ -hydroxybutyrate. *FEMS. Microbiol. Rev.* 103:251-256.

Huijberts,G.N.M., de Rijk,T.C., de Waard,P. and Eggink,G. 1994.  $^{13}\text{C}$  NMR studies of *Pseudomonas putida* fatty acids metabolic routes involved in poly (3-hydroxyalkanoate) synthesis. *J. Bacteriol.* 176:1661-1666.

Huijberts,G.N.M., Eggink,G., de waard,P., Huisman, G.W. and Witholt, B. 1992. *Pseudomonas putida* KT2442 cultivated on glucose accumulates poly (3-hydroxyalkanoates) consisting saturated and unsaturated monomers. *Appl. Environ. Microbiol.* 58:536-544.

Huijberts,G.N.M., van de Wal,H., Wilkinson,C. and Eggink,G. 1994. Gas-chromatographic analysis of poly(3-hydroxyalkanoates) in bacteria. *Biotechnol. Techn.* 8:187-192.

Huisman,G.W., de Leeuw,O., Eggink,G. and Witholt,B. 1989. Synthesis of poly-3-hydroxyalkanoates is a common feature of florescent pseudomonads. *Appl. Environ. Microbiol.* 55:1949-1954.

Iftikhar,A. 1984. Significance of palm oil and palm stearin as fatty raw materials for soap. PORIM Occ. Paper Palm Oil Res. Inst. Malaysia No.13. pp.3-9.

Jaeger,K.E., Ransac,S., Dijkstra,B.W., Cosson,C., van Hauvel,M. and Missel,O. 1994. Bacterial Lipases. *Fems Microbiol. Rev.* 15:29-61.

Kellarhals,M.B. Hazenberg,W. and Witholt,B. 1999. High cell density fermentations of *Pseudomonas oleovorans* for the production of mcl-PHAs in two-liquid phase media. *Enzyme and Microb. Technol.* 24:11-116.

Kim,B.S., Lee,S.Y. and Chang,H.N. 1993. Production of polyhydroxyalkanoate by fed-batch culture of recombinant *Escherichia coli*. *Biotechnol. Lett.* 14:811-816.

Kim, B.S., Lee,S.C. Lee,S.Y., Chang, Y.K. and Woo, S.I. 1993. Production of poly(3-hydroxybutyric acid) by fed-batch culture of *Alcaligenes eutrophus* with glucose concentration control. *Biotechnol. Bioeng.* 43:892-898.

Kim, S. W., Kim, P., Lee, H. S. and Kim, J. H. 1996. High production of poly- $\beta$ -hydroxybutyrate (PHB) from *Methylobacterium organophilum* under potassium limitation. *Biotechnol. Lett.* 18:25-30.

Kobayashi,G., Shiotani,T., Shima,Y. and Doi,Y. 1994. Biosynthesis and characterisation of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) from oils and fats by *Aeromonas* sp. OL-338 and *Aeromonas* sp. FA-440. In: Doi,Y. and Fukuda,K. (Eds). Biodegradable Plastics and Polymers. Elsvier Science B.V. pp.410-416.

5. Bacterial polyhydroxyalcanoates as biodegradable plastics with great future. *Chem. & Process Eng.* 1995, 16(10), 30-33.

Kominek, L.A. and Havolson, H.O. 1965. Metabolism of poly- $\beta$ -hydroxybutyrate and acetoin in *Bacillus cereus*. *J. Bacteriol.* 90(5):1251-1259.

Lageveen,R.G., Huisman,G.W., Preusting,H., Ketelaar,P., Eggink,G. and Witholt,B. 1988. Formation of polyesters by *Pseudomonas oleovorans*: Effect of substrates on formation and composition of poly-(R)-3-hydroxyalkanoates and poly-(R)-3-hydroxyalkenoates. *Appl. Environ. Microbiol.* 54:2924-2932.

Nishio,N., Tsuchiyu,Y., Hayashi,M. and Nagai,S. 1977. A fed-batch culture of methanol-utilising bacteria with pH-stat. *J. Ferment. Technol.* 55:151-155.

Nunn,W.D. 19886. A molecular view of fatty acid catabolism in *Escherichia coli*. *Microbiol. Rev.* 50:179-192.

Oeding,V., and Schlegel,H.G. 1973.  $\beta$ -Ketothiolase from *Hydrogemonas eutropha* H16 and its significance in the regulation of poly- $\beta$ -hydroxybutyrate metabolism. *Biochem. J.* 134:239-248.

Ostle,A.G. and Holt,J.G. 1982. Nile Blue A as a floorescent stain for poly- $\beta$ -hydroxybutyrate. *Appl. Environ. Microbiol.* 44:238-241.

Continuous p. 10

Page, W.J. 1995. Bacterial polyhydroxyalkanoates natural biodegradable plastics with great future. *Can. J. Microbiol.* 41(Suppl. 1):1-3.

Pirt, S. J. 1975. Principles of Microbe and Cell Cultivation. Blackwell Scientific Publication, Oxford. pp.5.

Preusting,H. Housten,R., van, Hoefs,A, Kool, Langenberghe, E.K. van, Favre-Bulle, O. and Witholt,B. 1992. High Cell Density Cultivation of *Pseudomonas oleovorans*: Growth and Production of Poly (3-hydroxyalkanoates) in Two-Liquid Phase Batch and Fed-Batch Systems. *Biotechnol. Bioeng.* 41:50-556.

Presuting, H., Nijenhuis, A. and Witholt, B. 1990. Physical characteristics of poly(3-hydroxyalkanoates) and Poly(3-hydroxyalkenoates) produced by *Pseudomonas oleovorans* grown on aliphatic hydrocarbons. *Macromol.* 23:4220-4224.

Ramsay, B.A., Ramsay, J.A. and Cooper, D.G. 1989. Production of poly- $\beta$ -hydroxyalkanoic acid by *Pseudomonas cepacia*. *Appl. Environ. Microbiol.* 55:584-589.

Ramsay, B.A., Saracovan, I., Ramsay, J.A. and Marchessault, R.H. 1991. Continous production of long-side-chain poly- $\beta$ -hydroxyalkanoates by *Pseudomonas oleovorans*. *Appl. Environ. Microbiol.* 57:625-629.

Redzwan,G., Gan,S.N. and Tan,I.K.P. 1997. Isolation of polyhydroxyalkanoate-producing bacteria from integrated-farming pond and palm oil-mill effluent ponds. *World J. Microbiol. Biotechnol.* 13:1-3.

Reusch, R. N. 1995. Low molecular weight complexed poly(3-hydroxybutyrate): a dynamic and versatile molecule in vivo. *Can. J. Microbiol.* 41 (Suppl. 1):50-54.

Senior,P.J., Beech,G.A., Richie,G.A.F. and Dawes,E.A. 1972. The role of oxygen limitation in the formation of poly- $\beta$ -hydroxybutyrate during batch and continuous culture of *Azotobacter beijerinckii*. *Biochem. J.* 128:1193-1201.

Schulz, H. and Kunau, W.H. 1987. Beta-oxidation of unsaturated fatty acids: a revised pathway. *Trends. Biochem. Sci.* 12:403-406.

Sheu,C.W. and Freese,E. 1972. Effects of fatty acids on growth and envelope proteins of *Bacillus substillis*. *J. Bacteriol.* 111:516-524.

Sheu,C.W.

Sheu,C.W. and Freese,E. 1973. Lipopolysaccharide layer protection of Gram-negative bacteria against inhibition by long chain fatty acids. *J. Bacteriol.* 115:869-875.

Shimamura,E., Kasuya,K., Kobayashi,G., Shiotani,T., Shima,Y. and Doi,Y. 1994. Physical properties and biodegradability of microbial poly(3-hydroxybutyrate-co-hydroxyhexanoate) synthesis. *Macromol.* 27:878-880.

Sirockin,G. and Cullimore,S. 1969. In:*Practical Microbiology*. Mc Graw-Hill, London, United Kingdom. pp.7-84.

Solorzono,L. 1969. Determination of Ammonia in natural waters by phenolhypochlorite method. *Limnol. Oceanogr.* 14: 799-801.

Spiekermann,P., Rehn,B.H.A., Kalscheuer,R., Baumeister,D. and Steinbuchel,A. 1999. A sensitive, viable-colony staining method using nile red for direct screening of bacteria that accumulate polyhydroxyalkanoic acids and other lipid storage compounds. *Arch. Microbiol.* 171:73-80.

Stainer,R.Y., Palleroni,N.J. and Dondorogff,M. 1966. The aerobic pseudomonads: A taxonomic study. *J. Gen. Microbiol.* 43:150-271.

Steinbuchel, A. and Schlegel,H.G. 1989. Excretion of pyruvate by mutants of *Alcaligenes eutrophus* which impaired in the accumulation of poly( $\beta$ -hydroxybutyrate)PHB under condition permitting synthesis of PHB. *Appl. Microbiol. Biotechnol.* 31:168-175

Suzuki,T., Yamane,T. and Shimizu,S. 1986(a). Mass production of poly beta hydroxybutyric acid by fed-batch culture with controlled carbon/nitrogen feeding. *Appl. Microbiol. Biotechnol.* 24:370-374.

Suzuki,T., Yamane,T. and Shimizu,S. 1986(b). Mass production of polyhydroxybutyric acid by fully fed-batch culture of methylotroph. *Appl. Microbiol. Biotechnol.* 23:322-329.

Suzuki,T., Yamane,T. and Shimizu,S. 1986(c). Kinetics and effects of nitrogen source feeding on production of polyhydroxybutyric acid by fed-batch culture. *Appl. Microbiol. Biotechnol.* 24:366-369.

Suzuki,T., Yamane,T. and Shimizu,S. 1990. Phenomenological background and some preliminary trials of automated substrate supply a pH-stat modal fed-batch culture using a set point of high limit. *J. Ferment. Bioeng.* 69:292-297.

Tan, I.K.P., Kumar,K.S., Theanmalar,M., Gan,S.N. and Gordon III,B. 1997. Saponified palm kernel oil and its major free fatty acids as carbon substrate for production of polyhydroxyalkanoate in *Pseudomonas putida* pGA1. *Appl. Microbiol. Biotechnol.* 47:207-211.

Tan, I.K.P., Stuart, E.S., Lenz, R.W. and Fuller, R.C. 1998. Compositional limitations in poly-3-hydroxyalkanoates produced by *Pseudomonas oleovorans*. *J. Environ. Polym. Degrad.* 6:67-71.

Theanmalar, M. 1996. Studies of some cultivation conditions on polyhydroxyalkanoates (PHA). Disser. for M.Biotechnol., Institute of Advanced Studies, University of Malaya. Malaysia.

Timm,A. and Steinbuchel,A. 1990. Formation of polyester consisting of medium-chain-length 3-hydroxyalkanoic acids from gluconate by *Pseudomonas aeruginosa* and other fluorescent pseudomonads. *Appl. Environ. Microbiol.* 56:3360-3367.

Zeng, F., Kondo, T. 1992.

using background information

Wallen,L.L. and Rohwedder,W.K. 1974. Poly- $\beta$ -hydroxyalkanoate from activated sludge. *Environ. Sci. Technol.* 8:576-579.

Week,G., Shapiro,M., Burns,R.O. and Wakil,S.J. 1969. Control of fatty acid metabolism. *J. Bacteriol.* 99:827-836.

Williams,D.H. and Fleming,I. 1989. Spectroscopic methods in organic chemistry. McGRAW-HILL Book Company Europe. pp. 29-62.

Williamson,D.H. and Wilkinson,J.F. 1958. The isolation and estimation of poly- $\beta$ -hydroxybutyrate inclusions of *Bacillus* sp. *J. Gen. Microbiol.* 19:198-209.

Witholt,B., Huisman,G.W. and Preusting,H. 1990. Bacterial poly(3-hydroxyalkanoates). Dawes, E.A. Novel Biodegradable Microbial Polymers.Kluwer Academic Publisher, Netherlands. Pp. 161-173.

Yamane,T., Sirifofe,P. and Shimizu,S. 1988. Economic significance of increasing cell mass concentration for metabolic significance. *J. Ferment. Technol.* 66(1):93-102.

Yano,T., Kobayashi,T. and Shimizu,S. 1978. Fed-batch culture of methanol utilising bacterium with DO-stat. *J. Ferment. Technol.* 56:416-420.