

**Tissue Culture and Protoplast
Isolation of *Gracilaria changii*
(Rhodophyta)**

By

Tan Eng Lai

Perpustakaan Universiti Malaya



A509841087

Dissertation submitted to the
Institute of Postgraduate Studies and Research,
University of Malaya,
Kuala Lumpur,
in partial fulfillment of
the requirement for
the degree of
Master of Biotechnology

November 1999

ACKNOWLEDGEMENTS

This dissertation would not have materialised without the support and assistance of several individuals. First and foremost, I would like to express my gratitude to my main supervisor, Associate Professor Dr. Phang Siew Moi for her strong encouragement and support during the time when I was working on this project. I would like to thank her for her invaluable guidance in every detail of the project and for all the knowledge and experience that she had cheerfully shared. Secondly, I would like to thank my co-supervisor Associate Professor Dr. Nurzulaani binti Khalid for her kind assistance and advice on the project.

My gratitude also goes to all my colleagues at the Algal Laboratory. I sincerely thank Annie, Baset, Gan, Melor, Miah, Muru, Qamrul, Rohani, Seok Min and Dr. Chu for creating such a friendly atmosphere in the laboratory making it simply one of the best in the institute. Though we worked on diverse individual projects, I truly cherish the moments when we worked together in cleaning the laboratory and entertaining visitors. The time that we shared during the many field trips will also be memorable.

I would like to thank my parents for their moral and financial support. I also thank them for encouraging me to pursue further in my education. Finally, my gratitude goes to the immortal, invisible, the only wise God who never fails to make a way for me when there seem to be no way and who works in ways I cannot see. All that I am and ever hoped to be, to God be the glory.

ABSTRACT

A system for tissue culture and protoplast isolation was investigated for *Gracilaria changii* (Rhodophyta) Abbott, Zhang and Xia. A plant sterilisation protocol for the generation of healthy and axenic explants required for both tissue culture and protoplast isolation was established. This method involved treating the seaweed thalli in ultra high quality (UHQ) water for an hour and sonicated for three to four minutes. This is then followed by treatment in 1% w/v povidone-iodine and incubated overnight in sterile seawater containing a series of antibiotics (0.1 g.L⁻¹ kanamycin, 0.3 g.L⁻¹ penicillin G, 0.22 g.L⁻¹ polymyxin B sulphate, 0.001 g.L⁻¹ nalidixic acid, 0.02 g.L⁻¹ cefotaxime and 1 g.L⁻¹ streptomycin sulphate).

Different forms and constituents of Murashige and Skoog (MS) and Provasoli's Enriched Seawater (PES) media were tested for callus induction and plant regeneration. Branching or shoot formation was only recorded in liquid PES + vitamins medium supplemented with different concentrations of growth substances. All the growth substances tested, except 2,4-D, induced branching. IAA (0.001 mg.L⁻¹) induced the highest incidence of branching (20%) followed by 0.001 mg.L⁻¹ kinetin, 0.1 mg.L⁻¹ NAA and 0.01 mg.L⁻¹ IAA in which 13.33% of the explants branched.

Friable callus-like structure (CLS) and shoots were initiated from cross-sections of explants in liquid PES + vitamins medium supplemented with different combinations and varying concentrations of 2,4-D and kinetin. A combination of 10 mg.L⁻¹ 2,4-D and 0.001 mg.L⁻¹ kinetin generated 100% incidence of CLS and shoot

formation. Incidence of CLS and shot formation was positively related to higher 2,4-D concentrations and lower kinetin concentrations.

Mannitol concentration of 1 M was found to be an excellent osmotic stabiliser for *G. changii* required for enzymatic cell wall digestion during protoplast isolation. Cells were sufficiently plasmolysed at this osmoticum concentration.

Enzymes such as abalone acetone powder, cellulase, pectinase, pectolyase and agarase were tested individually and in combinations in various concentrations and time of incubation for their effectiveness in protoplast isolation. Incubation of tissues in 1% w/v abalone acetone powder generated the highest protoplast yield (5.0×10^4 protoplast g^{-1} fresh weight). A combination of 1% w/v each of all the enzymes mentioned failed to achieve the maximum achieved using 1% w/v abalone acetone powder alone.

ABSTRAK

Sistem untuk pengkulturan tisu dan pengasingan protoplast *Gracilaria changii* (Rhodophyta) Abbott, Zhang dan Xia telah disiasat. Satu kaedah untuk menghasilkan bahagian tumbuhan yang sihat dan asenik telah berjaya ditetapkan. Kaedah ini melibatkan rendaman bahagian tumbuhan di dalam air berkualiti tinggi, UHQ (*ultra high quality*) selama satu jam dan sonikasi selama tiga hingga empat minit. Kemudian, bahagian tumbuhan tersebut dibersihkan dengan 1% w/v povidone-iodine dan direndamkan selama satu hari di dalam air laut yang mengandungi pelbagai jenis antibiotik (0.1 g.L⁻¹ kanamisin, 0.3 g.L⁻¹ penisilin G, 0.22 g.L⁻¹ polimisin B sulfat, 0.001 g.L⁻¹ asid nalidisik, 0.02 g.L⁻¹ sifotaksin dan 1 g.L⁻¹ streptomisin sulfat).

Medium Murashige dan Skoog (MS) dan *Provasoli's Enriched Seawater* (PES) dalam bentuk dan komposisi yang berlainan telahpun diuji untuk perkembangan kalus dan anak pokok. Hanya medium PES + vitamin yang mengandungi hormon-hormon pertumbuhan mengizinkan pertumbuhan anak pucuk. Semua hormon pertumbuhan yang telah diuji kecuali 2,4-D menyebabkan perkembangan anak pucuk. IAA (0.001 mg.L⁻¹) menyebabkan peratusan perkembangan anak pucuk yang paling tinggi (20%) diikuti dengan 0.001 mg.L⁻¹ kinetin, 0.1 mg.L⁻¹ NAA dan 0.01 mg.L⁻¹ IAA di mana 13.33% perkembangan anak pucuk telah direkodkan.

Struktur serupa kalus, CLS (*callus-like structure*) yang berbentuk longgar didapati berkembang daripada keratan rentas tumbuhan yang dikultur di dalam medium PES + vitamin yang mengandungi kombinasi kepekatan 2,4-D dan kinetin

yang berlainan. Kombinasi 10 mg.L⁻¹ 2,4-D dan 0.001 mg.L⁻¹ kinetin menyebabkan 100% perkembangan CLS dan anak pucuk. Peratusan perkembangan CLS dan anak pucuk bertambah dalam kepekatan 2,4-D yang tinggi dan kepekatan kinetin yang rendah.

Kepekatan manitol 1 M didapati bertindak sebagai penstabil osmotik yang baik untuk *G. changii* yang diperlukan semasa pencernaan dinding sel dalam proses pengasingan protoplast. Sel-sel telah diplasmolisiskan dengan sempurna dengan kepekatan manitol ini.

Enzim-enzim seperti serbuk *abalone acetone*, selulase, pektinase, pektoliase dan agarase telah diuji secara berasingan dan berkombinasi dalam kepekatan dan masa inkubasi yang berlainan semasa pengasingan protoplast. Tisu yang diinkubasikan dalam 1% w/v serbuk *abalone acetone* menghasilkan bilangan protoplast yang paling tinggi (5.0×10^4 protoplast g⁻¹ berat basah). Kombinasi 1% w/v setiap enzim yang tersebut di atas tidak berjaya mencecah bilangan protoplast yang terhasil apabila hanya 1% w/v serbuk *abalone acetone* digunakan.

TABLE OF CONTENTS

| | |
|--------------------|------|
| ACKNOWLEDGEMENTS | II |
| ABSTRACT | III |
| ABSTRAK | V |
| LIST OF TABLES | VII |
| LIST OF FIGURES | VIII |
| LIST OF PLATES | IX |
| LIST OF APPENDICES | XI |

CHAPTER 1 : INTRODUCTION

| | | |
|-------|---|---|
| 1.1 : | Importance of Seaweeds..... | 1 |
| 1.2 : | Economic Importance of Agar and <i>Gracilaria</i> | 3 |
| 1.3 : | Tissue and Protoplast Cultures..... | 5 |
| 1.4 : | Significance of Research..... | 6 |
| 1.5 : | Aims and Objectives of Research..... | 7 |

CHAPTER 2 : LITERATURE REVIEW

| | | |
|---------|---|----|
| 2.1 : | Rhodophyta – The Red Algae..... | 8 |
| 2.2 : | <i>Gracilaria changii</i> Abbott, Zhang and Xia..... | 8 |
| 2.3 : | Conventional Culture Techniques for <i>Gracilaria</i> species.. | 9 |
| 2.4 : | Important Factors Controlling Tissue Culture..... | 10 |
| 2.4.1 : | Surface Sterilisation of Tissue..... | 10 |
| 2.4.2 : | Callus Formation and Frond Regeneration..... | 12 |
| 2.4.3 : | Media and Growth Substances..... | 13 |
| 2.4.4 : | Growth and Maintenance of Culture..... | 16 |
| 2.5 : | Tissue Culture of Seaweeds..... | 17 |

| | | |
|---------|--|----|
| 2.5.1 : | Tissue Culture of <i>Gracilaria</i> | 19 |
| 2.6 : | Important Factors Controlling Protoplast Culture..... | 20 |
| 2.6.1 : | Structure and Organisation of Marine Algal Cell Wall.... | 20 |
| 2.6.2 : | Cell Wall-degrading Enzymes..... | 21 |
| 2.6.3 : | Source of Protoplasts..... | 24 |
| 2.6.4 : | Incubation Conditions During Protoplast Isolation..... | 24 |
| 2.7 : | Protoplast Culture of Seaweeds..... | 26 |
| 2.7.1 : | Protoplast Culture of <i>Gracilaria</i> species..... | 29 |

CHAPTER 3 : MATERIALS AND METHODS

| | | |
|-----------|---|----|
| 3.1 : | <i>Gracilaria changii</i> Abbott, Zhang and Xia | 31 |
| 3.2 : | Collection of <i>Gracilaria</i> | 33 |
| 3.2.1 : | Maintenance of <i>Gracilaria changii</i> in the Laboratory... | 33 |
| 3.3 : | Sterilisation Techniques..... | 34 |
| 3.3.1 : | Treatment 1..... | 34 |
| 3.3.2 : | Treatment 2..... | 34 |
| 3.3.3 : | Treatment 3..... | 34 |
| 3.3.4 : | Treatment 4..... | 35 |
| 3.3.5 : | Treatment 5..... | 35 |
| 3.3.6 : | Treatment 6..... | 35 |
| 3.3.7 : | Sterility Test..... | 36 |
| 3.4 : | Methods of Tissue Culture..... | 37 |
| 3.4.1 : | Source and Preparation of Sterile Seawater..... | 38 |
| 3.4.2 : | Tissue Culture on Solid MS and PES Media..... | 39 |
| 3.4.2.1 : | Murashige and Skoog Basal Salt Medium (MS) | 40 |
| 3.4.2.2 : | MS Basal Medium with Added Vitamins..... | 41 |

| | | |
|-----------|--|----|
| 3.4.2.3 : | Provasoli's Enriched Seawater Medium (PES)..... | 41 |
| 3.4.3 : | Effect of Solid MS and PES media with Growth Substances..... | 42 |
| 3.4.4 : | Tissue Culture in Semi-solid MS and PES Media..... | 43 |
| 3.4.4.1 : | Semi-solid Media with Growth Substances..... | 43 |
| 3.4.5 : | Liquid MS and PES Media..... | 44 |
| 3.4.5.1 : | Liquid Media with Growth Substances..... | 45 |
| 3.5 : | Methods of Protoplast Culture..... | 45 |
| 3.5.1 : | Determination of Mannitol (osmoticum) Concentration..... | 45 |
| 3.5.2 : | Protoplast Isolation..... | 46 |
| 3.5.3 : | Determination of the Effect of Individual Enzyme Concentration and Incubation Time on the Amount of Protoplast Isolated..... | 46 |
| 3.5.4 : | Effect of a Combination of Enzymes on the Amount of Protoplast Isolated at Various Times of Incubation..... | 47 |

CHAPTER 4 : RESULTS

| | | |
|---------|---|----|
| 4.1 : | Sterilisation of Explants..... | 49 |
| 4.2 : | Tissue Culture of <i>Gracilaria changii</i> Abbott, Zhang and Xia..... | 52 |
| 4.2.1 : | Solid Media..... | 52 |
| 4.2.2 : | Semi-solid Media..... | 54 |
| 4.2.3 : | Liquid Media..... | 56 |
| 4.2.4 : | Effects of Liquid Media with Growth Substances..... | 57 |
| 4.3 : | Determination of Optimal Mannitol (osmoticum) Concentration..... | 63 |
| 4.4 : | Protoplast Isolation Using Single Enzyme and Combination of Enzymes..... | 66 |

| | | |
|---------------------------|--|-----|
| CHAPTER 5 : | DISCUSSION | |
| 5.1 : | Sterilisation of <i>Gracilaria changii</i> Explants..... | 72 |
| 5.2 : | Tissue Culture of <i>Gracilaria changii</i> Abbott, Zhang and Xia..... | 74 |
| 5.2.1 : | Effects of the forms of Culture Medium..... | 74 |
| 5.2.2 : | Branching or Shoot Formation..... | 76 |
| 5.2.3 : | Regeneration of Callus-like Structures (CLS)..... | 78 |
| 5.3 : | Optimal Mannitol Concentration..... | 82 |
| 5.4 : | Protoplast Isolation Using Single and Combination of Enzymes..... | 83 |
| CHAPTER 6 : | CONCLUSION | 87 |
| BIBLIOGRAPHY | | 91 |
| APPENDICES | | 100 |

LIST OF TABLES

| | PAGE |
|---|------|
| Table 1. Present market value of seaweeds worldwide..... | 2 |
| Table 2. Sterilisation procedures used in obtaining seaweed tissues... | 11 |
| Table 3. Major components of culture media and their functions..... | 14 |
| Table 4. Growth substances and their functions..... | 15 |
| Table 5. Tissue culture of selected seaweed species..... | 18 |
| Table 6. Natural cell wall degrading enzymes and their organism sources..... | 22 |
| Table 7. Protoplast isolation and culture of selected seaweed species..... | 27 |
| Table 8. Protoplast isolation and culture of <i>Gracilaria</i> spp. | 30 |
| Table 9. Sterilisation of <i>G. changii</i> explants under different treatments..... | 49 |
| Table 10. Culture of explants in liquid PES medium (pH 7.8, 30 ppt salinity) with single growth substance. Each replicate (R) consisted of a culture vessel with five cross-sections of explants inoculated..... | 58 |
| Table 11. Culture of explants in liquid PES medium (pH 7.8, 30 ppt salinity) with combinations of growth substances. Each replicate (R) consisted of a culture vessel with five cross-sections of explants inoculated..... | 62 |
| Table 12. Maximum number of protoplast obtained using each enzyme at the respective isolating conditions..... | 69 |

LIST OF FIGURES

| | PAGE |
|--|------|
| Figure 1. Average percentage of plasmolysed cortical (C) and epidermal (E) cells of the thallus cross-sections in different mannitol concentrations..... | 64 |
| Figure 2. Number of protoplast isolated per gram fresh weight at different time of incubation using 1%, 2% and 3% w/v abalone acetone powder..... | 67 |
| Figure 3. Number of protoplast isolated per gram fresh weight at different time of incubation using 1%, 2% and 3% w/v cellulase..... | 67 |
| Figure 4. Number of protoplast isolated per gram fresh weight at different time of incubation using 1%, 2% and 3% w/v pectinase..... | 68 |
| Figure 5. Number of protoplast isolated per gram fresh weight at different time of incubation using 1%, 2% and 3% w/v pectolyase..... | 68 |
| Figure 6. Number of protoplast isolated per gram fresh weight at different time of incubation using 1%, 2% and 3% w/v agarase..... | 69 |
| Figure 7. Number of viable protoplast isolated per gram of fresh weight at different time of incubation using 1% w/v each of abalone acetone powder, cellulase, pectinase, pectolyase and agarase..... | 71 |

LIST OF PLATES

| | PAGE |
|---|------|
| Plate 1. <i>Gracilaria changii</i> Abbott, Zhang and Xia..... | 32 |
| Plate 2. Collection site at the mangrove area of Pulau Carey..... | 32 |
| Plate 3. Culture of explants in solid media in petri dish..... | 37 |
| Plate 4. "Baby Food Jar" culture vessel used for the culture of explants in semi-solid and liquid media..... | 38 |
| Plate 5. a. Plate of marine agar streaked with thalli from Treatments 3, 4 and 6 on Day 3; b. Plate of marine agar streaked with thalli from Treatments 1, 2 and 5 on Day 3..... | 51 |
| Plate 6. a. Explants on MS and PES solid media on the 28 th day. Complete loss of pigmentation was observed on most of the explants. b. Some explants (10%) managed to survive up to the 56 th days but without callus formation..... | 53 |
| Plate 7. Sample of explants cultured in semi-solid MS, MS + vitamins, MS + vitamins + growth substances, PES + vitamins, PES + vitamins + growth substances media on the 56 th day as shown on a petri dish..... | 55 |
| Plate 8. Sample of explants cultured in liquid MS, MS + vitamins, MS + vitamins + growth substances, PES + vitamins, PES + vitamins + growth substances media on the 56 th day as shown on a petri dish..... | 56 |
| Plate 9. Explants that branched when cultured in liquid PES media containing growth substances..... | 57 |
| Plate 10. a. Ball-like fraible CLS developed from explants in liquid PES media containing a combination of 2,4-D and kinetin. b. CLS as shown on a petri dish..... | 60 |
| Plate 11. a. Cross-section of the CLS under microscope; b. Cells of the CLS..... | 61 |
| Plate 12. a. Cells remained unplasmolysed for mannitol concentrations below 0.7 M. b. Sufficient plasmolysis was induced for cell in 1 M mannitol. c. Over-plasmolysed cells in mannitol concentration above 1.1 M..... | 65 |

Plate 13. Membrane of live protoplast not stained with Evans Blue after isolation..... 70

Plate 14. Membrane of dead protoplast stained with Evans Blue after isolation..... 70

LIST OF APPENDICES

| | PAGE |
|--|------|
| Appendix 1: Percentage of plasmolysed cells in samples of 10 cortical (C) or epidermal (E) cells from the three cross-sections of thallus (T) soaked in seawater (30 ppt) with the respective mannitol concentrations at 30 minutes. Five samples were taken for each cross-section..... | 100 |
| Appendix 2: Effect of time and single enzyme concentrations on protoplast yield..... | 101 |
| Appendix 3: Effect of time and combination of enzymes - 1% each of abalone acetone powder, cellulase, pectinase, pectolyase and agarase on protoplast yield..... | 106 |