

## **REFERENCES:**

Alias, S., Omar, S., Fuad, F. and Sabaratnam,V. (2005). Effect of temperature, media, sea water and pH on the growth of Antarctic fungi. Seminar ‘Antarctic research in the University of Malaya’27-28 June 2005 Kuala Lumpur,Malaysia.

Alexopoulos, C.J. and Beneke, E.S. (1952). Laboratory manual for introductory Mycology. US: Burgess Publishing Company.

Anke, H. and Sterner, O. (2002). Insecticidal and nematicidal metabolites from fungi. *Mycota* 10:109–127.

Azmi, O. and Seppelt, R. (1997). Fungi of the Windmill Islands, Continental Antarctica. Effect of temperature, pH and culture medium on the growth of selected microfungi. *Polar Biol* 18:128–134.

Barker, R.J. (1977). Abiological reconnaissance of the Bunguer Hills. *ANARE Tech Memo* 67:29-30.

Baker, S.J., Hui, X. and Maibach, H.I. (2005). Progress on new therapeutics for fungal nail infections. *AnnuRepMedChem* 40:322–335.

Balkovec, J.M. (1998). Non-azole antifungal agents. *Annu Rep Med Chem* 33:173–182.

Bauer, A., Kirby, W., Sherris, J. and Turck, M. (1966). Antibiotic susceptibility testing by a standardized single disk method. *American Journal of Clinical Pathology* 45: 493–496.

Basile, A., Sorbo, S., Giordano, S., Lavitola, A. and Castaldo Cobianci, R. (1998). Antibacterial activity in *Peurochaete squava* extract (Bryophyta). *International Journal of Antimicrobial Agents* 10: 169-172.

Bell, A.A. and Wheeler, M.H. (1986). Biosynthesis and functions of fungal melanins. *Annu Rev Phytopathol* 24:411-451.

Bills, G.F. (1995). Analysis of microfungal diversity from a user's perspective. *Can J Bot* 73 suppl 1:S33–S41.

Bridge, P.D. and Worland, M.R. (2004). First report of an entomophthoralean fungus on an arthropoda host in Antarctica. *Polar Biol* 27:190-192.

Broady, P.A. (1993). Soil heated by volcanism. In: Friedmann EI(ed) Antarctic microbiology. Wiley-Liss, Inc, New York, pp 413-432.

Broady, P.A. (1996). Diversity, distribution and dispersal of Antarctic terrestrial algae. *Biodivers Conserv* 5:1307-1335.

Brock, T. D. and Madigan, M. T. (1991). Biology of microorganisms. Prentice-Hall International, Inc., Englewood Cliffs N. J. 6<sup>th</sup> ed. p. 874.

Butler, M.J. and Day, A.W. (1998). Fungal melanins: a review. *Can J Microbiol* 44:1115-1113.

Bu'Lock, J. D. (1961). Intermediary metabolism and antibiotic synthesis. *Advances Applied Microbiology* 3:293–342.

Caporale, L. H. (1995). Chemical ecology: a view from the pharmaceutical industry. *Proc Natl Acad Sci USA* 92:75–82.

Caretta, G. and Piontelli, E. (1977). *Microsporium magellanicum* and *Cunninghamella Antarctica*, new species isolated from Australic and Antarctic soils of Chile. *Sabouradia* 15, 1-10.

Caretta, G., Del Frate, G. and Margiarotti, A. (1994). A record of *Arthrobotrys tortor* Jarowaja and *Engyodontium album* (Limber) de Hoog from Antarctica. *Bol Micol* 9:9-13.

Cameron, R.E. (1971). Antarctic soil microbial and ecological investigations. In: Quam LO, Porter HD (eds) Research in the Antarctic. American Association for the Advancement of Science, Washington DC, pp 137-189.

Chakravarti, R. and Sahai, V. (2004). Compactin – a review. *Appl Microbiol Biotechnol* 64:618–24.

Cooke, R.C. and Whipps, J.M. (1993). Ecophysiology of fungi. Oxford, UK: Blackwell Scientific.

Cockell, C. and Blaustein, A. (2001). Ecosystems, evolution and UV radiation. Springer, New York.

Corley, D.G. and Durley, R.C. (1994). Strategies for database dereplication of natural products. *J Nat Prod* 57:1484–1490.

Crowe, J.H., Crowe, L.M., carpenter, J.F. and Wistorn, C.A. (1987). Stabilisation of dry phospholipid bilayers and proteins by sugars, *Biochem J* 242:1-10.

Crowe, J.H., Hoekstra, F.A. and Crowe, L.M. (1992). Anhydrobiosis. *Annu Rev Physiol* 54:579-599.

Cragg, G.M. and Newman, D.J. (2005). Biodiversity: a continuing source of novel drug leads. *Pure Appl Chem* 77:7–24.

Del Frate, G. and Caretta, G. (1990). Fungi isolated from Antarctic material. *Polar Biol* 11:1-7.

De Hoog, G., Gottlich, E., Platas, G., Genilloud, O., Leotta, G. and Van Brummelen, J. (2005). Evolution, taxonomy and ecology of the genus *Thelebolus* in Antarctica. *Stud Mycol* 51:33-76.

Demain, A.L. (1992). Regulation of secondary metabolism. In: Finkelstein DB, Ball C (eds) Biotechnology of filamentous fungi: technology and products. Butterworth- Heinemann, Boston, MA, pp 89–112.

Demain, A.L. (1995). Microbial Secondary Metabolism: The new Frontier. Attempto Verlag Tubingen. Alle Rechte Vorbehalten.

Demain, A.L., Velasco, J.I and Adrio, J.L. (2005). Industrial mycology: past, present, and future. In: AnZ(ed)Mycology Series 22. *Handbook of Industrial Mycology*. Marcel Dekker, New York, pp 1–25.

Dreyfus, M.M. and Chapela, I.H. (1994). Potential of fungi in the discovery of novel low-molecular weight pharmaceuticals. In: Gullo VP (ed) The discovery of natural products with therapeutic potential. Butterworth-Heinemann, Boston, pp 49–80.

Drummond, A.J. and Waigh, R.D. (2000). Recent Research Developments in Phytochemistry. 2000;4:143–152.

Ellis-Evans, J.C. (1985). Microbial ecology in Antarctica. *Biologist* 32:171-176.

Ellen, M., Fox, I., Barbara, J. and Howlett, C. (2008). *Current Opinion in Microbiology*, 11:481–487.

Fenice, M., Selbmann, L., Zucconi, L. and Onofri, S. (1997). Production of extracellular enzymes by Antarctic fungal strains. *Polar Biol* 17:257-280.

Feofilova, E.P., Tereshina, V.M. and Gorova, I.B. (1994). Changes in carbohydrate composition of fungi during adaptation to thermostress. *Microbiology* 64:442-445.

Ferreira, G.A., Schloss, I.R. and Abele, D.I (1999). UV-absorbing compounds in surface waters of Potter Cove: preliminary results. *Berichte zur polarforschung* 299: 316-321.

Finotti, E., Moretto, D., Marsella, R. and Mercantini, R. (1993). Temperature effects and fatty acid patterns in *Geomycetes* species isolated from Antarctic soil. *Polar Biol* 13:127-130.

Foissner, W. (2006). Biogeography and dispersal of microorganisms: a review emphasizing protists. *Acta Protozool* 45:111-136.

Finotti, E., Paolino, C., Lancia, B. and Mercantini, R. (1996). Metabolic Differences between Two Antarctic Strains of *Geomycetes pannorum*. *Current Microbiology*. **32**:7–10.

Fletcher, L.D., Kerry, E.J. and Weste, G.M. (1985). Microfungi of Mac. Robertson and Enderby Lands, Antarctica, *Polar Biol* 4:81-92.

Friedmann, E.I. (1993). Antarctic microbiology. Wiley-Liss, New York, 634pp.

French, G.L. ( 2006). "Bactericidal agents in the treatment of MRSA infections-the potential role of daptomycin". *J. Antimicrob. Chemother.* **58** (6): 1107–17.

Gamundi, I.J. and Spinedi, H.A. (1987). *Sclerotinia Antarctica* sp.nov., the teleomorph of the first fungus described from Antarctica. *Mycotaxon*, 29:81-89.

Ghorbani, R., Leifert, C. and Seel, W. (2005). Biological control of weeds with antagonistic plant pathogens. *Adv Agron* 86:191–225.

Gloer, J.B. (2007). Applications of fungal ecology in the search for new bioactive natural products. In C.P.Kubicek and I.S.Druzhinina (eds) The Mycota vol IV, 2nd edn. Environmental and microbial relationships. Springer-Verlag Berlin Heidelberg, pp 257–275.

Gounot, A. M. (1991). Bacterial life at low temperature: physiological aspects and biotechnological aspects and biotechnological implications. *J of App Bactel* **71**, 386-397.

Goodrich, R.P., Handel, T.M. and Baldeschwieler, J.D. (1988). Modification of lipid phase behaviour with membrane-bound cryoprotectants. *Biochimica et Biophysica Acta* **938**: 143–154.

Gorbushina, A., Krumbein, W., Panin, L., Soukharjevsky, S., Wollenzien, U. and Hamann, K. (1993). On the role of black fungi in colour change and biodeterioration of antique marbles. *Geomicrobiol J* **11**:205-221

Gorbushina, A., Beck, A. and Schulte, A. (2005). Microcolonial rock inhabiting fungi and lichen photobionts: evidence for mutualistic interactions. *Mycol Res* **109**(11):1288-1296.

Gould, D. and Booker, C. (2000). Applied microbiology for nurses. Aardvark Editorial, Mcndham, Suffolk. pp. 75-94.

Grant, W.D. (2004). Life at low water activity. *Phil Trans R Soc London B* **359**:1249-1267.

Gunde-Cimerman, N., Sonjak, S., Zalar, P., Frisvad, J., Didderichsen, B. and Plemenitas, A. (2003). Extremophilic fungi in arctic ice: relationship between adaptation to low temperature and water activity. *Phy Chem Earth* **28**:1273-1278.

Hawkey, B. M. (1998). The origins and molecular basis of antibiotic resistance. *BMJ Journal* **317**: 657-60.

Helmholz, H., Etoundi, P. and Lindequist, U. (1998). Cultivation of marine basidiomycete *Nia vibrissae* (Moore & Meyers). *Journal of Biotechnology*, 70,203-206.

Holdgate, M.W. (1977). Terrestrial ecosystems in the Antarctic. *Philosophical Transactions of the royal society B*. 279:5-25.

Jennings D.H. (1984). Polyol metabolism in fungi. *Advances in Microbial.*

Jennings, D.H. (1995). The physiology of fungal nutrition. Cambridge Univ. Press, Cambridge.

Jorgensen, J. H., Turnidge, J. D. and Washington, J. A. (1999). Antibacterial susceptibility test: Dilution and disk diffusion method. In: (Murray, P. R., Baron, E. J., Pfaller, M. A, Ternover, F. C. and Yolken, R. H. eds.). *Manual of Clinical Microbiology* 7<sup>th</sup> Edition Washington. ASM Press p. 1526-1543.

Jumpponen, A., Newsham K. and Neises, D. (2003). Filamentous ascomycetes inhabiting the rhizoid environment of the liverwort *Cephaloziella varians* in Antarctica are assessed by direct PCR and cloning. *Mycologia* 95:457-466.

Kappen, L. (1993). Lichens in Antarctic region. In: Friedmann EI (ed) *Antarctic Microbiology*. Wiley Liss, New York, pp433-490.

Karentz, D. (1994). Ultraviolet tolerance mechanisms in Antarctic marine organisms. In: Weiler CS, Penhale PA (eds) Ultraviolet radiation in Antarctica: measurements and biological effects. American Geophysical Union, Washington, DC, pp 92-110.

Kerry, E. (1990). Effects of temperature on growth rates of fungi from subantarctic Macquarie Island and Casey, Antarctica. *Polar Biol.* 10:293–299.

Kerry, E. (1990). Microorganisms colonizing plants and soil subjected to different degree of human activity, including petroleum contamination in the Vestfold Hills and Mac. Robertson Land, Antarctica. *Polar Biol* 10:423-430.

Kogej, T., Wheeler, M., Rizner, T. and Gunde-Cimerman, N. (2004). Evidence for 1,8-dihydroxynaphthalene melanin in three halophilic black yeasts grown under saline and non-saline conditions. *FEMS Microbiol Lett* 232: 203-209.

Koltin, Y. (1990). Targets for antifungal drug discovery. *Annu Rep Med Chem* 25:141–148.

Kumar, V., Chauhan, N., Padh, H. and Rajani, M. (2006). Search for antibacterial and antifungal agents from selected Indian medicinal plants, *J. of Ethnopharmacology* **107**: 182–188.

Liu, X., Li, S. (2005). Fungal secondary metabolites in biological control of crop pests. In: An Z (ed)Mycology Series 22. Handbookof Industrial Mycology. MarcelDekker,New York, pp 723–747.

Leotta, G., Pare, J., Sigler, L., Montalti, D., Vigo, G., Petruccelli, M. and Reinoso, E. (2002). *Thelebolus microsporus* mycelial mats in the trachea of wild brown skua (*Catharacta antarctica lonnbergi*) and south polar skua ( *C. maccormicki*) carcasses. *J Wildlife Dis* 38 (2):443-447.

Lewis Smith, R.I. (1984). Terrestrial plant biology of the sub-Antarctic and Antarctic. In:Laws R.M., 8<sup>th</sup> edition, *Antarctic Ecology* 1:61-162. Academic press, London.

Lyutskanova, D., Ivanova, V., Stoilova-Disheva, M., Kolarova, M., Aleksieva, K., Raykovska, V. and Peltekova, V. (2009). Isolation, characterization and screening for antimicrobial activities of psychrotolerant *streptomyces* isolated from polar permafrost soil. Bulgaria Academy of Sciences, The stefan angeloff institute of microbiology, Acad, G. Bonchev str. B1.26,1113 Sofia, Bulgaria.

Maggi, O., Persiani, A., Fabbri, A., De Luca, C., Lunghini, D., Quadraccia, L. and Fanelli, C. (1991). Differenze nella comoposizione fosfolipidica di diversi taxa fungini. Giorn Bot Ital 125(3):259.

Marshall, W.A. (1998). Aerial transport of keratinaceous substrate and distribution of the fungus *Geomycetes pannorum* in Antarctic soils. *Microbial Ecology* **36**: 212–219.

Masurekar, P.S. (2005). Strain improvement for the production of fungal secondary metabolites. In: An Z (ed) Mycology Series 22. Handbook of Industrial Mycology. Marcel Dekker, New York, pp 539–561.

Marinelli, F., Brunati, M., Songa, F., Ciciliato, I., Losi, D., van Trappen, S., Mergaet, J., Swings, J., Gottlich, E., Hoog, G., Rojas, J. and Genilloud, O. (2004). Biotechnological exploitation of heterotrophic bacteria and filamentous fungi isolated from benthic mats of Antarctic lakes. In:Kurtboke I & Swings J.(eds), Microbial genetic resources and biodiscovery : 163-184.

Mercantini, R., Maarsella, R., Moretto, D. and Finotti E. (1993). Keratinophilic fungi in the Antarctic environment. *Mycopathologia* 122:169-175.

Miller, J.D. (1991). Mycology, mycologists, and biotechnology. In: Hawksworth DL, Regensburg CA (eds). *Frontiers in mycology. Honorary lectures 4th Int Mycological Congr.* CABI, Wallingford, pp 225–240.

Miao, L., Theresa, F. and Qian, P. (2006). Effect of culture conditions on mycelial growth, antibacterial activity, and metabolite profiles of the marine-derived fungus *Arthrinium* c.f. *saccharicola*. *Appl Microb and Cell Physiology* 72: 1063-1073.

Morita, R. Y. (1975). Psychrophilic bacteria. *Bacteriol. Rev.* **39**, 144–167.

Montermartini Corte, A. (1991). Fungi di ambienti acquatici. In: Battaglia B, Bisol PM, Varotto V (eds) proceeding of the 1st meeting on Biology in Antarctica (English summaries). Roma CNR 22-23 June 1989, Scienza e Culture, Edizioni Universitarie Patavine, pp 67-76.

Montermartini Corte, A., Caretta, G. and Del Frate, G. (1993). Notes on Thelebolus microspores isolated in Antarctica. *Mycotaxon* 48:343-358.

Montermartini Corte, A. and Gestro, C. (1994). Fungi from mud of Antarctic ponds. In: Battaglia, B, Bisol, PM. and Varotto, V. (eds) Scienza e cultura. Proceeding of the 2<sup>nd</sup> meeting on Antarctic Biology, Padova, 26-28 February 1992. Edizioni Universitarie Patavine, Italy, pp 33-42.

Montermartini, A., Liotta, M., Venturi, C. and Calegari, L. (2000). Antibacterial activity of *Penicillium* spp. Strains isolated in extreme environments. *Polar Biol* 23(4), 294-296. Retrieved June 10, 2008, from the springer link database.

Montiel, P.O. (2000). Solubile carbohydrates (Trehalose in particular) and cryoprotection in polar biota. *Cryo-Lett* 21:83-90.

Monaghan R.L. and Tkacz, J.S. (1990). Bioactive microbial products: focus upon mechanism of action. *Annu Rev Microbiol* 44:271–301.

Moss, M. O. (1984). The mycelial habit and secondary metabolite production. In: *Advances in Applied Microbiology: The Ecology and Physiology of the fungal Mycelium* at Symposium of the Brittish Mycological Society, Bath University (Jennings, D. H. and Rayno, A. D. M, eds). Cambridge University Press 34: 127-141.

NCCLS. (2000). Method for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Approved standard M7-A5.NCCLS,Wayne,Pa.

Newman D.J., Cragg, G.M. and Snader, K.M. (2000). The influence of natural products upon drug discovery. *Nat Prod Rep* 17:215–234.

Nedialkova Denitsa and Mariana Naidenova. (2004). Screening the antimicrobial activity of *Actinomycetes* strains isolated from Antarctica. *J. of cult collections*, Volume 4, 2004-2005, pp. 29-39.

Nienow, J.A. and Friedmann, E.I. (1993). Terrestrial lithophytic (rock) communities. In: Friedmann, E.I. (Ed.), Antarctic Microbiology. Wiley-Liss, New York, pp. 343–412.

Niederer, M., Pankow, W. and Wiemken, A. (1992). Seasonal changes of soluble carbohydrates in mycorrhizas of Norway spruce and changes induced by exposure to frost desiccation. *Eur. J. of Forest Pathology* **22**: 291–299.

Oh, D. C., Jensen, P. R., Kauffman, C. A. and Fenical, W.I. (2005). Libertellenones A-D: Induction of cytotoxic diterpenoid biosynthesis by marine microbial competition. *Bioorganic and Medicinal Chemistry* **13**: 5267-5273.

Onofri, S., Rambelli, A., Maggi, O., Persiani, A., Riess, S., Tosi, S. and Grasselli, E. (1991). Mycologia del suolo. In: Battaglia B, Bisol PM, Varotto V (eds) proceeding of the 1st meeting on Biology in Antarctica (English summaries). Roma CNR 22-23 June 1989, Scienza e Culture, Edizioni Universitarie Patavine, pp 55-65.

Onofri, S. and Tosi, S. (1992). *Anthrobotrys ferox* sp. Nov. asprigtail-capturing hyphomycete from continental Antarctica. *Mycotaxon* **44**:445-451.

Onofri, S., Tosi, S., Persiani, A., Maggi, O., Riess, S. and Zucconi, L. (1994). Mycological research in Victoria Land terrestrial ecosystem. Proceeding of the 2<sup>nd</sup> meeting on Antarctic Biology, Padova, 26-28 February 1992. Scienza e Culture Edizioni Universitarie Patavine, padova, pp 19-32.

Onofri, S. (1999). Antarctic microfungi. In:Seckbach, J.(Ed.), Enigmatic Microorganisms and Life in Extreme Environments. Kluwer Academic\Publishers, Dordrecht, Boston, London, pp. 323–336.

Onofri, S., Fenice, M., Cicalini, A., Tosi, S., Magrino, A., Pagano, S., Selbmann, L., Zucconi, L., Vishniac, H., Ocampo-Friedmann, R. and Friedmann, E. (2000). Ecology and biology of microfungi from Antarctic rocks and soil. *Italy J ZOOL* 67 (suppl. 1):163-168.

Onofri, S., Selbmann, L., Zucconi, L., Fenice, M., Barreca, D. and Ruisi, S. (2005).Studies on Antarctic fungi. *Polar net Tech Rep.*1:49-52.

Onofri, S., Zucconi, L. and Tosi, S. (2007). Continental Antarctic fungi. I HW-Verlag, Eching (in press).

Øvstedal, D.O. and Lewis Smith, R.I. (2001).Lichens of Antarctica and South Georgia. *A guide to their identification and ecology.* Studies in polar research. Cambridge University Press, Cambridge, UK.

Patrick, R. and Murray. A. (2003). Manual of clinical microbiology; editors, Ellen Jo Baron.(et al)-8<sup>th</sup> ed.ASM press p.1120-1121.

PetroviVc, U., Gunde-Cimerman, N. and Zalar, P. (2000). Xerotolerant mycobiota from high altitude Anapurna soil, Nepal. *FEMS Microbiol. Lett.* 182, 339–342.

Pickard, J. and Seppelt, R. (1984). Phytogeography of Antarctica. *J Biogeogr* 11:83-102

Pugh, G. and Allsop, D. (1982). Microfungi on Signy Island, South Orkney Island. *Br Antarct Surv Bull* **57**, 55-67.

Purvis, W. (2000). *Lichens*. Smithsonian Institution Press, Washington, D.C., 112pp.

Rivkina, E.P., Friedmann, E.I., McKay, C.P. and Gilichinsky, D.A. (2000). Metabolic activity of permafrost bacteria below the freezing point. *Appl Environ. Microbiol* 66:3230-3233.

Richardson, K. and Marriott, S. (1987). Antifungal agents. *Ann Rep Med Chem* 22:159–167.

Rounsevell, D.E. (1981). A population of *Nanochestes antarcticus* (Acari: Prostigmata) at the Vestfold Hills, Antarctica. *ANARE Sci. Rep. Ser. B* (1) Zool 131:100.

Robinson, H. (2001). Cold adaptation in Arctic and Antarctic fungi. Division of life sciences, King's college, University of London.

Rusisi S., Barreca D., Selbmann L., Zucconi, L. and Onofr, S. (2007). Fungi in Antarctica .*Rev Environ Sci Biotechnology* 6:127-141.

Russell, N.J. (1990). Cold adaptation of microorganisms. *Philos Trans Roy Soc London Ser B* 326:595-611.

Ribal, C., Platas, G. and Bills, G. (2005). Isolation and characterization of melanized fungi from limestone in Mallorca. *Mycol Progr* 4: 23-38

Samson, R.A., Hoekstra, E.S., Frisvad, J.C. and Filtenborg, O. (1995). Introduction to Food-borne Fungi. CBS, Baarn, the Netherlands.

Selbmann, L., Onfori, S., Fenice, M., Federici, F. and Petruccioli, M. (2002). Production and structural characterization of the exopolysaccharide of the Antarctic fungus *Phoma herbarum* CCFEE 5080. *Rec Microniol* 153:585-592.

Selbmann, L., de Hoog, G., Mazzaglia A, Friedmann, E. and Onofri, S. (2005). Fungi at the edge of life: cryptoendolithic black fungi from Antarctic desert. *Stud Mycol* 51:1-32.

Shick, J.M. and Dunlap, W.C. ( 2002). Mycosporine-like amino acids and related gadusols: biosynthesis, accumulation, and UV-protective functions in aquatic organisms. *Annu Rev of Physiology* 64: 223-262.

Singh, S.B., Pelaez, F.A., Hazuda, D.J. and Lingham, R.B. (2005). Discovery of natural product inhibitors of HIV-1 integrase at Merck. *Drugs Future* 30:277–299.

Sinclair, N.A. and Stokes, J.L. (1965). Obligatory psychrophilic yeasts from the Polar Regions. *C. J. microbiology* 11:259-269.

Sterflinger, K. de Hoog G. and Haase, G. (1999). Phylogeny and ecology of meristematic ascomycetes. *Stud Mycol* 43: 5-22.

Sterflinger, K. (2005). Black yeast and meristematic fungi:ecology, diversity and identification. In: Rosa C, Gabor P (eds) Yeast handbook: biodiversity and ecophysiology of yeasts. Springer, New York, pp 505-518.

Strongman, D.B., Miller, J.D., Calhoun, L., Findlay, J.A. and Whitney, N.J. (1987) The biochemical basis for interference competition among some lignicolous marine fungi. *Bot Mar* 30:21–26.

Thevelein, J.M. (1984). Regulation of trehalose mobilisation in fungi. *Microbiological Reviews* 48: 42–59.

Tosi, S., Begona, C., Gerdol, R. and Caretta, G. (2002). Fungi isolated from Antarctic mosses. *Polar Biol* 25:262-268.

Tubak, K. (1961). On some fungi isolated from the Antarctic materials. *Biol Results JARE*, No 14, pp9.

Tubaki, K. and Asano, I. (1965). Additional species of fungi isolated from the Antarctic materials. *JARE 1956-62 Sci Rep, Ser E*, No 27, pp 1-12.

Vishniac, H.S. (1993). The microbiology of Antarctic soils. In: Friedmann EI (ed) Antarctic microbiology. Wiley, New York, pp 297-341.

Vishniac, H. and Onfri, S. (2002). *Cryptococcus antarcticus* var. *circumpolaris* var. nov, a basidiomycetous yeast from Antarctica. *Antonie van Leeuwenhoek* 83:233-235.

Vincent, W.F. (2000). Evolutionary origins of Antarctic microbiota: invasion, selection and endemism. *Antarct Sci* 12(3): 374-385.

Volkmann, M., Whitehead, K., Ruttgers, H., Rullkotter, J. and Gorbushina, A. (2003).

Mycosporine-glutamicol-glucoside: a natural UV absorbing secondary metabolite of rock-inhabiting microclonal fungi. *Rapid Comm Mass Spectr* 17:897-902.

Weinstein, R., Montiel, P. and Johnstone, K. (2000). Influence of growth temperature on lipid and soluble carbohydrate synthesis by fungi isolated from fellfield soil in the maritime Antarctic. *Mycologia* 92:222-229.

Wicklow, D.T. (1968). Aspergillus fumigatus Fresenius isolated from ornithogenic soil collected at Hallet Station, Antarctica. *Can. J. Microbiol.* 14:717-719.

Wong-Beringer, A. and Kriengkauykiat, J. (2003). Systemic antifungal therapy: new options, new challenges. *Pharmacotherapy* 23:1441–1462.

Wynn-Williams, D.D. (1990). Ecological aspects of Antarctic microbiology. *Advan in Microbial Ecology* 11: 71-146.

Wynn-Williams, D.D. and Edwards, H.G. (2001). In: Horneck G, Baumstark-Khan C (eds) Environmental UV radiation: biological strategies for protection and avoidance, inastrobiology: the quest for the conditions of life. Springer-Verlag, Berlin, pp 244-259.

Zucconi, L., Pagano, S., Fenice, M., Selbmann, L., Tosi, S. and Onofri, S. (1996). Growth temperature preferences of fungal strains from Victoria Land. Antarctica. *Polar Biol.* 16, 53–