

REFERENCES

- Abbott, I.A., Zhang, C.F. and Xia, B.M. (1991) *Gracilaria mixta*, sp. nov. and other Western Pacific species of the genus Rhodophyta: Gracilariaceae. **Pacific Science** 45(1):12-27.
- Alberte, R.S., Suba, G.K., Procaccini, G.A., Zimmerman, R.C. and Fain, S.R. (1994). Assessment of genetic diversity of eel grass (*Zostera marina* L.) populations using DNA fingerprinting : implications for population stability and management. **Proc. Nat. Acad. Sci. USA** 91:1049-1053.
- Alberto, F., Santos, R. and Leitao, J.M. (1997) DNA extraction and RAPD markers to assess the genetic similarity among *Gelidium sesquipedale* (Rhodophyta) populations. **J. Phycol** 33 : 706-710.
- Allen, R.C., Graves, G. and Budowle, B. (1989). Polymerase chain reaction amplification products separated on rehydratable polyacrylamide gels and stained with silver. **Biotechniques** 7:736-744.
- Armisen, R. (1995). World-wide use and importance of *Gracilaria*. **Journal of Applied Phycology** 7:231-243.
- Armisen, R., and Galatas, F. (1987). Production, properties and uses of agar. **In:** D.J. McHugh (ed.). **Production and Utilisation of Products from Commercial Seaweeds**. FAO Fish. Tech. Pap. 288. Food and Agriculture Organisation of the United Nations, Rome. pp. 1-57.
- Bakker, F.T., Olsen, J.L., Stam, W.T. and Van der Hoek, C. (1992). Nuclear ribosomal DNA internal transcribed spacer regions (ITS1 and ITS2) define discrete biogeographic groups in *Cladophora albida* (Chlorophyta). **J. Phycol.** 28:839-845.
- Baril, C.P., Verhaegen, D., Vigneron, P., Bauret, J.M. and Kremer, A. (1997). Structure of the specific combining ability between two species of *Eucalyptus* I. RAPD Data. **Theor. Appl. Genet.** 94:796-803.
- Batchvarov, S. (1993). Garlic *Allium sativum* L. **In:** G. Kalloo and B.O. Bergh (eds.). **Genetic Improvement of Vegetable Crops**. Pergamon Press, New York. pp. 15-27.
- Besnard, G., Griveau, Y., Quillet, M.C., Serieys, H., Lambert, P., Vares, D. and Berville, A. (1997). Specifying the introgressed regions from *H. argophyllus* in cultivated sunflower (*Helianthus annuus* L.) to mark phomopsis resistance genes. **Theor. Appl. Genet.** 94:131-138.

- Bhattacharya, D. and Druehl, L.D. (1990). Restriction enzyme analysis of variation and taxonomy in the kelp genus *Laminaria* (Laminariales, Phaeophyta). **Hydrobiologia** 204/205:105-110.
- Bhattacharya, D., Elwood, H.J., Goff, L.J. and Sogin, M.L. (1990). Phylogeny of *Gracilaria lemaneiformis* (Rhodophyta) based on sequence analysis of ITS small subunit ribosomal RNA coding region. **J. Phycol.** 26:181-186.
- Bird, C.J. and McLachlan, J. (1982). Some underutilized taxonomic criteria in *Gracilaria* (Rhodophyta, Gigartinales). **Bot. Mar.** 25:557-562.
- Bird, C.J., and McLachlan, J. (1984). Taxonomy of *Gracilaria* : Evaluation of some aspects of reproductive structure. **Hydrobiologia** 115/116:41-62.
- Bird, C.J., Murphy, C.A., Rice, E.L. and Ragan, M.A. (1992a). The 18S rRNA gene sequence of four commercially important seaweeds. **Journal of Applied Phycology** 4:379-384.
- Bird, C.J. and Rice, E.L. (1990) Recent approaches to the taxonomy of the Gracilariaceae (Gracilariales, Rhodophyta) and the *Gracilaria verrucosa* problem. **Proc. Int. Seaweed Symp.** 13 :111-118.
- Bird, C.J., Rice, E.L., Murphy, C.A., Liu, Q.Y. and Ragan, M.A. (1990a). Nucleotide sequences of 18S ribosomal RNA genes from the red algae *Gracilaria tikvahiae* McLachlan, *Gracilaria verrucosa* (Hudson) Papenfuss and *Gracilariopsis* sp. **Nucleic Acids Res.** 18(13):4023-4024.
- Bird, C.J., Rice, E.L., Murphy, C.A. and Ragan, M.A. (1992b). Phylogenetic relationships in the Gracilariales (Rhodophyta) as determined by 18S rDNA sequences. **Phycologia** 31(6) :510-522.
- Bird, C.J., Nelson, W.A., Rice, E.L. and Ragan, M.A. (1990b). A critical comparison of *Gracilaria chilensis* and *G. sordida* (Rhodophyta, Gracilariales). **Journal of Applied Phycology** 2:375-382.
- Bird, C.J., Ragan, M.A., Critchley, A.T., Rice, E.L. and Gutell, R.R. (1994). Molecular relationships among the Gracilariaceae (Rhodophyta) : further observations on some undetermined species. **Eur. J. Phycol.** 29 :195-202.
- Bird, C.J. and Van der Meer, J.P. (1993). Systematics of economically important marine algae: A Canadian perspective. **Can. J. Bot.** 71:361-369.
- Bird, K.T., Hanisak, M.D. and Ryther, J.H. (1981a). Chemical quality and production of agars extracted from *Gracilaria tikvahiae* grown in different nitrogen enrichment conditions. **Bot. Mar.** 24:441-444.

- Bird, K.T., Hanisak, M.D. and Ryther, J.H. (1981b). Changes in agar and other chemical constituents of the seaweed, *Gracilaria tikvahiae*, when used as a substrate in methane digesters. **Resour. Conserv.** 6 : 321-327.
- Bommineni, V.R., Jauhar, P.P., Peterson, T.S., Chibbar, R.N. and Almouslem, A.B. (1997). Analysis of hybrids of durum wheat with *Thinopyrum junceiforme* using RAPD markers. **Theor. Appl. Genet.** 95 :757-763.
- Brasch, D.J., Chang, H.M., Chuah, C.T. and Melton, L.D. (1981). The galactan sulfate from the edible, red alga *Porphyra columbina*. **Carbohydrate Res.** 97:113-125.
- Brown, L.M. and McLachlan, J. (1982). A typical carotenoid for the Rhodophyceae in the genus *Gracilaria* (Gigartinales). **Phycologia** 21:9-16.
- Buchheim, M.A. and Chapman, R.L. (1991). Phylogeny of the colonial green flagellates: a study of 18S and 26S rRNA sequence data. **Biosystems** 25: 85-100.
- Caetano-Anolles, G., Bassam, B.J., and Gresshoff, P.M. (1991). DNA amplification using very short arbitrary oligonucleotide primers. **Biotechnology** 9:553-557.
- Chan, K.F. and Sun, M. (1997). Genetic diversity and relationships detected by isozyme and RAPD analysis of crop and wild species of *Amaranthus*. **Theor. Appl. Genet.** 95:865-873.
- Chang, C.F. and Xia, B.M. (1963). *Polycavernosa*, a new genus of the *Gracilariaceae*. **Studies Marina Sinica** 3:119-126.
- Chang, C.F. and Xia, B.M. (1976). Studies on Chinese species of *Gracilaria*. **Studies Marina Sinica** 11: 157-163.
- Chapman, R.L. and Buchheim, M.A. (1992). Green algae and the evolution of land plants: inferences from nuclear-encoded rRNA gene sequences. **Biosystems** 28:127-137.
- Chauvat, F., Labarne, J., Ferino, F., Thuriaux, P. and Fromageot, P. (1988). gene transfer to the *Cyanobacterium synechocystis* PCC 6803. **In** : T. Stadler, J. Mollion, M.C. Verdu, Y. Karamanos, H. Morvan and D. Christiaen (eds.). **Algal Biotechnology** Elsevier Applied Science. New York. pp. 89-98.
- Chen, L.F.O., Kuo, H.Y., Chen, M.H., Lai, K.N. and Chen, S.C.G. (1997). Reproducibility of the differential amplification between leaf and root DNAs in soybean revealed by RAPD markers. **Theor. Appl. Genet.** 95:1033-1043.
- Chiang, M.S., Chong, C., Landry, B.S. and Crete, R. (1993). Cabbage *Brassica oleracea* subsp. *capitata* L. **In** : G. Kalloo and B.O. Bergh (eds.). **Genetic Improvement of Vegetable Crops.** Pergamon Press, New York. pp. 113-155.

- Chirapart, A. and Ohno, M. (1993). Seasonal variation in the physical properties of agar and biomass of *Gracilaria* sp. (chorda type) from Tosa Bay, Southern Japan. **Hydrobiologia** 260/261:541-547.
- Coat, G., Dion, P., Noailles, M.C., De Reviers, B., Fontaine, J.M., Berger-Perrot, Y. and Loiseaux-De Boer, S. (1998). *Ulva armoricana* (Ulvales, Chlorophyta) from the coasts of Brittany (France) II. Nuclear. DNA ITS sequence analysis. **Eur. J. Phycol.** 33: 81-86.
- Coleman, A.W. and Goff, L.J. (1991). DNA analysis of eukaryotic algal species. **J. Phycol.** 27:463-473.
- Coyer, J.A., Olsen, J.L. and Stam, W.T. (1997). Genetic variability and spatial separation in the sea palm kelp *Postelsia palmaeformis* (Phaeophyceae) as assessed with M13 fingerprints and RAPDs. **J. Phycol.** 33:561-568.
- Coyer, J.A., Robertson, D.L. and Alberte, R.S. (1994). Genetic variability within a population and between diploid/haploid tissue of *Macrocystis pyrifera* (Phaeophyceae). **J. Phycol.** 30 : 545-552.
- Coyer, J.A. and Steller, D.L. (1995) A field-compatible method for extraction of fingerprint-quality DNA from *Macrocystis pyrifera* (Phaeophyceae). **J. Phycol** 31: 177-180.
- Craigie, J.S., Wen, Z.C. and Van der Meer, J.P. (1984). Interspecific, intraspecific and nutritionally-determined variations in the composition of agars from *Gracilaria* sp. **Bot. Mar.** 27:55-61.
- Crisp, P. and Tapsell, C.R. (1993). Cauliflower *Brassica oleracea* L. **In:** G. Kalloo and B.O. Bergh (eds.), **Genetic Improvement of Vegetable Crops**. Pergamon Press, New York. pp. 157-178.
- Critchley, A.T. (1993). Introduction : seaweed resources. **In:** M. Ohno and A. T. Critchley (eds.), **Seaweed Cultivation and Marine Ranching**. Japan International Fisheries Training Center, Japan International Cooperation Agency. pp.1-6.
- Crueger, W. and Crueger, A. (1989). **In:** D.B. Thomas (ed.), **Biotechnology: A textbook of industrial microbiology**. Science Tech. Publishers, USA.

- Curtsinger, J.W., Fukui, H.H., Resler, A.S., Kelly, K. and Khazaeli, A.A. (1998). Genetic analysis of extended life span in *Drosophila melanogaster* I. RAPD screen for genetic divergence between selected and control lines. **Genetica** 104:21-32.
- Dawson, E.Y. (1949). Studies on the Northeast Pacific *Gracilariaceae*. Allan Hancock Found. Publ., **Occ. Papers** 7:1-54. **In** : Federicq, S. and Norris, J. N. (1985). Morphological studies on some tropical species of *Gracilaria* Grev. (*Gracilariaceae*, Rhodophyta): Taxonomic concepts based on reproductive morphology. **In** : I. A. Abbott and J. N. Norris (eds.). **Taxonomy of economic seaweeds-with references to some Pacific and Caribbean species**. California Sea Grant College, La Jolla, California. pp. 137-155.
- Dawson, E.Y. (1954). Notes on tropical Pacific marine algae. **Bull. So. Calif. Acad. Sci.** 53: 1-7.
- Dehghan-shoar, M., Hampton, J.G. and Gardiner, S.E. (1997). Genetic analysis among and within populations forming ecotypes and cultivars of Lucerne, *Medicago sativa* (Leguminosae), using RAPD fragments. **Plant Systematics and Evolution** 208: 107-119.
- De Roeck-holtzhauer, Y (1991). Uses of seaweeds in cosmetics. **In**: M.D. Guiry and G. Blunden (eds.). **Seaweed Resources in Europe: Uses and Potential**. John Wiley and Sons, Chichester, England. pp. 83-93.
- ✓ Doyle, J.J. and Dickson, E.E. (1987). Preservation of plant samples for DNA restriction endonuclease analysis. **Taxon** 36: 715-722.
- Doyle, J.J. and Doyle, J.L. (1990) Isolation of plants DNA from fresh tissue. **Focus** 12(1): 13-15.
- Dubouzet, J.G., Shinoda, K. and Murata, N. (1997). Phylogeny of *Allium* L. subgenus *Rhizirideum* (G. Don ex Koch) Wendelbo according to dot blot hybridization with randomly amplified DNA probes. **Theor. Appl. genet.** 95: 1223-1228.
- Dutcher, J.A., Kapraun, D.F. and Sizemore, R.K. (1990). Inter- and intraspecific variation of nuclear DNA reassociation kinetics in the Gracilariales (Rhodophyta). **Journal of Applied Phycology** 2: 259-267.
- Dutcher, J.A. and Kapraun, D.F. (1994). Random amplified polymorphic DNA (RAPD) identification of genetic variation in three species of *Porphyra* (Bangiales, Rhodophyta). **Journal of Applied Phycology** 6:267-273.
- Dweikat, I., Ohm, H., Patterson, F. and Cambron, S. (1997). Identification of RAPD markers for 11 Hessian fly resistance genes in wheat. **Theor. Appl. Genet.** 94:419-423.

- Emebiri, L.C., Devey, M.E., Matheson, A.C. and Slee M.U. (1997). Linkage of RAPD markers to NESTUR, a stem growth index in radiata pine seedlings. **Theor. Appl. genet.** 95:119-124.
- Fain, S.R., Druehl, L.D. and Braille, D.L. (1988) Repeat and single copy sequences are differentially conserved in the evolution of Kelp chloroplast DNA. **J. Phycol.** 24:292-302.
- Fredericq, S. and Hommersand, M.H. (1989). Comparative morphology and taxonomic status of *Gracilariopsis* (Gracilariales, Rhodophyta). **J. Phycol.** 25:228-241.
- Fredericq, S. and Norris, J.N. (1985). Morphological studies on some tropical species of *Gracilaria* Grev. (Gracilariaceae, Rhodophyta): taxonomic concepts based on reproductive morphology. **In**: I. A. Abbott and J.N. Norris (eds.). **Taxonomy of economic seaweeds with reference to some Pacific and Caribbean species.** California Sea Grant College Program, La Jolla C.A. pp.137-155.
- Freshwater, D.W., Fredericq, S., Butler, B.S., Hommersand, M.H. and Chase, M.W. (1994). A gene phylogeny of the red algae (Rhodophyta) based on plastid-encoded rbcL. **Proc. Nat. Acad. Sci. USA** 91: 7281-7285.
- Fujimura, T. and Kajiwara, T. (1990). Production of bioflavor by regeneration from protoplasts of *Ulva pertusa* (Ulvales, Chlorophyta). **Hydrobiologia** 204/205:143-149.
- Fujita, Y. and Migita, S. (1987). Fusion of protoplasts from thalli of two different colour types in *Porphyra yezoensis* and development of fusion products. **Japan J. Phycol.** 35 : 201-208.
- Gaiotto, F.A., Bramucci, M. and Grattapaglia, D. (1997). Estimation of outcrossing rate in a breeding population of *Eucalyptus urophylla* with dominant RAPD and AFLP markers. **Theor. Appl. Genet.** 95: 842-849.
- Ganesan, E.K. (1993). Morphological and taxonomic studies on some little-known species of *Gracilaria* (Gracilariales: Rhodophyta) from Venezuela. I. *G. damaecornis*. **Hydrobiologia** 260/261:91-96.
- Gargiulo, G.M., De Masi, F. and Tripodi, G. (1992). Morphology, reproduction and taxonomy of the mediterranean species of *Gracilaria* (Gracilariales, Rhodophyta). **Phycologia** 31:53-80.
- Gavino, C.T. (1997). Rhodophyceae. **In**: **Field guide and atlas of the seaweed resources of the Philippines.** Bookmark, Inc. Philippines. pp. 158-271.

- Gehrig, H.H., Rosicke, H. and Kluge, M. (1997). Detection of DNA polymorphisms in the genus *Kalanchoe* by RAPD-PCR fingerprint and its relationships to infrageneric taxonomic position and ecophysiological photosynthetic behavior of the species. **Plant Science** 125:41-51.
- George, A. (1998). Survey of potential disease causing organisms associated with *Gracilaria changii*. M. Biotech. thesis. University of Malaya, Kuala Lumpur, Malaysia.
- Gerwick, W.H., Bernart, M.W., Moghaddam, M.F., Jiang, Z.D., Solem, M.L. and Nagle, D.G. (1990). Eicosanoids from the Rhodophyta : New metabolism in the algae. **Hydrobiologia** 204/205 : 621-628.
- Gibbs, R.A. (1990). DNA amplification by the polymerase chain reaction. **Anal. Chem.** 62:1202-1214.
- Gilbert, D.A., Lehman, N., O'Brian, S.J. and Wayne, R.K. (1990). Genetic fingerprinting reflects population differentiation in the California Channel Island fox. **Nature** 344:764-767.
- ✓ Goff, L.J. and Coleman, A.W. (1988). The use of plastid DNA restriction endonuclease patterns in delineating red algae species and populations. **J. Phycol.** 24:357-368.
- Goff, L.J. and Moon, D.A. (1993). PCR amplification of nuclear and plastid genes from algal herbarium specimens and algal spores. **J. Phycol.** 29:381-384.
- Goff, L.J., Moon, D.A. and Coleman, A.W. (1994). Molecular delineation of species and species relationships in the red algal agarophytes *Gracilariopsis* and *Gracilaria* (Gracilariales). **J. Phycol.** 30:521-537.
- Gonzalez, M., Montoya, R., Candia, A., Gomez, P. and Cisternas, M. (1996). Organellar DNA restriction fragment length polymorphism (RFLP) and nuclear random amplified polymorphic DNA (RAPD) analyses of morphotypes of *Gracilaria* (Gracilariales, Rhodophyta) from Chile. **Hydrobiologia** 326/327 :229-234.
- Grainger, J. and Madden, D. (1993). The polymerase chain reaction: turning needles into haystacks. **Biologists.** 40:197-200.
- Haglund, K., Bjorklund, M., Gunnare, S., Sandberg, A., Olander, U. and Pedersen, M. (1996). New method for toxicity assessment in Marine and brackish environments using the macroalga *Gracilaria tenuistipitata* (Gracilariales, Rhodophyta). **Hydrobiologia** 326/327:317-325.
- Hamby, R.K., Sims, L., Issel, L. and Zimmer, E.A. (1988). Direct ribosomal RNA sequencing of extraction and sequencing methods for work with higher plants. **Plant Mol. Biol. Rep.** 6 : 75-92.

- Hansen, J.E. (1984). Strain selection and physiology in the development of *Gracilaria* mariculture. **Hydrobiologia** 116/117 : 89-94.
- Hardys, H., Balick, M. and Schierwater, B. (1992). Applications of random amplified polymorphic DNA (RAPD) in molecular ecology. **Mol. Eco.** 1:55-63.
- Haring, M.A., Schuring, F., Urbanus, J., Musgrave, A. and Van den Ende, H. (1996). Random amplified polymorphic DNA markers (RAPDs) as tools for gene mapping in *Chlamydomonas eugametos* (Chlorophyta). **J. Phycol** 32:1043-1048.
- Harlin, M.M., Thorn-Miller, B. and Thusby, G.B. (1979). Ammonium uptake by *Gracilaria* spec. (Florideophyceae) and *Ulva lactuca* (Chlorophyceae) in closed system fish culture. **In** : A. Jensen and J.R. Stein (eds.). **Proceedings of the IXth International Seaweed Symposium** . Priceton Science Press. pp. 285-292.
- Hilconida, P., Calumpang, E and Menez, G. (1997). **Field guide to the common mangroves, seagrasses and algae of the Philippines**. Bookmark, Inc. Philippines.
- Ho, C.L. (1995). Molecular taxonomic studies of selected *Sargassum* species (Phaeophyta). M. Biotech thesis, University of Malaya, Kuala Lumpur, Malaysia.
- Ho, C.L., Phang, S.M. and Pang, T. (1995a) Molecular Characterization of *Sargassum polycystum* and *S. siliquosum* (Phaeophyta) by polymerase chain reaction (PCR) using random amplified polymorphic DNA (RAPD) primers. **Journal of Applied Phycology** 7:33-41.
- Ho, C.L., Phang, S.M. and Pang, T. (1995b) Application of polymerase chain reaction (PCR) using random amplified polymorphic DNA (RAPD) primers in the molecular identification of selected *Sargassum* species (Phaeophyta, Fucales). **Eur. J. Phycol** 30: 273-280.
- ↪Hong, Y.K., Coury, D.A., Polne-Fuller, M. and Gibor, A. (1992). Lithium Chloride extraction of DNA from the seaweed *Porphyra perforata* (Rhodophyta). **J. Phycol.** 28:717-720.
- Hong, Y.K., Kim, S.D., Polne-Fuller, M. and Gibor, A. (1995). DNA extraction conditions from *Porphyra perforata* using LiCl. **Journal of Applied Phycology** 7:101-107.
- Hong, Y.K., Sohn, C.H., Lee, K.W. and Kim, H.G. (1997). Nucleic acid extraction from seaweed tissues for polymerase chain reaction. **J. Mar. Biotechnology** 5:95-99.

- Hongtrakul, V., Huestis, G.M. and Knapp, S.J. (1997). Amplified fragment length polymorphisms as a tool for DNA fingerprinting sunflower germplasm : genetic diversity among oilseed inbred lines. **Theor. Appl. Genet.** 95 : 400-407.
- Houvenaghel, G.T. and Mathot, J.F. (1983). The production of marine green algae in coastal waters and their culture in ponds enriched with waste water. **In** : A. Strub, A. Chartier, P. Schleser and G. Schleser (eds.). **Energy from Biomass** Elsevier Applied Science, London. pp. 308-312.
- Hu, X.Y., Ohm, H.W. and Dweikat, I. (1997). Identification of RAPD markers linked to the gene PM1 for resistance to powdery mildew in wheat. **Theor. Appl. Genet.** 94:832-840.
- Hurtado-Ponce, A.Q. and Umezaki, I. (1988). Physical properties of gels from *Gracilaria* (Rhodophyta). **Bot. Mar.** 31:171-174.
- Intasuwan, S., Gordon, M.E., Daugherty, C.H. and Lindsay, G.C. (1993). Assessment of allozyme variation among New Zealand populations of *Gracilaria chilensis*. (Gracilariales, Rhodophyta) using starch-gel electrophoresis. **Hydrobiologia** 260/261: 159-165.
- Iqbal, M.J., Aziz, N., Saeed, N.A., Zafar, Y. and Malik, K.A. (1997). Genetic diversity evaluation of some elite cotton varieties by RAPD analysis. **Theor. Appl. Genet.** 94 : 139-144.
- Janssen, P., Coopman, R., Huys, G., Swings, J., Bleeker, M., Vos, P., Zabeau, M. and Kersters, K. (1996). Evaluation of the DNA fingerprinting method AFLP as a new tool in bacterial taxonomy. **Microbiology** 142 : 1881-1893.
- Jeffreys, A.C., Wilson, A. and Thein, S.L. (1985). Hypervariable "minisatellite" regions in human DNA. **Nature** 332:278-281.
- Jin, H.J., Kim, J.H., Sohn, C.H., Dewreede, R.E., Choi, T.J., Towers, G.H.N., Hudson, J.B. and Hong, Y.K. (1997). Inhibition of Taq DNA polymerase by seaweed extracts from British Columbia, Canada and Korea. **Journal of Applied Phycology** 9 :383-388.
- Kain, J.M. and Destombe, C. (1995). A review of the life history, reproduction and phenology of *Gracilaria*. **Journal of Applied Phycology** 7 : 267-281.
- Kalnoski, M.H., McCoy-Haman, M.F. and Hollis, G.F. (1991). A rapid method for purifying PCR products for direct sequence analysis. **Biotechnology** 11:246-249.
- Kapraun, D.F. (1993). Karyology and cytophotometric estimation of nuclear DNA content variation in *Gracilaria*, *Gracilariopsis* and *Hydropuntia* (Gracilariales, Rhodophyta). **Eur. J. Phycol.** 28:253-260.

- Kapraun, D.F. and Dutcher, J.A. (1991). Cytophotometric estimation of inter- and intraspecific nuclear DNA content variation in *Gracilaria* and *Gracilariopsis* (Gracilariales, Rhodophyta). **Bot. Mar.** 34:139-144.
- Kapraun, D.F., Duthier, J.A. and Freshwater, W.D. (1993). Quantification and characterisation of nuclear genomes in commercial red seaweeds : Gracilariales and Gelidiales. **Hydrobiologia** 260/261 :679-688.
- Kapraun, D.F., Lopez-Bautista, J. and Bird, K.T. (1996a). DNA base composition heterogeneity in some agarophytes (Gracilariales, Rhodophyta) from Mexico and the Philippines. **Journal of Applied Phycology** 8: 229-237.
- Kapraun, D.F., Lopez-Bautista, J., Trono, G. and Bird, K.T. (1996b). Quantification and characterization of nuclear genomes in commercial red seaweeds (Gracilariales) from the Philippines. **Journal of Applied Phycology** 8: 125-130.
- Keeton, W.T. and Gould, J.L. (1986). **Biological Science (4th ed.)**. W.W. Norton and Company pp. 910-915.
- King, I.P., Purdie, K.A., Rezanoor, H.N., Kaebner, R.M.D., Miller, T.E., reader, S.M. and Nicholson, P. (1993). Characterization of *Thinopyrum bessarabicum* chromosome segments in wheat using random amplified polymorphic DNAs (RAPDs) and genomic in situ hybridization. **Theor. Appl. Genet.** 86:895-900.
- Klein-Lankhorst, R.M., Vermunt, A., Weide, R., Liharska, T. and Zabel, P. (1991). Isolation of molecular markers for tomato (*L. esculentum*) using RAPD. **Theor. Appl. Genet.** 83 :108-114.
- Klekowski, E.J. (1998). Mutation rates in mangroves and other plants. **Genetica** 102/103: 325-331.
- Kleppe, K., Ohtsuka, F., Kleppe, R., Molineux, I. and Khorana, H.G. (1971). Studies on polynucleotides. **J. Mol. Biol.** 56:341-361.
- Kooistra, W.H., Stam, W.T., Olsen, J.L. and Van der Hoek, C. (1992). Biogeography of the green alga *Cladophoropsis membranacea* (Chlorophyta) based on nuclear rDNA ITS sequences. **J. Phycol.** 28:660-668.
- Lee, R.E. (1980). Basic characteristics of the Algae. **In** : Phycology, International Student Edition, Cambridge pp. 1-31, 287-350. **In** : Sinnappah, N.D. (1994). **Molecular taxonomic studies of selected *Gracilaria* species (Rhodophyta)**. M. Biotech thesis, University of Malaya, Kuala Lumpur, Malaysia.

- Lee, M.H. and Nicholson, P. (1997). Isolation of genomic DNA from plant tissues. **Nature Biotechnology** 15:805-806.
- Lee, Y.K. and Tan, H.M. (1988). Genetic transformation through protoplast fusion in Algae. **In** : T. Stadler, J. Mollion, M.C. Verdu, Y. Karamanos, H. Morvan and D. Christiaen (eds.). **Algal Biotechnology**. Elsevier Applied Science. New York. pp. 101-109.
- Lerceteau, E., Robert, T., Petiard, V. and Crouzillat, D. (1997). Evaluation of the extent of genetic variability among *Theobroma cacao* accessions using RAPD and RFLP markers. **Theor. Appl. Genet.** 95:10-19.
- Li, N. and Cattolico, R.A. (1987). Chloroplast genome characterization in the red alga *Griffithsia pacifica*. **Mol. Gen. Genet.** 209:343-351.
- Li, H., Gyllensten, U.B., Cui, X., Saiki, R.k., Ehrlich, H.A. and Arnheim, N. (1988). Amplification and analysis of DNA sequences in single human sperm and diploid cells. **Nature** 335: 414-417.
- Linz, U., Delling, U. and Rubsamen-Waigmann, H. (1990). Systematic studies on parameters influencing the performance of the polymerase chain reaction. **J. Clin. Chem. Clin. Biochem.** 28:5-13.
- Lin, J.J. and Kua, J.(1997). AFLP: A novel PCR-based assay for plant and bacterial DNA fingerprinting. **Focus** 17(2):66-70.
- Liu, X.W. and Gordon, M.E. (1987). Tissue and cell culture of New Zealand *Pterocladia* and *Porphyra* species. **Hydrobiologia** 151/152: 147-154.
- Liu, Y.S., Thomas, R.J.S. and Phillips, W.A. (1995). Single-step direct PCR amplification from solid tissues. **Nucleic Acids Res.** 23(9): 1640.
- Lluisma, A.O. and Ragan, M.A. (1998a). Cloning and characterization of a nuclear gene encoding a starch-branching enzyme from the marine red alga *Gracilaria gracilis*. **Current Genetics** 34(2):105-111.
- Lluisma, A.O. and Ragan, M.A. (1998b). Characterization of a galactose-1-phosphate uridylyltransferase gene from the marine red alga *Gracilaria gracilis*. **Current Genetics** 34(2):112-119.]
- Lluisma, A.O. and Ragan, M.A. (1999). Characterization of the UDP-glucose pyrophosphorylase gene from the marine red algae *Gracilaria gracilis*. **Journal of Applied Phycology** 10(6):581-588.

- Lopez-Bautista, J. and Kapraun, D.F. (1995). Agar analysis, nuclear genome quantification and characterisation of four agarophytes (*Gracilaria*) from the Mexican Gulf Coast. **Journal of Applied Phycology** 7:351-357.
- Lorenz, M., Weihe, A. and Borner, T. (1997). Cloning and sequencing of RAPD fragments amplified from mitochondrial DNA of male-sterile and male-fertile cytoplasm of sugar beet (*Beta vulgaris* L.). **Theor. Appl. Genet.** 94:273-278.
- Maggs, C.A. and Cheney, D.P. (1990). Competition studies of marine macroalgae in laboratory culture. **J. Phycol.** 26: 18-24.
- Manhart, J.R. and McCourt, R.M. (1992). Minireview : Molecular data and species concepts in the algae. **J. Phycol.** 28:730-737.
- Maniatis, T., Fritsch, E.F. and Sambrook, J.J. (1989). **Molecular Cloning : A laboratory manual, 2nd Ed.**. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press.
- Marks, J.C. and Cummings, M.P. (1996). DNA sequence variation in the ribosomal internal transcribed spacer region of freshwater *Cladophora* species (Chlorophyta). **J. Phycol.** 32 : 1035-1042.
- Matsunaga, T. and Takeyama, H. (1995). Genetic engineering in marine cyanobacteria. **Journal of Applied Phycology** 7 : 77-84.
- Maxam, A.M. and Gilbert, W. (1977). A new method for sequencing DNA. **Proc. Nat. Acad. Sci. USA.** 74:560-564.
- Maxam, A.M. and Gilbert, W. (1980). Sequencing end-labeled DNA with base-specific chemical cleavages. **Methods Enzymol** 65:499-560.
- Mayes, C., Saunders, G.W., Tan, I.H. and Druehl L.D. (1992). DNA extraction methods for Kelp (Laminariales) Tissue. **J. Phycol** 28 : 712-716.
- Meglecz, E., Pecsénye, K., Peregovits, L. and Varga, Z. (1997). Allozyme variation in *Parnassius mnemosyne* (L.) (Lepidoptera) populations in North-East Hungary: variation within a subspecies group. **Genetica** 101:59-66.
- Meneses, I. (1996). Assessment of populations of *Gracilaria chilensis* (Gracilariales, Rhodophyta) utilizing RAPDs. **Journal of Applied Phycology** 8:185-192.
- Meneses, I. and Abbott, I.A. (1987). *Gracilaria* and *Polycavernosa* (Rhodophyta) from Micronesia. **Micronesia** 20:187-200.

- Meneses, I., Aedo, D., Sanchez, P. and Santelices, B. (1998a). Intraclonal genetic variability in *Gracilaria chilensis* : I. A result of active growth environmental conditions. **In: Abstracts for papers presented at XVIth International Seaweed Symposium, Cebu, Philippines.** April 12-17, 1998.
- Meneses, I., Aedo, D., Sanchez, P., Hormazabal, M. and Santelices, B. (1998b). Intraclonal genetic variability in *Gracilaria chilensis* :II. Induction by culture conditions. **In: Abstracts for papers presented at XVIth International Seaweed Symposium, Cebu, Philippines.** April 12-17, 1998.
- Mercier, B., Gaucher, C., Feugeas, O. and Mazurier, C. (1990). Direct PCR from whole blood, without DNA extraction. **Nucleic Acids Res.** 18:5908.
- Milliow, P. and Salleh, A. (1994). The taxonomy of *Gracilaria* of Morib, Selangor. **In: S.M. Phang, Y.K. Lee M.A. Borowitzka and B.A. Whitton (eds.). Algal Biotechnology in the Asia-Pacific Region.** University of Malaya. pp 262-267.
- Mitchell, D.H.F. (1998). Phycocolloids [Online]. Available HTTP : [http://www.botany.uwc.ac.za/algae/studentAssignments/DaleNorman98/\[1988,October 27\]](http://www.botany.uwc.ac.za/algae/studentAssignments/DaleNorman98/[1988,October 27])
- Mizukami, Y., Kito, H., Kunimoto, M. and Kobayashi, M. (1998). Effect of DNA preparation from laver (*Porphyra yezoensis*) thalli on reproducibility of RAPD (random amplified polymorphic DNA) patterns. **Journal of Applied Phycology** 10 : 23-29.
- Mizukami, Y., Okauchi, M., Kito, H., Ishimoto, S., Ishida, T. and Fuseya, M. (1995). Culture and development of electrically fused protoplasts from red marine algae, *Porphyra yezoensis* and *P. suborbiculata*. **Aquaculture** 132:361-367.
- Morand, P., Carpentier, B., Charlier, R.H., Maze, J., Orlandini, M., Plunkett and Jacob de Waart, B.A. (1991). Bioconversion of seaweeds. **In: M.D. Guiry and G. Blunden (eds.). Seaweed Resources in Europe : Uses and Potential.** John Wiley and Sons, Chichester, England. pp. 95-148.
- Morand, P., Charlier, R.H. and Maze, J. (1990). European bioconversion projects and realizations for macroalgal biomass : Saint-Cast-Le-Guildo (France) experiment. **Hydrobiologia** 204/205 : 301-308.
- Mullis, K.B., Faloona, F., Scharf, S, Saiki, R.K., Horn, G. and Ehrlich, H. (1986). Specific enzymatic of DNA *in vitro* : The polymerase reaction. Cold Spring Harbor symp. **Quant. Biol.** 51 :263-273.
- Munoz, A.A. and Santelices, B. (1994). Quantification of the effects of sporeling coalescence on the early development of *Gracilaria chilensis* (Rhodophyta). **J. Phycol** 30:387-392.

- Murano, E. (1995). Chemical structure and quality of agars from *Gracilaria*. **Journal of Applied Phycology** 7: 245-254.
- ✓Murray, M.G. and Thompson, W.T. (1980). Rapid isolation of high molecular weight plant DNA. **Nucleic Acids Res.** 8(19): 4321-4325.
- Ng, S.M. (1998). Molecular characterisation of two *Sargassum* species (Phaeophyta) using the random amplified polymorphic DNA (RAPD) technique. Master of Philosophy thesis. University of Malaya, Kuala Lumpur, Malaysia.
- Noda, H., Amano, H., Arashima, K. and Nisizawa, K. (1990). Antitumor activity of marine algae. **Hydrobiologia** 204/205 : 577-584.
- Olsen, J.L. (1990). Nucleic acid in algae systematics. **J. Phycol.** 26:209-214.
- O'Neill, N.R., Van Berkum, P., Lin, J.J., Kuo, J., Ude, G.N., Kenworthy, W. and Saunders, J.A. (1997). Application of amplified restriction fragment length polymorphism for genetic characterization of *Colletotrichum* pathogens of Alfalfa. **Host Genetics and Resistance** 8(7):745-749.
- Otto, S.P. and Hastings, I.M. (1998). Mutation and selection within the individual. **Genetica** 102/103 : 507-524.
- Pacak, A., Fiedorow, P., Dabert, J. and Szweykowska-kulinska, Z. (1998). RAPD technique for taxonomic studies of *Pellia epiphylla*-complex (Hepaticeae, Metzgeriales). **Genetica** 104:179-187.
- Palmer, J.D. (1985). Comparative organization of chloroplast genomes. **Annu. Rev. Genet.** 19 : 325-354.
- Panaccio, M., Georgesz, M., Hollywell, C. and Lew, A. (1993). Direct PCR from solid tissues without DNA extraction. **Nucleic Acids Research** 21(19): 4656.
- Papenfuss, G.F. (1953). Outstanding problems in the morphology & taxonomy of marine algae of the tropical and south Pacific. 7th Pacif. Sci. Congress 5:27-39. **In** : Fredericq, S. and Norris, J. N. (1985). Morphological studies on some tropical species of *Gracilaria* Grev. (*Gracilariaceae*, Rhodophyta) : Taxonomic concepts based on reproductive morphology. **In** : I. A. Abbott and J. N. Norris (eds.). **Taxonomy of economic seaweeds-with references to some Pacific and Caribbean species**. California Sea Grant College, La Jolla, California. Pp. 137-155.

- Papenfuss, G.F. (1967). Notes on Algal Nomenclature-V. Various *Chlorophyceae* & *Rhodophyceae*. **Phycos** 5(1-2):95-105. **In** : Fredericq, S. and Norris, J. N. (1985). Morphological studies on some tropical species of *Gracilaria* Grev. (*Gracilariaceae*, Rhodophyta) : Taxonomic concepts based on reproductive morphology. **In** : I. A. Abbott and J. N. Norris (eds.). **Taxonomy of economic seaweeds-with references to some Pacific and Caribbean species**. California Sea Grant College, La Jolla, California. Pp. 137-155.
- Parsons, T.J., Maggs, C.A. and Douglas, S.E. (1990). Plastid DNA restriction analysis links the heteromorphic phases of an apomictic red algal life history. **J. Phycol** 26:495-500.
- Patwary, M.U., Mackay, R.M. and Van der Meer, J.P. (1993). Revealing genetic markers in *Gelidium vagum* (Rhodophyta) through the random amplified polymorphic DNA (RAPD) techniques. **J. Phycol** 29:216-222.
- Patwary, M.U. and Van der Meer, J.P. (1992). Genetics and breeding of cultivated seaweeds. **Korean J. Phycol.** 7:281-318.
- Patwary, M.U. and Van der Meer, J.P. (1994). Application of RAPD markers in an examination of heterosis in *Gelidium vagum* (Rhodophyta). **J. Phycol** 30:91-97.
- Paul, S., Wachira, F.N., Powell, W. and Waugh, R. (1997) Diversity and genetic differentiation among populations of Indian and Kenyan tea (*Camellia sinensis* (L) O. Kuntze) revealed by AFLP markers. **Theor. Appl. Genet.** 94: 255-263.
- Phang, S.M., Shaharuddin, S., Noraishah, H, and Sasekumar, A. (1996). Studies on *Gracilaria changii* (Gracilariales, Rhodophyta) from Malaysia mangroves. **Hydrobiologia** 326/327 : 347-352.
- Phang, S.M. (1994). Some species of *Gracilaria* from Peninsular Malaysia and Singapore. **In** : I. A. Abbott (ed.). **Taxonomy of economic seaweeds – With references to some Pacific species volume IV**. California Sea Grant College, La Jolla, California, pp. 125-133.
- Pillmann, A., Woolcott, G.W., Olsen, J.L., Stam, W.T. and King, R.J. (1997). Inter- and intraspecific genetic variation in *Caulerpa* (Chlorophyta) based on nuclear rDNA ITS sequences. **Eur. J. Phycol.** 32:379-386.
- Plastino, E.M. and Oliveira, E.C. (1996). Approaches to the identification of terete Brazilian Gracilariaceae (Gracilariales, Rhodophyta). **Hydrobiologia** 326/327 : 145-148.
- Polne-Fuller, M. and Gibor, A. (1987). Calluses and callus-like growth in seaweeds : Induction and culture. **Hydrobiologia** 151/152:131-138.

- Powell, W., Morganate, M., Andre, C., Hanafey, M., Vogel, J., Tingey, S. and Rafalski, A. (1996). The comparison of RFLP, RAPD, AFLP and SSR (microsatellite) markers for germplasm analysis. **Mol. Breed** 3:225-238.
- Primrose, S.B. (1987). **Modern Biotechnology**. Blackwell Scientific Publications, London. Pp. 149-159.
- Rafalski, J.A., Tingey, S.V. and Williams, J.G.K. (1991). RAPD markers – a new technology for genetic mapping and plant breeding. **Ag. Biotech. News Info** 3:645-648.
- Reeve, H.K., Westneat, D.F., Noon, W.A., Sherman, P.W. and Aquadro, C.F. (1990). DNA “Fingerprinting” reveals high levels of inbreeding in colonies of the eusocial naked mole-rat. **Proc. Nat. Acad. Sci. USA** 87:2496-2500.
- Rehm, H.J., Reed, G., Puhler, A. and Stadler, P. (1993). **Biotechnology : Genetic Fundamentals and Genetic Engineering vol. 2** (edited by A. Puhler). VCH Publishers Inc., New York.
- Renn (1984). Agar and agarose : indispensable partners in biotechnology. **Ind. Eng. Chem. Prod. Res. Dev.** 23: 17-21.
- Rice, E.L. and Bird, C.J. (1990). Relationships among geographically distant populations of *Gracilaria verrucosa* (*Gracilariales*, Rhodophyta) and related species. **Phycologia** 29:501-510.
- Roa, A.C., Maya, M.M., Duque, M.C., Tohme, J., Allem, A.C. and Bonierbale, M.W. (1997). AFLP analysis of relationships among *Cassava* and other *Manihot* species. **Theor. Appl. Genet.** 95:741-750.
- ✓Roell, M.K. and Morse, D.E. (1991) Fractionation of nuclear, chloroplast, and mitochondrial DNA from *Polysiphonia boldii* (Rhodophyta) using a rapid and simple method for the simultaneous isolation of RNA and DNA. **J. Phycol.** 27:299-305.
- Rogstad, S.H., Patton, J.C., and Schaal, B.A. (1988). M13 repeat probe detects DNA microsatellite-like sequences in gymnosperms and angiosperms. **Proc. Nat. Acad. Sci. USA** 85:9176-9178.
- ✓Ruano, G., Brash, D.E. and Kidd, K.K. (1991) PCR: the first few cycles. **Amplifications** 7:1-4.
- Ryther, J.H., DeBoer, J.A and Lapointe, B.E. (1979). Cultivation of seaweeds for hydrocolloids, waste treatment and biomass for energy conversion. **Proc. Int. Seaweed symp.** 9:1-16.

- Saiki, R.K., Scharf, S., Faloona, F., Mullis, K.B., Horn, G.T. and Ehrlich, H.A. (1985). Enzymatic amplification of β -globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. **Science** 230: 1350-1354.
- Saiki, R.K., Gelfand, D.H., Stoffel, S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, K.B. and Ehrlich, H.A. (1988a). Primer directed enzymatic amplification of DNA with thermostable DNA polymerase. **Science** 239:487-491.
- Saiki, R.K., Chang, C.A., Levenson, C.H., Warren, T.C., Boehm, C.D., Kazazian, H.H. and Ehrlich, H.A. (1988b). Diagnosis of sickle cell anemia and β -thalassemia with enzymatically amplified DNA and nonradioactive allele-specific oligonucleotide probes. **N. Engl. J. Med.** 319 :537.
- Sane, A.P., Seth, P., Ranade, S.A., Nath, P. and Sane, P.V. (1997). RAPD analysis of isolated mitochondrial DNA reveals heterogeneity in elite wild abortive (WA) cytoplasm in rice. **Theor. Appl. Genet.** 95:1098-1103.
- Sanger, F., Nicklen, S. and Coulson, A.R. (1977). DNA sequencing with chain-terminating inhibitors. **Proc. Nat. Acad. Sci. USA.** 74:5463-5467.
- Santelices, B., Aedo, D. and Varela, D. (1995). Causes and implications of intra-clonal variation in *Gracilaria chilensis* (Rhodophyta). **Journal of Applied Phycology** 7: 283-290.
- Santelices, B., Correa, J.A., Meneses, I, Aedo, D. and Varela, D. (1996). Sporeling coalescence and intraclonal variation in *Gracilaria chilensis* (Gracilariales, Rhodophyta). **J. Phycol.** 32:313-322.
- Santelices, B. and Doty, M.S. (1989). A review of *Gracilaria* farming. **Aquaculture** 78:95-133.
- Santelices, B. and Varela, D. (1993). Intraclonal variation in the red seaweed *Gracilaria chilensis*. **Mar. Biol.** 116: 543-552.
- Saunders, G.W. (1993) Gel purification of red algae genomic DNA : an inexpensive and rapid method for the isolation of polymerase chain reaction-friendly DNA. **J. Phycol.** 29:251-254.
- Schierwater, B. and Ender, A. (1993). Different thermostable DNA polymerase may amplify different RAPD products. **Nucleic Acids Res.** 21(19): 4647-4648.
- Scholfield, C.I., Gacesa, P., Price, J.H., Russell, S.J. and Bhoday, R. (1991). Restriction fragment length polymorphisms of enzymically-amplified small-subunit rRNA-coding regions from *Gracilaria* and *Gracilariopsis* (Rhodophyta)-a rapid method for assessing 'species' limits. **Journal of Applied Phycology** 3:329-334.

- Scholten, O.E., Klein-Lankhorst, R.M., Esselink, D.G., De Bock, T.S.M. and Lange, W. (1997). Identification and mapping of random amplified polymorphic DNA (RAPD) markers linked to resistance against beet necrotic yellow vein virus (BNYVV) in *Beta* accessions. **Theor. Appl. Genet.** 94:123-130.
- Schramm, W. (1991). Cultivation of unattached seaweeds. **In:** M.D. Guiry and G. Blunden (eds.). **Seaweed Resources in Europe: Uses and Potential.** John Wiley and Sons pp. 379-408.
- Schut, J.W., Qi, X. and Stam, P. (1997). Association between relationship measures based on AFLP markers, pedigree data and morphological traits in barley. **Theor. Appl. Genet.** 95:1161-1168.
- Scowcroft, W.R. and Larkin, P.J. (1988). Somaclonal variation. **In:** G. Bock and J. Marsh (eds.). **Applications of plant cell and tissue culture.** John Wiley and Sons Ltd., U.K. pp. 21-35.
- Shafeei, S, Phang, S.M. and Sasekumar, A. (1994). Agar quality of *Gracilaria changii* extracted under various acid and alkali treatments. **In:** S.M. Phang, Y.K. Lee, M.A. Borowitzka, B.A. Whitton (eds.). **Algal Biotechnology in the Asia-Pacific region.** University of Malaya. pp. 64-69.
- Shi, S.Y., Zhang, Y.X., Li, Z.E. and Liu, W.Q. (1984). The yield and properties of agar extracted from different life stages of *Gracilaria verrucosa*. **Hydrobiologia** 116/117: 551-553.
- Shin, J.A., Kiyokawa, S., Kidachi, Y and Miura, A. (1995). A simple and rapid method for isolation of total DNA from free-living conchocelis of *Porphyra* (Bangiales, Rhodophyta). **The Korean Journal of Phycology** 10(2): 117-119.
- Shen, D. and Wu, M. (1995). Chromosomal and mutagenic study of the marine macroalga, *Gracilaria tenuistipitata*. **Journal of Applied Phycology** 7:25-31.
- ✓ Sinnappah, N.D. (1994). Molecular taxonomic studies of selected *Gracilaria* species (Rhodophyta). M. Biotech thesis. University of Malaya, Kuala Lumpur, Malaysia.
- Sosa, P.A., Cabrera-Perez, M.A. and Garcia-Reina, G. (1996). Genetic variation of *Gracilaria cervicornis* (Rhodophyta) gametophytes from the Canary Islands. **Eur. J. Phycol.** 31:143-147.
- Steane, D.A., McClure, B.A., Clarke, A.E. and Kraft, G.T. (1991). Amplification of the polymorphic 5.8S rRNA gene from selected Australian gigartinean species (Rhodophyta) by polymerase chain reaction. **J. Phycol.** 27:758-762.

- Steiner, J.J., Poklemba, C.J., Fjellstrom, R.G and Elliot, L.F. (1995). A rapid one-tube genomic DNA extraction process for PCR and RAPD analyses. **Nucleic Acids Res.** 23(13): 2569-2570.
- Stevens, D.R. and Purton, S. (1997). Genetic engineering of eukaryotic algae: Progress and prospects : **J. Phycol** 33:713-722.
- Tomiuk, J., Guldbrandtsen , B. and Loeschcke, V. (1998). Population differentiation through mutation and drift- a comparison of genetic identity measures. **Genetica** 102/103 : 545-558.
- Tseng, C.K. (1984). Phycological research in the development of the Chinese seaweed industry. **Hydrobiologia** 116/117:7-8.
- Ur-Rahman, H., James, D.J., Hadonou, A.M. and Caligari, P.D.S. (1997). The use of RAPD for verifying the apomictic status of seedlings of *Malus* species. **Theor. Appl. Genet.** 95:1080-1083.
- Van der Meer, J.P. (1988). The development and genetic improvement of marine crops. **In:** T. Stadler, J. Mollion, M.C. Verdus, Y. Karamonos, H. Morvan and D. Christiaen (eds.). **Algal Biotechnology**. Elsevier Applied Science, New York.
- Van der Meer, J.P. and Zhang, X.C. (1988). Similar unstable mutations in three species of *Gracilaria* (Rhodophyta). **J. Phycol** 24:198-202.
- Van der Meer, J.P. and Todd, E.R. (1977). Genetics of *Gracilaria* sp. (Rhodophyceae, Gigartinales) IV. mitotic recombination and its relationship to mixed phases in the life history. **Can. J. Bot.** 55:2810-2817.
- Van Oppen, M.J.H., Klerk, H., De Graaf, M., Stam, W.T. and Olsen, J.L. (1996). Assessing the limits of random amplified polymorphic DNAs (RAPDs) in seaweed biogeography. **J. Phycol** 32: 433-444.
- Van Oppen, M.J.H., Olsen, J.L. and Stam, W.T. (1995). Genetic variation within and among North Atlantic and Baltic populations of the bethic alga *Phycodryis rubens* (Rhodophyta). **Eur. J. Phycol.** 30: 251-260.
- Villemur, R. (1990). Circular plasmid DNAs from the red alga *Gracilaria chilensis*. **Current Genetics** 18:251-257.
- Vos, P., Hogers, R., Bleeker, M., Reljans, M., Van de Lee, T., Hornes, M., Frijters, A., Pot, J., Peleman, J., Kuiper, M. and Zabeau, M. (1995). AFLP: a new technique for DNA fingerprinting. **Nucleic Acids Res.** 21 : 4407-4414.

- Vosberg, H.P. (1989). The PCR: an improved method for the analysis of nucleic acids. **Hum. Genet.** 83: 1-15.
- Warren, W., Wheat, T. and Knudsen, P. (1991). Rapid analysis and quantitation of PCR products by high-performance liquid chromatography. **Biotechniques** 11:250-254.
- Wattier, R., Dallas, J.F., Destombe, C., Saumitou-Laprade, P and Valero, M. (1997). Single locus microsatellites in Gracilariales (RHodophyta): high level of genetic variability within *Gracilaria gracilis* and conservation in related species. **J. Phycol.** 33:868-880.
- Welsh, J. and McClelland, M. (1990). Fingerprinting genomes using PCR with arbitrary primers. **Nucleic Acids Res.** 18:7213-7218.
- White, T.J., Arnheim, N. and Erlich, H.A. (1989). The PCR. **Trends in genetics** 6: 185-188.
- ✓ Wilkie, S.E., Isaac, P.G. and Slater, R.J. (1993). Random amplified polymorphic DNA(RAPD) markers for genetic analysis in *Allium*. **Theo. Appl. Genet.** 86:497-504.
- Williams, J.G.K., Kubelik, A.R., Livak, K.J., Rafalski, J.A. and Tingey, S. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. **Nucleic Acids Res.** 18:6531-6535.
- Winberg, G. (1991). A rapid method for preparing DNA from blood, suited for PCR screening of transgenes in mice. **PCR Methods Appl.** 1:72-74.
- ✓ Wolff, K., Schoen, E.D. and Peters-van Rijn, J. (1993). Optimizing the generation of random amplified polymorphic DNAs in Chrysanthemum. **Theor. Appl. Genet.** 86:1033-1037.
- Wu, C.Y. and Lin, G.H. (1987). Progress in the genetics and breeding of economic seaweeds in China. **Hydrobiologia** 151/152: 57-61.
- Wu, D.Y., Ugozzoli, L., Pal, B.K., Qian, J. and Wallace, B.R. (1991). The effect of temperature and oligonucleotide primer length on the specificity and efficiency of amplification by the polymerase chain reaction. **DNA Cell Bio.** 10:233-238.
- Wynne, M.J. (1989). The re-instatement of *Hydropuntia* Montagne (Gracilariaceae, Rhodophyta). **Taxon** 38: 476-479.
- Xia, B.M. (1986). On *Gracildria salicornia* (C. Agardh) Dawson. **Chin. J. Oceanol. Limnol.** 4 : 100-105.

- Xia, B.M. and Abbott, I.A. (1985). The genus *Polycavernosa* Chang et Xia (Gracilariaceae, Rhodophyta): A comparison with *Gracilaria* Grev. and a key to the species. **In**: I. A. Abbott and J. N. Norris (eds.). **Taxonomy of economic seaweeds – With references to some Pacific and Caribbean species**. California Sea Grant College, La Jolla, California, pp. 157-162.
- Xia, B.M. and Abbott, I.A. (1987). New species of *Polycavernosa* Chang et Xia (*Gracilariaceae*, Rhodophyta) from the Western Pacific. **Phycologia** 26(4):405-418.
- Yabu, H. and Yamamoto, H. (1988). Chromosome number of *Gracilaria verrucosa* (Huds.) Papenfuss in the vicinity of Hakodate, Hokkaido. **Bull. Fac. Fish. Hokkaido Univ.** Vol. 39(1): 4-5.
- Yabu, H. and Yamamoto, H. (1989). Chromosome number of *Gracilaria chorda* and *G. vermiculophylla*. **The Japanese journal of Phycology** 37(3): 236-238.
- Yamado, O., Matsumoto, T., Nakashima, M. and Hagari, S. (1990). A new method for extracting DNA or RNA for polymerase chain reaction. **J. Virol. Methods** 27:203-210.
- Yamamoto, H. (1975). The relationship between *Gracilariopsis* and *Gracilaria* from Japan. **Bull. Fac. Fish. Hokkaido Univ.** 26:217-222.
- Yamamoto, H. (1978). Systematic and anatomical study of the genus *Gracilaria* in Japan. **Mem. Fac. Fish. Hokkaido Univ.** 25(2):97-152.
- Yamamoto, H. (1984). An evaluation of some vegetative features and some interesting problems in Japanese populations of *Gracilaria*. **Hydrobiologia** 116/117:51-54.
- Yamamoto, H. (1985). *Gracilaria* from Japan : vegetative and reproductive keys and list of the species. **In**: I.A. Abbott and J.N. Norris (eds.). **Taxonomy of economic seaweeds with reference to some Pacific and Caribbean species**. California Sea Grant College Program, La Jolla, CA. pp. 77-80.
- Yamamoto, H. (1994). Review on *Gracilaria sublittoralis* Yamada et Segawa (nom. nud.), Gracilariaceae, Rhodophyta. **Japan J. Phycol.** (Sorui) 42: 421-424.
- Yamamoto, H. and Sasaki, J. (1987). Crossing experiments between populations of so-called *Gracilaria verrucosa* (Huds.) Papenfuss from two localities, Shinori and Kikonai in Hokkaido. **Bull. Fac. Fish. Hokkaido Univ.** 38(4) : 335-338.
- Yamamoto, H. and Sasaki, J. (1988). Interfertility between so-called *Gracilaria verrucosa* (Huds.) Papenfuss and *G.vermiculophylla* (Ohmi) Papenfuss in Japan. **Bull. Fac. Fish. Hokkaido Univ.** 39(1):1-3.

Yaphe, W. (1984). Properties of *Gracilaria* agars. **Hydrobiologia** 116/117 : 171-186.

✓ Yu, K and Pauls, K.P. (1992). Optimisation of the PCR program for RAPD analysis. **Nucleic Acids Research** 20(10):2606.