

REFERENCES

- Addyanis, A. (1995). The abundance of stomatopoda species in the Klang Strait-Angsa Bank water. Malaysia thesis, of BSc. Degree 1994-1995, Zoology Department Faculty of Science, University of Malaya.
- Alongi, D. M. (1990). Effect of mangrove detrital outwelling on nutrient regeneration and oxygen fluxes in coastal sediments of the Central Great Barrier Reef Lagoon. *Estuar. Coast. Mar. Sci.*, **31**: 581-598.
- Alongi, D. M. and Christofferson, P. (1992). Benthic infauna and organism sediment relations in a shallow tropical coastal area: influence of outwelled mangrove detritus and physical disturbance. *Mar. Ecol. Prog. Ser.*, **81**: 229-245.
- Ambrose, R. F. (1982). Shelter utilisation by the molluscan cephalopoda *Octopus bimaculatus*. *Mar. Ecol. Prog. Ser.*, **7**: 67-73.
- Ambrose, R. F. (1988). Population dynamic of *Octopus bimaculatus*: influence of life history patterns synchronous reproduction and recruitment. *Malacologia*, **29**: 23-39.
- Anderson, T. J. (1997). Habitat selection and shelter use by *Octopus tetricus*. *Mar. Ecol. Prog. Ser.*, **150**: 137-148.
- Anderson, T. W. (1994). Role of macroalgal structure in the distribution and abundance of a temperate reef fish. *Mar. Ecol. Prog. Ser.*, **133**: 279-290.
- Andrewartha, H. G. and Birch, L. C. (1954). The distribution and abundance of animals. Chicago University Press, Chicago.
- Annual Fisheries Statistics. (1997). Department of Fisheries, Ministry of Agriculture, Malaysia.
- Ansell, A. D., Connely, C. A. and Robb, L. (1999). Distribution, movements and diet of macrocrustaceans on a Scottish sandy beach with particular reference to predation on juvenile fishes. *Mar. Ecol. Prog. Ser.*, **176**: 115-130.
- Aoyama, T. (1973). The demersal fish stocks and fisheries of the South China Sea. FAO, Rome.(SCS/DEV/73/3).
- Arnold, P.W. and Birtles R.A. (1989). Soft-sediment marine invertebrates of Southeast Asia and Australia: A Guide to Identification. 272 pp.
- Aronson, R. B. (1986). Life history and den ecology of *Octopus briareus* (Robson) in a marine lake. *J. Exp. Mar. Bio. Eco.*, **95**: 37-56.
- Auster, P. J., Malatesta, R. J., Langton, R. W., Watling, L., Valentine, P.C., Donaldson, C. S., Langton, E. W., Shepard, A. N. and Babb, I. G. (1996). The impacts of mobile fishing gear on low topography benthic habitats in the gulf

of marine (Northwest Atlantic): implications for conservation of fish populations. Rev. Fish. Sci., **4(2)**:185-202

Australian Institute of Marine Science (AIMS), (1992). A Guide of Penaeid Shrimps found in THAI waters, 77 pp.

Bailey, K. M. and Houde, E. D. (1989). Predation on eggs and larvae of marine fishes and the recruitment problem. Advance in Marine Biology., **25**:1-83.

Banse, K. (1986). Vertical distribution and horizontal transport of planktonic larvae of echinoderms and benthic polychaetes in an open coastal sea. Bull. Mar. Sci., **39**:162-175.

Barmuta, L. A. (1987). Polemics aquatic insects and biomonitoring an appraisal. Pp 65-72., In: Majer, J. D. (Editor) The role of Invertebrates in Conservation and Biological Survey, Western Australian Department of land Management Report. Western Australian Department of Land Management, Perth, Western Australia.

Barrett, J. C., Grossman, G. D. and Rosenfeld, J. (1992). Turbidity induced changes in reactive distance of rainbow trout. Trans Am. Fish Soc., **121**:437-443.

Barry, J. P. and Dayton, P. K. (1991). Physical heterogeneity and the organisation of Marine communities. P.230-270. In: (J. Kolasa and T. A. Pickett, eds.) Ecological Heterogeneity. Springer-Verlage, New York.

Bell and Galizin, R. (1984). Influence of live coral cover on coral reef fish communities. Mar.Ecol.Prog.Ser.,**15**:265-274.

Birkeland, C. (1988). The influence of echinoderms on coral reef communities. Echinoderm Studies, **3**: 1-79.

Blaber, S. J. M. (1981). The zoogeographic affinities of estuarine fishes in South-East Africa. S. Afr. J. Sci., **77**:305-307.

Blaber, S. J. M. and Blaber, T. G. (1980). Factors affecting the distribution of juvenile estuarine and inshore fish. J.Fish Biol., **17**:143-162.

Blaber, S. J. M. and Milton, D. A. (1990). Species composition structure and zoogeography of fisheries of mangrove estuaries in the Solomon Islands. Mar. Biol., **105**:759-767.

Blaber, S. J. M., Brewer D. T. and Salini., J. P. (1989). Species composition and biomass of fisheries in different habitats of a tropical Northern Australian estuary: their occurrence in the adjoining sea and estuarine dependence. Estuarine, Coastal and Shelf Science., **29**:509-531.

Boehlert, G.W. and Morgan, J.B. (1985). Turbidity enhances feeding abilities of larval pacific herring, *Clupea harengus pallasi*. Hydrobiologia., **123**: 161-170.

- Boesch, D. F. (1974). Diversity, stability and response to human disturbance in estuarine ecosystems. Proceedings of the 1st International Congress on Ecology. The Hague, Pp.109-114.
- Boto, K. G. and Bunt, J. S. (1981). Tidal export of particulate organic matter from a Northern Australia mangrove system. *Estuarine Coastal and Shelf Science.*, **13**:247-255.
- Bourget, E. D., Duise, J. and Daigle, G. (1994). Scales of substratum heterogeneity structural complexity, and the early establishment of a marine epibenthic community. *J. Exp.Mar.Biol.Ecol.*, **181**:31-51.
- Bouyocos, G. J. (1951). A recalibration of the hydrometer method for making mechanical analysis of soils . *Agron. J.* **43**:435-438.
- Branch, G. M., Barkai, A., Hockey, P. A. R. and Hutchings, L. (1987). Biological implications: causes and effects of variability in the Benguela ecosystem. *S.Afr.J.Mar.Sci.*, **5**: pp. 425-445.
- Brandimarte, A. L. and Shimizu, G. Y. (1996). Temporal and spatial variations in littoral benthic communities of Paraibuna reservoir (Sao Paulo, Brazil). *Tropical Ecology.*, **37(2)**:215-222.
- Bray, R. J. and Curtis, J. T. (1957). An ordination of the upland forest communities of Southern Wisconsin. *Ecol. Monogr.*, **27**: 325-347. In: Numerical Ecology. Legendre P. and Legendre L.(eds). (1998)., Elsevier Science B.V. Amsterdam, Netherlands. 853 pp.
- Brazeiro, A. and Omar, D. (1999). Effects of harvesting and density dependence on the demography of sandy beach populations: the yellow clam *Mesodesma mactroides* of Uruguay. *Mar. Ecol. Prog. Ser.*, **182**:127-135.
- Breitburg, D. L. (1994). Behavioral response of fish larvae to low dissolved oxygen concentrations in a stratified water column. *Marine Biology.*, **120**:615-625.
- Breitburg, D. L., Rose, K. A. and Cowan, J. H. (1999). Linking water quality to larval survival: predation mortality of fish larvae in an oxygen-stratified water column. *Mar. Ecol. Prog. Ser.*, **178**:39-54.
- Breitburg, D.L., Steiberg, N., DuBeau, S., Cooksey, C. and Houde, E. D. (1994). Effects of low dissolved oxygen on predation on estuarine fish larvae. *Mar Ecol. Prog. Ser.*, **104**:235-246
- Broom, M. J. (1982). Structure and seasonality in a Malaysian mudflat community. *Estuarine, Coastal and Shelf Science.* **15**:135-150.
- Brown, A. C. and McLachlan, A. (1991). Ecology of sandy shores. Elsevier, Amsterdam.

- Buchanan, J. B. (1984). Chapter 3: Sediment analysis. In: Holmes, N.A. and McIntyre, A.D. (ed.), Methods for the Study of Marine Benthos. 2nd Edition. Oxford: Blackwell Scientific Publications, pp. 41-65.
- Carpenter, R. C. (1986). Partitioning herbivory and its effects on coral reef algal communities. *Ecol. Monogr.*, **56**:345-363.
- Carpenter, S. R. (ed.) (1988). Complex interactions in lake communities. Springer-Verlag, New York., 283 pp.
- Carr, M. H. (1994). Effects of macroalgal dynamics on the recruitment of a temperate reef fish. *Ecology*, **75**:1320-1333.
- Cervetto, G., Gaudy, R., Pagano, M. (1999). Influence of salinity on the distribution of *Acartia tonsa*. *J. Exp. Mar. Biol. Ecol.*, **239**: 33-45.
- Cheng, B. K. (1988a). A summary of the results of studies conducted during 1972 - 1988 on northeast monsoon in Malaysia. Malaysian Meteorological Service Technical Note No.32.
- Cheng, B. K. (1988b). A summary of the results of studies conducted during 1972 - 1988 on northeast monsoon in Malaysia. Malaysian Meteorological Service Technical Note No.31.
- Chesney, E. J. (1989). Estimating the food requirements of striped bass larvae *Morone saxatilis*: effects of light, turbidity and turbulence. *Mar. Ecol. Prog. Ser.*, **53**:191-200.
- Chong, V. C. (1980). The biology of the white prawn *Penaeus merguiensis* de Man (Crustacea: Penaeidae) in the Pulau Angsa Klang Strait Waters (Strait of Malacca). M. Sc. Thesis. Department of Zoology. University of Malaya, Kuala Lumpur. 141 pp.
- Chong, V. C. (1984). Prawn resource management in the West Coast of Peninsular Malaysia. *Worldaceana* (University Malaya) **37**:3-6.
- Chong, V. C., Sasekumar, A. and Wolanski, E. (1996). The role of mangroves in retaining Penaeid prawn larvae in Klang Strait, Malaysia. *Mangroves and Salt Marshes*, **1**(1): 11-22.
- Chong, V. C., Sasekumar, A. and Lim, K. H. (1994). Distribution and abundance of prawns in a Malaysian mangrove system. In: Sudara, S., Wilkinson, C. R. and Chou, L. M. (ed) Proceedings Third ASEAN-Australia Symposium on Living Coastal Resources. Vol:2, Research papers, Chulalongkorn University, Bangkok Thailand .May 1994.
- Chong, V. C., Sasekumar, A., Leh, M. V. C. and D'cruz, R. (1990). The fish and prawn communities of a Malaysia Coastal Mangrove system with comparison to adjacent mudflates and inshore waters. *Estuarine Coastal and Shelf Science*, **31**:703-722.

- Chong, V.C. (1993). Studies on the biology and taxonomy of prawn larvae of the genera *Penaeus*, *Metapenaeus*, *Parapenaeopsis*, *Trachypenaeus* and *Metapenaeopsis* (Crustacea: Decapoda: Penacina) PhD. Thesis. Zoology Department. University of Malaya. 490 pp.
- Chong, V.C., Wee, G.H. and Sasekumar, A. (1989). The inshore fish and prawn communities of a mangrove shore. In: Asian-Australia Living Resources in Coastal Areas. First Regional Symposium, 1989. Manila, Philippines.
- Choudhury, A., Das, A., Bhattacharvay, S. and Blhunia, A. B. (1984). A quantitative assessment of benthic macrofauna in the intertidal mudflates of Sagar Island ,Sunderbans ,India. Proc. As.Mangr.Env.Res.and Manag., 298-310.
- Chua, T. E., Ross, S.A. and Yu, H. (1997). Malacca Straits; Environmental profile; In: GEF/UNDP/IMO Regional Programme for the Prevention and Management of Marine Pollution in the East Asia Seas-Quezon City. Philippines., Pp.259
- Claridge, P. N., Potter, I. C., (1986). Seasonal changes in movements, abundance, size composition and diversity of the fish fauna of the Severn estuary. J. Mar. Biol. Assoc. U.K., **66**:229.
- Cohen, A. S. (1986). Distribution and fauna associations of benthic invertebrates at Lake Turkana, Kenya. *Hydrobiologia*, **141**:179-197.
- Coleman, J. M., Gagliano, S.M., and Smith, W. G. (1970). Sedimentation in a Malaysia high tide tropical delta. In: Chong, V.C. (1993). Studies on the biology and taxonomy of prawn larvae of the genera *Penaeus*, *Metapenaeus*, *Parapenaeopsis*, *Trachypenaeus* and *Metapenaeopsis* (Crustacea: Decapoda: Penaeina) PhD. Thesis. Zoology Department. University of Malaya. 490 pp.
- Connell, S., and Jones G. P. (1991). The influence of habitat complexity on postrecruitment processes in a temperate reef fish population. *J. Exp. Mar. Bio. Ecol.*, **151**:271-294
- Crisp, D. J. and Costlow, J. D. (1963). The tolerance of developing cirripede embryos to salinity and temperature. *Oikos* **14**:22-34.
- Currin, B. M., Reed, J. P. and Miller, J. M. (1984). Growth, production, food consumption, and mortality of juvenile spot and croaker: a comparision of tidal and nontidal nursery areas. *Estuaries* **7**:451-459.
- Cushing, D. H. (1975). "Marine Ecology and Fisheries". Cambridge University Press, London. 278 pp.
- Cushing, D. H. (1990). Plankton production and year-class strength in fish populations: an update of the match / mismatch hypothesis. *Advances in Marine Biology*, **26**:249-293.

- Cyr, H. (1998). Effects of wave disturbance and substrate slope on sediment characteristics in the littoral zone of small lakes. Can. J. Fish. Aquat. Sci., **55**(4):967-976.
- Dauwe, B., Herman, P. M. J. and Heip, C. H. R. (1998). Community structure and bioturbation potential of macrofauna at four North Sea stations with contrasting food supply. Mar. Ecol. Prog. Ser., **173**:67-83.
- Davenport, J., Gruffydd, L. D. and Beaumont, A. R. (1975). An apparatus to supply water of fluctuating salinity and its use a study of the salinity tolerances of larvae of the scallop *Pectin maximus*. L. J. Mar. Biol. Assoc., U.K. **55**:391-409.
- Day, J. H. (1981). The nature, origin and classification of estuarines. In: Estuarine ecology with particular reference to southern Africa (ed. J. H. Day). A. A. Balkema, Cape Town, South Africa.. pp.1-6.
- Day, J. H., Blaber, S. J. M. and Wallace, J. H. (1981). Estuarine fishes. Pp.197-221. In: Day, J. H.(ed.) Estuarine Ecology with Particular Reference to Southern Africa, A.A. Balkema, Cape Town.
- Day, J. W., Hall, C. A., Kemp, W. M. and Yanez-Arancibia, A. (1989). Estuarine Ecology. Wiley Interscience, New York., 558 pp.
- Day, P. R. (1965). Particle fractionation and particle-size analysis. 545-567 In: Black, C. A.. et.al. (ed) Methods of Soil Analysis, Part I. Agronomy., **9**:545-567.
- Day, R. W. (1977). Two contrasting effects of predation on species richness in coral reef habitats. Marine Biology. **44**:1-5.
- De Freitas, A. J. (1986). Selection of nursery areas by six Southeast African Penaeidae, Estuar. Coast. Shef. Sci., **23**:901-908.
- De Groot, S. J. (1984). The impact of bottom trawling on benthic fauna of the North Sea. Ocean Manage., **9**: 177-129
- Dendrinos, P., Dewan, S., and Thorpe, J. P. (1984). Improvement in the feeding efficiency of larval and juvenile Dover sole (*Solea solea*) by the use of staining to improve the visibility of Artemia used as food. Aquaculture., **38**: 137-144.
- Dichl, S. (1988). Foraging efficiency of three freshwater fishes: effects of structural complexity and light. Oikos **53**:207-214.
- Doherty, P. J. And Williams, D. M. (1988). The replenishment of coral reef fish populations. Oceanogr.Mar.Biol.Annu.Rev., **26**: pp. 487-551.
- Donald L. Lovett. (1981). A Guide to the Shrimps, Prawns, Lobsters, and Crabs of Malaysia and Singapore. 156 pp.

- Drouin, G., Himmelman, J. H. and Beland, P. (1985). Impact of tidal salinity fluctuations on echinoderm and mollusc populations. *Can. J. Zool.*, **63**:1377-1387
- Ebenman, B. and Persson, L. (ed.) (1988). Size-structured populations: ecology and evolution. Springer-Verlag, Berlin., 284 pp.
- Ebert, T. A., Russell, S. C., Dixon, J. D. and Kalvass, P. (1994). Settlement patterns of red and purple sea urchins (*Strongylocentrotus franciscanus* and *S. purpuratus*) in California, USA. *Mar. Ecol. Prog. Ser.*, **111**:41-52
- Efford, E. and Hall, K. J. (1975), Marion-lake: and analysis of a lake ecosystem pp.199-219. In: T. W. M. Cameron and L. M. Billingsley (eds) Energy-flow-It's Biological Dimensions: A summary on the IBP in Canada., 1964-1974. Royal Society of Canada, Ottawa.
- Emson, R. H. and Faller-Fritsch, R. J. (1976). An experimental investigation into the effect of crevice availability on abundance and size-structure in a population of *Littorina rudis* (Maton): Gastropoda: Prosobranchia. *J. Exp. Mar. Biol. Ecol.*, **23**:285-297.
- Errington, P. L. (1946). Predation and vertebrate populations. *Q. Rev. Biol.*, **21**:144-177
- FAO. Fishery Resources and Environment Division, FAO. (1974). FAO species identification sheets for fishery purposes. Eastern Indian Ocean Fishing Area 57 and Western Central Pacific Fishing Area 71. Vol. I, II, III and IV.
- Farrell, T. M., Bracher, D. and Roughgarden, J. (1991). Cross-shelf transport causes recruitment to intertidal populations in central California. *Limnol. Oceanogr.*, **36**:279-288
- Felley, J. D. (1987). Nekton assemblages of three tributaries to the Calcasieu estuary, Louisiana. *Estuaries*, **10**:321-329.
- Fishscher, W. and Whitehead, P. J. P. (1974). FAO species Identification sheets for Fisheries purposes. Eastern Indian ocean and western Central Pacific (Fishing Area 71). Vol. I-V.
- Fleming, M., Lin, G. and Steinberg, S. L. (1990). Influence of mangrove detritus in an estuarine ecosystem. *Bull. mar. Sci.*, **47**:663-669.
- Folk, R. L. (1974). Petrology of sedimentary rocks, Hemphills, Austin, Texas, 170 pp.
- Folkvord, A. and Hunter, J. R. (1986). Size - specific vulnerability of northern anchovy, *Engraulis mordax*, larvae to predation by fishes. *Fish Bull.*, **84**:859-869.
- Forrester, G. E. (1990). Factors influencing the juvenile demography of a coral reef fish. *Ecology*, **71**:1666-1681.

- Fowler, A. J. (1990). Spatial and temporal patterns of distribution and abundance of chaetodontid fishes at One Tree Reef, southern GBR. Mar.Ecol.Prog.Ser., **64**:39-53.
- Fraser, T. H. (1997). Abundance, seasonality, community indices, trends and relationships with physicochemical factors of trawled fish in upper Charlotte harbour, Florida. Bulletin of Marine Science, **60**(3):739-963.
- Freese, L., Auster, P. J., Heifetz, J. and Wing, P. L. (1999). Effects of trawling on seafloor habitat and associated invertebrate taxa in the Gulf of Alaska. Mar. Ecol. Prog. Ser., **182**:119-126.
- Gaines, S. and Roughgarden, J. (1985). Larval settlement rate: a leading determinant of structure in an ecological community of marine intertidal zone. Proc. Natl. Acad. Sci. USA **82**:3707-3711.
- Gardner, M. B. (1981). Effects of turbidity on feeding rates and selectivity in bluegills. Trans. Am. Fish. Soc., **110**:446-450.
- Genin, A., Dayton, P. K., Lonsdale, P. F. and Spiess, F. N. (1986). Corals on seamount peaks provide evidence of current acceleration over deep-sea topography. Nature., **322**:59-61.
- George, H.P.De Bruin; Barry C. Russell and Andre Bogusch (1994). The marine fishery resources of Sri Lanka. 400 pp.
- Gerry, A. (1997). Marine Fishes of South East Asia. 292 pp.
- Gislason, H. (1994). Ecosystem effects of fishing activities in the North Sea. Mar. Pollut. Bull., **29**:520-527.
- Golley, F. B., Odum, H. T. and Wilson R. F. (1962). The structure and metabolism of a Puerto Rico red mangrove forest in May. Ecology., **43**:9-19.
- Gomes, E. D., Deocadiz, H., Hungspreugs, A., Jothy, A., Kuan, K. J., Soegiarto, A. and Wn. R. S. S. (1990). State of the marine environment in the East Asia Seas region. UNEP Regional seas Reports and studies No. **126**, 63p.
- Gonzalez, A. E., Trathan, P. N., Yau, C. and Rodhouse, P. G. (1997). Interactions between oceanography, ecology and fishery biology of the ommastrephid squid *Martialia hyadesi* in the South Atlantic. Mar.Ecol.Prog.Ser., **152**:205-215.
- Gouldig, M., Caravalho, M.L. and Ferreira, E. G. (1988). Rio Negro, rich life in poor waters. SPB academic. The Hague The Netherlands.
- Grady, J. R. (1971). The distribution of sediment properties and shrimp catch on two shrimping grounds on the continental shelf of the Gulf of Mexico. Proc.Gulf and Caribbean Fisheries Inet.23rd Ann. Sessions, pp. 139-148.

- Gray, J. S. (1974). Animal-sediment relationships. *Oceanogr.Mar.Biol.Ann.Rev.*, **12**:223-261.
- Gray, J. S. (1989). Effects of environmental stress on species rich assemblages. *Biol. J. Linn. Soc.*, **37**: 19-32.
- Gray, S. J., Poore, B. C. G., Ugland, I. K., Wilson, S. R., Olsgard, F. and Johannessen, Q. (1997). Coastal and deep-sea benthic diversities compared. *Mar. Ecol. Prog. Ser.*, **158**:97-103
- Greenstreet, S. P. R. and Hall, S. J. (1996). Fishing and ground-fish assemblage structure in the north-western North Sea: an analysis of long-term and spatial trends. *J. Anim. Ecol.*, **65**:577-598.
- Gregory, R. S., and Northcote, T. G. (1993). Surface, planktonic, and benthic foraging by juvenile chinook salmon (*Oncorhynchus tshawytscha*) in turbid laboratory conditions. *Can. J. Fish. Aqual. Sci.*, **50**: 233-240.
- Grigg, R. W. (1994). Effects of sewage discharge, fishing pressure and habitat complexity on coral ecosystems and reef fishes in Hawaii. *Mar.Ecol.Prog.Ser.*, **103**:25-34.
- Hall, S. J. (1994). Physical disturbance and marine benthic communities life in unconsolidated sediments. *Oceanogr.Mar.Biol.A.Rev.*, **32**:179-239.
- Hartwick, E. B. and Thorarinsson G. (1978). Den associates of the giant Pacific octopus *Octopus dofleini* (Wulker). *Ophelia.*, **17**:163-166.
- Hartwick, E. B., Ambrose R. F., Robinson S. M. C. (1984). Den utilisation and the movements of tagged *Octopus dofleini*. *Mar. Behav. Physiol.*, **11**:95-110.
- Hartwick, E. B., Robinson, S. M. C., Ambrose R. F., Trotter, D., Walsh, M. (1988). Inshore-offshore comparisons of *Octopus dofleini* with special reference to abundance growth and physical conditions during winter. *Malacologia*, **29**:57-68.
- Haywood, M. D.E. and Staples, D. J. (1993). Field estimates of growth and mortality of juvenile banana prawns (*Penaeus merguiensis*). *Mar.Biol.*, **116**:407-416.
- Hecht, T., Van der lingen, C. D. (1992). Turbidity-induced changes in feeding strategies of fish in estuaries. *S. Afr. J. Zool.*, **27**:95-107.
- Holthuis, L. B. (1980). FAO species catalogue Vol.I. shrimps and prawns of the world: an annotated catalogue of species interst to fisheries. FAO Fish Symposium, (125), **3**: 261 pp.
- Houde, E.D. (1987). Fish early life dynamics and recruitment variability. American Fisheries Society Symposium., **2**:17-29.

- Hutchings, P. (1990). Review of the effects of trawling on macrobenthic epifaunal communities. *Aust. J. mar. Freshwat. Res.*, **1**: 20: 1-85.
- Hynes, H. B. N. (1970). The ecology of running water. University of Toronto Press, Toronto.
- Ioannis, K. and Anastasios, E. (1997). The continental shelf of Crete: structure of macrobenthic communities. *Mar. Ecol. Prog. Ser.*, **160**:185-196.
- Jackson, G. and Jones, G. K. (1999). Spatial and temporal variation in nearshore fish and macroinvertebrate assemblages from temperate Australian estuary over a decade. *Mar. Ecol. Prog. Ser.*, **182**:253-268.
- Jansson, B. O., McIntyre, A. D., Nixon, S. W., Pammatmat, M. M., Zeitzschel, B. and Zijlstra, J. J. (1988). Coastal-offshore interactions- an evaluation of presented evidence. In : B. O. Jansson (ed.). *Coastal-offshore Ecosystem Interactions* Springer-Verlag: 357-363.
- Jennings, S., Greenstreet, S. P. R. and John D. R. (1999). Structural change in an exploited fish community: a consequences of differential fishing effects on species with contrasting life histories. *J. Anim. Ecol.*, **68**:617-627.
- Jobling, M. (1981). Temperature tolerance and the final preferendum rapid methods for the assessment of optimum growth temperature. *Journal of Fish Biology.*, **19**:439-455.
- Johannes, R. E. (1978). Reproductive strategies of coastal marine fishes in the tropics. *Environmental Biology of Fishes.*, **3(1)**:65-84.
- Jones, G. P. (1991). Postrecruitment processes in the ecology of coral reef fish populations: a multifactorial perspective. In: Sale, P. F. (ed). *The ecology of fishes on coral reefs*. Academic Press., San Diego, p 294-328.
- Jones, J. B. (1992). Environmental impact of trawling on the sea bed: a review. *NZ. J. Mar. Freshwater Res.*, **26**:59-67
- Joseph, E. G. (1972). The status of sciaenid stocks of the middle Atlantic coast. Chesapeake CT. 302 pp.
- Joseph, E.G. (1973). Analysis of a nursery ground pp. 118-121. In: Pacheco, A. L. (ed). *Proceedings of a Workshop on Egg, Larval and Juvenile Stages of fish in Atlantic Coast Estuaries*. Tech Publ. No. 1, Mid-Atlantic Coastal Fish Ctr., Highlands, NJ.
- Kamaruzaman, Raja Malik bin Raja (1995). Safety in the Strait of Malacca. Paper presented at Asia-Pacific shipping Conference, 13-14 september 1995, Singapore. 17p.
- Karakassis, I., and A. Eleftherion. (1997). The continental shelf of crete: structure of macrobenthic communities. *Mar. Ecol. Prog. Ser.*, **160**:185-196.

- Kaufman, L. S. and Ebersole, J. P. (1984). Microtopography and the organisation of two assemblages of coral reef fishes in the West Indies. *J. Exp. Mar. Biol. Ecol.*, **78**:253-268.
- Kelley, C. A., Martens, C. S. and Chanton, J. P. (1990). Variations in sedimentary carbon remineralisation rates in the White Oak River estuary, North Carolina. *Limnol. Oceanogr.*, **26**:891-897
- Kennish, J. Michael (1990). Ecology of Estuaries., Volume II Biological Aspects. By CRC press, Inc., 291pp.
- Kerfot, W. C. and Sih, A. (1987). Predation: direct and indirect impacts on aquatic communities. University Press of New England. Hannover., 386pp.
- Kramer, D. L. (1987). Dissolved oxygen and fish behavior. *Envir. Biol. Fish.*, **18**:81-92
- Krishnamurthy, K., and Jeyaseelan, M. J. P., (1981). The early life history of fishes from Pichavaram mangrove ecosystem of India. *Reports et proces-verbaux des Reunions. Conseil Permanent International pour l'Exploration de la Mer.*, **178**:416-423.
- Kristensen, E., Holmer, M. and Bussarawit, N. (1991). Benthic metabolism and sulfate reduction in a Southeast Asia mangrove swamp. *Mar. Ecol. Prog. Ser.*, **73**: 93-103.
- KrÖneke, I., Dippner, J. W., Heyen, H. and Zeiss, B. (1988). Long-term changes in macrofaunal communities of Nordeney (East Frisia, Germany) in relation to climate variability. *Mar. Ecol. Prog. Ser.*, **167**:25-36
- Kumlu, M. and Jones, D. A. (1995). Salinity tolerance of hatchery-reared postlarvae of *Penaeus indicus* H. Milne Edwards originating from India. *Aquaculture.*, **130**:287-296
- Kurihara, T. (1999). Effects of sediment type and food abundance on the vertical distribution of the starfish *Asterina pectinifera*. *Mar. Ecol. Prog. Ser.*, **181**:269-277.
- Lack, D. (1954). The natural regulation of animal numbers. Oxford University Press. London
- Langton, R. W. and Robinson, W. E. (1990). Faunal associations on scallop grounds in the western Gulf of Maine. *J. Exp. Mar. Biol. Ecol.*, **144**:157-171
- Legendre, L. and Legendre, P. (1998). Numerical Ecology. 2nd edition, Elsevier. New York., 853 pp.
- Levin, P. S. (1991). Effects of microhabitat on recruitment variation in a Gulf of Marine reef fish. *Mar. Ecol. Prog. Ser.*, **75**:183-189.

- Levin, P. S. (1993). Habitat structure, conspecific presence and spatial variation in the recruitment of a temperate reef fish. *Oecologia*, **94**:176-185.
- Leyu, C. H. and Ling, L. K. (1988). Diurnal and seasonal rainfall variations in the Klang Valley and the adjacent sea. Malaysian Meteorological Service Technical Note No.**35**.
- Lim, Kelvin K. P and Low, Jeffrey K. Y. (1998). A Guide to common marine fishes of Singapore., Singapore Science Centre Pp. 163.
- Little, M. C., Reay, P. J. and Grove, S. J. (1988). The fish community of an east African mangrove creek. *Jounal of Fish Biology*, **32**:729-747.
- Livingston, R. J. (1990). Inshore marine habitats. Pp 549-573 In: Myers, R. L. and Ewel, J. J. (eds). *Ecosystems of Florida*. University of Central Florida Press, Orlando, Florida.
- Lobban, C. S. and Harrison, P. J. (1994). *Seaweed ecology and physiology*. Cambridge University Press, U.S.A.
- Longhurst, A. R. and Pauly, D. (1987). *Ecology of tropical oceans*. Academic Press Inc., San Diego., 407 p.
- Longley, W. L.. (1994). Freshwater inflows to Texas bays and estuaries: ecological relationships and methods for determination of needs. Texas Water Development Board and Texas Parks and Wildlife Department. Austin
- Luckhurst, B. E. and Luckhurst, K. (1978). Analysis of the influence of substrate variables on coral reef fish communities. *Mar.Biol.*, **49**:317-323.
- MacArthur, R. H. and Mac Arthur, J. W. (1961). On bird species diversity. *Ecology*, **42**:594-598.
- Macintosh, D.J. (1984). Ecology and production of Malaysia mangrove crab populations (Decapoda: Brachyura). in: Socpadmo, E.. Roa, A.N. and Macintosh, D.J. (eds), Proceeding of the UNESCO Asia Symposium on Mangrove Environment: Research and Management. pp 354-377.
- Macnae, W. (1968). A general account of the fauna and flora of mangrove swamps and forests in the Indo West Pacific region. *Adv. Mar. Biol.*, **6**:73-270.
- Macnae, W. (1974). Mangrove forest and fisheries. Food and Agriculture Organisation of the United Nation. Rome. IOFC/DEU/ 74/34.1-35.
- Magnuson, J., Crowder, L. B. and Medvick, P. A. (1979). Temperature as an ecological resource. *American Zoologist*, **19**:331-343.
- Malthus, T. R., (1966). First essay on population. Macmillan. London. 396pp.

- Mangold, K. (1983). *Octopus vulgaris*. In: Boyle P. R. (ed.) Cephalopod life cycles, Vol. I, Species accounts. Academic Press, London, pp.335-364.
- Mann, K. H. (1993). Physical oceanography, food chains and fish stocks: a review. ICES J. Mar. Sci., **50**, pp. 105-119.
- Maravelias, C. D. (1999). Habitat selection and clustering of pelagic fish: effects of topography and bathymetry on species dynamics. Canadian Journal of Fisheries and Aquatic Sciences., **56**:337-450.
- Margalef, R. (1968). Perspective in Ecological Theory. University of Chicago Press, Chicago, 111 pp.
- Mather, J. A. and O'Dor, R. K. (1991). Foraging strategies and predation risk shape the natural history of juvenile *Octopus vulgaris*. Bull. Mar. Sci., **49**:256-269.
- Matthews, W. J. and Maness, J. D. (1979). Critical thermal maxima, oxygen tolerance and success of cyprinid fishes in a southwestern river. American Midland Naturalist., **102**:374-377.
- Mazeaud, M. M., Mazeaud, M. F. and Donaldson, E. M. (1977). Primary and secondary effects of stress in fish: some new data and a general review. Trans. Am. Fish. Soc., **106**:201-222
- McClanahan, T. R. (1992). Resource utilization, competition and predation: a model and example from coral reef grazers. Ecol. Mod., **61**:195-215.
- McClanahan, T. R. and Kurtis, J. K. (1991). Population regulation of the rock-boring, sea urchin., *Echinometra mathaei* (de Blainville). J. Exp. Mar. Biol. Ecol., **147**: 121-146.
- McClanahan, T. R. and Shafir, S. H. (1990). Causes and consequences of sea urchin.. abundance and diversity in Kenyan reef lagoons. Oecologia, **83**:362-270.
- McCormick, M. I. (1994). Comparison of field methods for measuring surface topography and their associations with a tropical reef fish assemblage. Mar. Ecol. Prog. Ser., **112**:87-96.
- McCoy, E. D. and Bell, S. S. (1991). Habitat structure: the evolution and diversification of complex topic. In: Bell, S. S., McCoy, E. D. and Mushinsky, H. R. (Eds.), Habitat structure: the physical arrangement of objects in Space. Chapman and Hall, New York:3-27.
- McGehee, M. A. (1994). Correspondence between assemblages of coral reef fishes and gradients of water motion, depth, and substrate size off Puerto Rico. Mar. Ecol. Prog. Ser., **105**:243-255.
- Menge, B. A. and Sutherland, J. P. (1976). Species diversity gradients: synthesis of the roles of predation, competition and temporal heterogeneity. American Naturalist., **110**:351-369.

- Menge, B. A., Lubchenco, J. and Ashkenas, L. R. (1985). Diversity, heterogeneity and consumer pressure in a tropical rocky intertidal community. *Oecologia*, **65**:349-405.
- Messieh, S. N., Rowell, T. W., Peer, D. L. and Cranford, P. J. (1991). The effects of trawling, dredging and ocean dumping on the eastern Canadian continental shelf seabed. *Contshelf Res.*, **11**: 1237-1263
- Minello, T. J. and Zimmerman, R. J. (1983). Fish predation on juvenile brown shrimp, *Penaeus aztecus* Ives:the effect of simulated *Spartina* structure on predation rates. *J.Exp.Mar.Biol.Ecol.*, **72**: 211-231.
- Miner, J. G. and Stein, R. A. (1993). Interactive influence of turbidity and light on larval bluegill (*Lepomis macrochirus*) foraging. *Can. J. Fish Aquat. Sci.*, **50**:781-788.
- Mohsin, A.K.M. and Ambak, M.A. (1996), Marine Fishes and Fisheries of Malaysia and Neighbouring Countries. Universiti Pertanian Malaysia Press. 244 p.
- Montague, C. L. and Ley, J. A. (1993). A possible effect of salinity fluctuation on abundance of benthic vegetation and associated fauna in Northeastern Florida Bay. *Estuaries*, **16**:707-717
- Moore, H. B. (1972). Aspects of stress in the tropical marine environment. *Adv. Mar. Biol.*, **10**:217-269
- Moore, R. (1978). Variations in the diversity of summer estuarine fish populations in Arkansas Bay, Texas. 1966-1973. *Estuarine Coastal Mar. Sci.*, **6**:495-1978.
- Moranta, J., Stefanesan, C., Massuti, E., Beatriz, M., and Lioris, D. (1998). Fish community structure and depth-related trends on the continental slope of the Bllearic Islands (Algerian basin. Western Mediterranean). *Mar. Ecol. Prog. Ser.*, **171**:247-259.
- Mossa, K. M. (1988). Strait of Malacca: environment, living resources and transnational problems. In: *Coastal zone management in the Strait of Malacca. Proceedings of Symposium on Environmental Research and Coastal Zone Management in the Strait of Malacca*. School for Resource and Environmental Studies, Dalhousie University, Halifax, Nova Scotia.
- Moyle, P. B. and Cech, J. J. (1982). *Fishes: An Introduction to Ichthyology*, Prentice-Hall, Englewood Cliffs, NJ.
- Myers, R. A., Hutchings, J. A. and Barrowman, N. J. (1996). Hypothesis for the decline of cod in the North Atlantic. *Mar. Ecol. Prog. Ser.*, **138**:243-308.
- Nelson. (1984). *Fishes of the world*.. John Wiley and Sons, N.Y. 523p.
- Nelson, W.B., Sasekumar, A., and Ibrahim, Z.Z. (1994). Neap-spring tide effects on Dissolved Oxygen in the Malaysian estuaries. *Hydrobiologia*, **285**: 7-17.

- Norris, R. H. and Georges, A. (1986). Design and analysis for assessment of water quality. Pages 555-572., In: De Deckker, P. and Williams, W. D. (eds). Limnology in Australia. Commonwealth Scientific and Industrial Research Organization, Melbourne, Australia, and Junk Publs .Dordrecht, The Netherlands.
- Norton, S. F. (1991). Habitat use and community structure in an assemblage of cottid fishes. *Ecology*, **72**:2181-2192.
- Odum, W. E. (1988). Comparative ecology of tidal freshwater and salt marshes. *Ann. Rev. Ecol. Syst.*, **19**:147-176.
- Odum, W. E. and Heald, E. J. (1972). Trophic analysis of an estuarine mangrove community. *Bull. Mar. Sci.*, **22**: 671-738.
- Pauly, D. (1979). Theory and management of tropical multispecies stocks: a review, with emphasis on the Southeast Asian demersal fisheries. *ICLARM Stud. Rev.*, **1**:1-37.
- Pauly, D. and Navaluna, N. A. (1983). Monsoon-induced seasonality in the recruitment of Philippine fishes. *FAO Fishery Report.*, **3(291)**:824-833.
- Pearson, T. H. and Rosenberg, R. (1978). Macrofaunal succession in relation to organic enrichment and pollution of the marine environment. *Oceanogr. Mar. Biol. A Rev.*, **16**: 229-311.
- Per Sparre, P. and Venema, S. C. (1989). Introduction to tropical fish stock assessment. part1. Manual FAO Fisheries Technical paper **No.306.1**. Rev.1.Rome. FAO.376p.
- Pielou, E. C. (1969). An introduction to mathematical ecology. John Wiley and Sons. New York.
- Pinto, L. (1987). Environmental factors influencing the occurrence of juvenile fish in the mangroves of Pagbilao, Philippines. *Hydrobiologia.*, **150**:283-301.
- Pinto, L. (1988). Population dynamics and community structure of fish in the mangroves of Pagbilao, Philippines. *J. Fish. Biol.*, **3**:35-43.
- Poiner, I. R. and Harris, A. (1986). The effect of commercial prawn trawling on the demersal fish communities of the South Eastern Gulf of Carpentaria. In: Haines AK. Williams
- Prena, J., Schwinghamer, P., Rowell, J. W., Gordon, C. G., Gilkinson, K. D., Vass, W. P. and McKeown, D. L. (1999). Experimental otter trawling on a sandy bottom ecosystem of the Grand Banks of Newfoundland: analysis of trawl by catch and effects on epifauna. *Mar. Ecol. Prog. Ser.*, **181**:107-124.
- Qasim, S. Z. (1974). Productivity of backwaters and estuaries. In: Zeitzschel, B. (ed.) The biology of the Indian Ocean., pp 143-154. Springer-Verlag, Berlin.

- Raffaelli, D. G., Hughes, R. N. (1978). The effects of crevice size and availability on populations of *Littorina rudis* and *Littorina neritoides*. J Anim. Ecol., **47**:71-83.
- Rao, G. S., (1990). An assessment of the penaeid prawn seed resource of the Godavari estuary and the adjacent backwaters. Indian J. Fish., **37**:99-108.
- Rau, N. and Rau, A. (1980). Commercial fishes of the Philippines. 623 pp.
- Reddering, J. S.V. (1988). Prediction of the effects of reduced river discharge on estuaries of the south-eastern Cape Province, South Africa. S. Afr. J. Sci., **84**:726-730
- Reise, K. (1982). Long-term changes in the macrobenthic invertebrate fauna of the Wadden Sea: are polychaetes about to take over? Neth. J. Sea. Res., **16**:29-36
- Rice, A. L., Aldred, R. G., Darlington, E., Wild, R. A. (1982). The quantitative estimation of the deep-sea megabenthos: a new approach to an old problem. Oceanol. Acta., **5**:63-72.
- Roberts, C. M. and Ormond, R. F. G. (1987). Habitat complexity and coral reef fish diversity and abundance on Red Sea fringing reefs. Mar.Ecol.Prog.Ser., **41**:1-8.
- Robertson, A. I. (1988a). Abundance, diet and predators of juvenile banana prawns, *Penaeus merguiensis*, in a tropical mangrove estuary. Aust. J. mar. Freshwat. Res., **39**:467-478.
- Robertson, A. I. (1988b). Abundance, diet and predators of juvenile banana prawn *Penaeus merguiensis* in tropical mangrove estuary. Aust. J. Mar. Freshw. Res., **39**: 467-478
- Robertson, A. I. and Duke, N. C. (1987). Mangroves as nursery sites: comparisons of the abundance and species composition of fish and crustaceans in mangroves and other nearshore habitats in tropical Australia. Mar. Biol., **96**: 193-205.
- Robertson, A. I. and Duke, N. C. (1990). Mangrove fish communities in tropical Australia: spatial and temporal patterns in densities, biomass and community structure. Mar. Biol., **104**: 369-379.
- Robertson, A. I., Dixon, P. and Daniel, P. A. (1988). Zooplankton dynamics in mangrove and other nearshore habitats in tropical Australia. Mar.Ecol.Prog.Ser., **43**:139-150.
- Rothlisberg, P. C., Jackson, P. J. and Pendrey, R. C. (1985). Distribution and abundance of early penaeidae larvae in the Gulf of Carpentaria, Australia. In: "Second Australia National Prawn Seminar" (P. C. Rothlisberg, B. J. Hill and D. J. Staples, eds), NPS2, Cleveland, Australia, pp. 23-30.

- Roughgarden, J., Gaines, S. and Possingham, H. (1988). Recruitment dynamics in complex life cycles. *Scince*, **241**:1460-1466
- Roughgarden, J., Pennington, J. T., Stoner D., Alexander, S. (1991). Collisions of upwelling fronts with the intertidal zone: the cause of recruitment pulses in barnacle populations of central California. *Acta. Oecol.*, **12**:35-51.
- Rozas, L. P. and Odum, W. E. (1987). Use of tidal freshwater marshes by fishes and macrofaunal crustaceans along a marsh stream-order gradient. *Estuaries*, **10**:36-43.
- Russ, G. R. (1991). Coral reef fisheries: effects and yields pp.601-635. In: Sale, P. F. (ed) *The Ecology of fishes on Coral Reefs* Academic Press. San Diego.
- Sagathevan, K. (1989). Tidal mudflats – An invaluable natural resource. Proc. 12th Annual Seminar of the Malaysia Society of Marine Science.
- Saila, S. B. (1975). Some aspects of fish production and cropping in estuarine systems in estuarine Research 1:473.
- Sainsbury, K. J. (1987). Assessment and management of the demersal fishery on the continental shelf of Northwestern Australia. In: Povolina, J. J., Ralston, S. (eds) *Tropical snappers and groupers;biology and fishery management*. Westview Press. London. pp 465-503
- Sale, P. F. (ed.) (1991). *The ecology of fish on coral reefs*. Academic Press, New York.
- Sanchez, P. A. and Miller, R. H. (1986). Organic matter and soil fertility management in acid soils of the tropics. International Society of Soil Science Transactions 13th Congress., **6**: 609-625.
- Sasekumar, A., Chong, V. C. and Singh, H. R. (1994). The physical and chemical characteristics of Matang mangrove waters. In: Sudara, S., Wilkinson, C. R. and Chou, I. M. (ed). *Proceedings Third ASEAN-Australia Symposium on living Coastal Resources*. Vol:2. Research papers. Chulalongkorn University. Bangkok Thailand May 1994.
- Sasekumar, A., Chong, V. C., Charles leh, M. U. (1989). Fishery resources in mangrove and inshore waters of Selangor Living Resources in Coastal Area (Malaysia). Dept. Zoology, Univ. Malaya, Kuala Lumpur. 190pp.
- Sasekumar, A., Chong, V.C., Leh, M.U. and D'Cruz, R. (1992). Mangroves an habitat for fish and prawns. *Hydrobiologia*, **247**: 195-207.
- Sasekumar, A., Ong, T. L. and Thong, K. L. (1984). Predation of mangrove fauna by marine fishes. In: Soepadmo, E., Rao, A. N. and Macintosh, D. J. (eds). *Proc. Asian Symp. on Mangrove Environment: Research and Management*. University of Malaya. pp.378-384.

- Schlacher, T. A., Newell, P., Clavier, J., Schlacher-Hoenlinger, M. A., and Chevillon, J. B. (1998). Soft-sediment benthic community structure in a coral reef lagoon- the prominence of spatial heterogeneity and spot endemism. Mar. Ecol. Prog. Ser., **174**:159-174.
- Schneider, D. C., Gagnon, J. M. and Gilkinson, K. D. (1987). Patchiness of epibenthic megafauna on the outer Grand Banks of Newfoundland. Mar. Ecol. Prog. Ser., **39**:1-13
- Scott, J. S. (1959). An Introduction to the Sea fishes of Malaya. Pp. 180
- Selvarajah, V. (1961). A study of drift in the north Malacca Straits from salinity determinations. Proceedings of the Indo-Pacific Fisheries Council. **9(2)**:1-6.
- Shanks, A. L. (1995). Mechanisms of cross-shelf dispersal of larvae invertebrates and fish. In: McEdward L. (ed). Ecology of marine invertebrates. CRC Publications, Boca Raton, pp.323-368.
- Shannon, C. E. and Weaver, W. (1963). The mathematical theory of communication. University of Illinois press, Urbana.
- Sheridan, P. F. and Livingston, R. J. (1979). Cyclic tropic relationships of fishes in an unpolluted, river-dominated estuary in north Florida, in Ecological Processes in Coastal and Marine System. Plenum Press, New York. pp.143.
- Sinclair, M., Tremblay, M. J., Bernal, P. (1985). El Nino events and variability in Pacific mackerel (*Scomber japonicus*) survival index: support for Hjort's hypothesis. Canadian Journal of Fisheries and Aquatic Sciences.. **42**:602-608.
- Singh, H. R. and Sasekumar, A. (1994). Distribution and abundance of marine catfish (Family: Ariidae) in the Matang mangrove waters. In: Sudara, S., Wilkinson, C. R., and Chou, L. M. (eds), (1994). Proceedings, Third ASEAN-Australia Symposium on living Coastal Resources. 2: Research Papers. Chulalongkorn university, Bangkok, Thailand.
- Smith, J.L.B. (1950). The sea fishes of Southern Africa. 550 pp.
- Soegiarto, A. (1985). Oceanographic assessment of the East Asia Seas.p 173-184. In: Environment and resources in the Pacific. UNEP Regional Seas Reports and Studies No.69. United Nations Environment Programme.
- Soeriaatmadja, R. E. (1956). Surface salinities in the Strait of Malacca. Marine Research in Indonesia., **2**: 27-48.
- Sogard, S. (1992). Variability in growth rates of Juvenile fishes in different estuarine habitats. Mar.Ecol.Prog.Ser., **85**:35-53.
- Sokal, R. and Rohlf, R. J. (1974). Biometry. W.H.Freeman, San Francisco.775pp.

- Springer, S. and Bullis, H. R. (1954). Exploratory shrimp fishing in the Gulf of Mexico. Summary report for 1952-54 Comm.Fish. Rev., **16**:1-16.
- Springer, V. G., (1982). Pacific Plate biogeography with special reference to shore fishes. Smithsonian contributions, Zoology., **367**:1-182.
- Staples, D. J., Vance, J. D., and Heales, D. S. (1985). Habitat requirement of juvenile penaeid prawns, pp.47-54. In Rothlisberg, P. C., B. J. Hill and D. S. Staples (eds.). Second National Prawn Seminar. NSP2, Cleveland, Australia.
- Stein, R. A., Threlkeld, S. T., Sandgren, C. D., Sprules, W. G., Persson, L., Werner, E. E., Neill, W. H. and Dodson, S. I. (1988). Size structured interactions in lake communities. Pp. 161-179. In: Ebenman, B. and Persson, L. (ed.) Size-structured populations:Ecology and Evolution. Springer-Verlag. Berlin.
- Stoner, A. W. and Zimmerman, R. J. (1988). Food pathways associated with penaeid shrimps in a mangrove-fringed estuary. Fishery Bulletin, U.S., **86**:543-551.
- Tan Leo W. H and Ng Peter K. L. (1988). A guide to seashore life. Singapore Science Centre 160 pp.
- Tan, S. R. Munro (1955). The marine and fresh water fishes of Ceylon. 349 pp.
- Tejerina-Garro, F.L., Fortin, R. and Rodriguez, M. A. (1998). Fish community structure in relation to environmental variation in flood plain lakes of the Araguaia River, Amazon Basin. Environmental Biology of Fishes., **51**: 399-410.
- Ter Braak, J.F. and Petr S. (1998). Canoco Reference Manual and User's Guide to Canoco for Windows: Software for Canonical Community Ordination (version 4). Microcomputer Power (Ithaca, NY, USA), 352 pp.
- Terek, J. (1980). Zonation of the macrozoobenthos and ecological conditions in the Velky Viholart lake. Biologia., **35**:111-120
- Thresher, R. E. (1983). Environmental correlates of the distribution of planktivorous fishes in the One Tree Reef Lagoon. Mar.Ecol.Prog.Ser., **10**:137-145.
- Thurow, F. (1997). Estimation of the total fish biomass in the Baltic Sea during the 20th Century. Journal of Marine Science., **54**:444-461.
- Trewavas (1977). The sciaenid fishes "Croakers or drums" of the Indo-West-Pacific.. Published by Academic press for Transactions of Zoological Society of London.. **33**:pp 253-541
- Tupper, M. and Boutilier, R.G. (1995). Effects of habitat on settlement, growth, and postsettlement survival of Atlantic cod (*Gadus morhua*). Can. J. Fish. Aquat. Sci., **52**:1834-1841.

- Tyson, R. V. and Pearson, T. H. (1991). Modern and ancient continental shelf anoxia. Geological Society Special Publication., **58**. London, UK.
- Uktolseyea, H. (1988). Physical and biological characteristics of the Strait of Malacca in the framework of coastal management, P. 118-131. In: Coastal zone management in the Strait of Malacca. Proceedings of Symposium on Environmental Research and Coastal Zone Management in the Strait of Malacca. School for Resource and Environmental Studies, Dalhousie University, Halifax, Nova Scotia.
- Uktolseyea, H. (1990). Physical conditions of the Indonesia waters, p. 34-45. In: H. T. Yap, M. Bohle-carbonell and E. D. Gomez (eds.) Oceanography and marine pollution: an ASEAN-EC perspective. Marine Science Institute Hamburg.
- Vance, D. J., Haywood, M. D. E. and Staples, D. J. (1990). Use of a mangrove estuary as a nursery area for post larval and juvenile banana prawns, *Penaeus merguiensis* de Man, in northern Australia, Estuarine, Coastal and Shelf Science., **31**:689-701.
- Vance, J. D., Haywood, M. D. E., Heales, D. S., Kenyon, A. R., and Loneragan, R. R. (1998). Seasonal and annual variation in abundance of postlarval and juvenile banana prawns *Penaeus merguiensis* and environmental variation in two estuaries in tropical northeastern Australia: a six year study. Mar. Ecol. Prog. Ser., **163**:21-36
- Villoso, E. P. and Hermosa, G. V. (1982). Demersal trawl fish resources of Samar Sea and Carigara Bay. Philippines. Fishery Res J Univ Phillip. Quezon City., **7**: 59-78
- Vinyard, G. L. and O'Brien, W. J. (1976). Effects of light and turbidity on the reactive distance of bluegill (*Lepomis macrochirus*). J. Fish. Res. Bd. Can., **33**:2845-2849.
- Voight, J. R. (1988). A technique for trapping sandflat octopuses. Am. Malacol. Bull., **6(1)**:45-48.
- Voight, J. R. (1992). Movement injury and growth of members of a natural population of Pacific pygmy octopus *Octopus digueti*. J. Zool. Lond., **228**:247-263.
- Warwick, R. M. and Uncles, R. J. (1980). Distribution of benthic macrofauna associations in the Bristol Channel in relation to tidal stress. Marine Ecol. Prog. Ser., **3**:97-103.
- Weinstein, M. P., Weiss, S. L., Hodson, R. G. and Gerry, L. R. (1980). Retention of three taxon of postlarval fishes in an intensively flushed tidal estuary, Cape Fear River, North Carolina. Fishery Bulletin, U.S. **78**:419-436.
- Welcomme, R. L. (1985). River fisheries. FAO Fish. Tech. Pap., **262**, Rome.330 pp.

- Wells, F. E. (1990). Distribution of marine invertebrates in a Hong Kong mangrove with emphasis on Molluscs. Proceedings of the second International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China, Hong Kong. (1986). (Ed.B.Morton) Hong Kong: Hong Kong Press:783-793.
- Wells, M. J. (1978). Octopus: Physiology and behavior of an advanced invertebrate. Chapman and hall. London.
- Wetzel, R. G. (1983). Limnology. Saunders. Philadelphia. Pennsylvania, USA.
- Whitfield, A. K. and Bruton, M. N. (1989). Some biological implications of reduced freshwater inflow into eastern Cape estuaries: a preliminary assessment. S. Afr. J. Sci., **85**:691-694.
- Whittaker, R. H. (1969). Evolution of diversity in plant communities. Brookhaven Symposium on Biology., **22**:178-196.
- Wilbur, H. M. (1988). Interaction between growing predators and growing prey. Pp. 157-172. In: Ebenman, B. and Persson, L. (ed.) Size-structured Populations: Ecology and Evolution. Springer-Verlag.Berlin.
- Williams, H. B. (1958). Substrates as a factor in shrimp distribution. Limnol.Oceanogr., **3**:283-290.
- Wing, S. R., Botsford, L. W. and Largier, J. L. (1995a). Spatial structure of relaxation events and crab settlement in the northern California upwelling system. Mar. Ecol. Prog. Ser., **128**:199-211
- Wing, S. R., Largier, J. L., Botsford, L. W. and Quinn, J. F. (1995b). Settlement and transport of benthic invertebrates in an intermittent upwelling zone. Limnol. Oceanogr., **40**:316-329
- Yanez-Arancibia, A., (ed.). (1985). Fish community ecology in estuarines and coastal lagoons. Dirección General de Publicaciones. Universidad Nacional Autónoma de México, UNAM Press Mexico City, 653pp.
- Yap, Y. N. (1995). The biology of sciaenidae in the Matang Mangrove waters. Malaysia. Thesis of master of science. Department of zoology University of Malaya, Kuala Lumpur. Malaysia.
- Yarnell, J. L. (1969). Aspect of the behavior of *Octopus cyanea* (Gray). Anim. Behav., **17**:747-754.
- Yoon, W. B. and Benner, R. (1992). Denitrification and oxygen consumption in sediments of two south Texas estuaries. Mar. Ecol. Prog. Ser., **90**:157-167
- Zar Jerrold H. (1984). Biostatistical analysis. Second edition by Prentice Hall, Inc. A Simon and Schuster company England Cliffs, New Jersey pp 718.

Zieman, J. C., Macko, S. A. and Mills, S. L. (1984). Role of sea grass and mangrove in estuarine food webs: temporal and spatial changes in stable isotope composition and amino acid content during decomposition. Bull. mar. Sci., **35**: 380-392.

Zimmerman, A. R. and Benner, R. (1994). Denitrification, nutrient regeneration and carbon mineralisation in sediments of Galveston Bay, Texas, USA. Mar. Ecol. Prog. Ser., **114**:275-288