

# A PROFILE OF DENGUE PATIENTS AT UNIVERSITY MALAYA MEDICAL CENTRE

ť

BY

# CHRIS NG FOOK SHENG

(EGE030024)

# SUBMITTED TO THE FACULTY OF ECONOMICS AND ADMINISTRATION UNIVERSITY OF MALAYA IN PARTIAL FULFILLMENT OF THE REQUIREMENT FOR THE DEGREE OF MASTER OF APPLIED STATISTICS

MAY 2005



#### ACKNOWLEDGEMENT

My utmost gratitude to Associate Professor Dr Noor Azina Ismail for her penetrating insights, invaluable guidance and assistance throughout this paper.

Special thanks to Professor Dr Lucy Lum for her generosity in allowing me the access to the data for this research paper.

I am endlessly indebted to Dr David Yeoh for his selfless support and unceasing motivation much needed to complete this paper. You inspired me!

Finally, my heartfelt appreciation to my family for their kind understanding and patience without which all of these would have been impossible.

#### ABSTRACT

The paper aims to study the prevalence of various dengue symptoms in children and adults diagnosed as having dengue fever (DF) and dengue haemorrhagic fever (DHF). Effort to predict and classify patients serologically is also attempted. Adults and children suspected of dengue infection are classified clinically into DF and DHF via a model developed in the paper. Part of the paper also looks at the various aspect of patient management in relation to the dengue outbreak experienced at the University Malaya Medical Centre (UMMC).

Findings reveal the low notification rate of the clinical dengue cases at UMMC which are made notifiable by law as part of the disease surveillance effort to ensure prompt action during the course of an outbreak.

In this dengue episode, close to one third of the suspected dengue cases at UMMC were not tested serologically despite the high percentage of positive outcome for those tested.

The reclassification of the clinical dengue cases by the World Health Organisation (WHO) guidelines implies that about 11.4% of the clinically cases were potentially "misdiagnosed".

Symptoms significant in the differential diagnosis between serologically confirmed dengue cases and other viral infections are skin rash, platelet count at admission and abdominal pain. Patients citing such symptoms have greater odds of being tested positive of such infection.

The clinical diagnosis of DHF among the pediatric patients appears to conform to the WHO definition. However, results show the otherwise for the adult patients.

### CONTENTS

			Page	
Acknowledge	ement		i	
Abstract			ii	
List of Table	ist of Tables			
List of Figur	es		viii	
Chapter 1.	Intro	duction	1	
	1.1 1.2 1.3 1.4 1.5 1.6 1.7	Overview History of Dengue Dengue Infection Global Impact of Dengue Situation of Dengue in Malaysia Objectives of the Study Significance of the Study Organization of the Paper	1 4 5 6 12 12	
Chapter 2.	Liter	ature Review	15	
	2.1 2.2 2.3 2.4	Scope of Review Profile and Management of Dengue Patients Clinical and Laboratory Findings on Dengue Infection Summary	15 15 17 25	
Chapter 3.	Data	and Methodology	26	
	3.1 3.2 3.3 3.4 3.5 3.6 3.7	Overview Data Missing Data New Variables General Statistical Method Multiple Logistic Regression Other Consideration	26 28 29 30 31 35	

Chapter 4.	The P	Profile and Management of Dengue Patients	36
	4.1	Overview	36
	4.2	Demographics of Patients	37
	4.3	Clinical Diagnosis of Dengue	44
	4.4	Length of Stay	47
	4.5	Admission of Dengue Patients	49
	4.6	Dengue Serology Test	52
	4.7	Notification of Clinical Dengue Cases	54
	4.8	Clinical and Laboratory Observation of Child and Adult	
		Dengue Patients	55
	4.9	Clinical Diagnosis of DF/DHF and the WHO Guidelines	58
	4.10	Summary	66
Chapter 5.	The P	Prediction of Dengue Serology Test Results	67
	5.1	Introduction	67
	5.2	Model Estimation - Logistic Regression	68
	5.3	Model Interpretation	71
	5.4	Model Fit Assessment	76
	5.5	Model Diagnostic	81
	5.6	Concluding Remarks	82
Chapter 6.	The C	Classification of Clinical Dengue Cases	83
	6.1	Introduction	83
	6.2	Classification of the Clinical Diagnosis of Dengue Infection	83
	0.2	6.2.1 Classification of Dengue Infection in Child Patients	84
		6.2.2 Classification of Dengue Infection in Adult Patients	93
	6.4	Concluding Remarks	100
Chapter 7.	Conc	lusion	102
	7.1	Overview	102
	7.1	Summary of Findings	102
	7.3	Discussion and Limitations of the Study	107
	7.4	Future Study	110
	7.4	and one	
Reference			111
Annendix			114

# LIST OF TABLES

Table		Page
1.1	Reports of dengue and dengue haemorrhagic fever for the world, $1956-1995$	5
1.2	Number of reported dengue cases and death in Malaysia, 1997-2002	10
2.1	World Health Organization case definition for dengue fever, dengue haemorrhagic fever and dengue shock syndrome	20
2.2	Clinical and laboratory manifestations of adult DF/DHF cases in the 1973-74 and 1982 outbreaks in Malaysia	22
2.3	Comparison of DHF manifestations in 34 children admitted to University Hospital, Kuala Lumpur in the 1982 outbreak with the findings in Thai children diagnosed as DHF	23
2.4	Clinical features compared in 61 DHF cases (1962-64) and 102 DF cases (1967-68) in Penang, Malaysia	24
3.1	Listing of variables and description	27
3.2	Threshold value specification for the diagnostic test	35
4.1	Percentage distribution of cases by ethnicity and gender	38
4.2	Descriptive statistics of age for all cases (Valid $N = 730$ )	39
4.3	Percentage distribution of child and adult cases by gender	42
4.4	Percentage distribution of child and adult cases by ethnicity	43
4.5	Percentage distribution of clinical diagnosis by gender (N = 714)	46
4.6	Percentage distribution of clinical diagnosis by ethnicity (N = 715)	46
4.7	Length of stay for all admitted patients (N = $406$ ), adult and child cases	47
4.8	Length of stay by clinical diagnosis (N = 399)	48
4.9	Percentage distribution of child and adult by admission	49
4.10	Percentage distribution of clinical diagnosis by admission	49

Table		Page
4.11	Percentage distribution of admission and thrombocytopenia of 50,000 per $\rm mm^3$ or less (column percentage)	50
4.12	Percentage distribution of admission and thrombocytopenia of 50,000 per mm <sup>3</sup> or less (row percentage)	50
4.13	Percentage distribution of admission and hematocrit change of 50% or more (column percentage)	51
4.14	Percentage distribution of admission and hematocrit change of 50% or more (row percentage) $$	51
4.15	Frequency and percentage distribution for the notification of DF/DHF to the District Health Office	54
4.16	Percentage distribution of DF/DHF notification and final dengue serology test results (row percentage)	54
4.17	Percentage of clinical and laboratory findings of child and adult clinically diagnosed as having dengue infection	57
4.18	Mann-Whitney $U$ test for symptoms grouped by child or adult	58
4.19	Cross-tabulation of clinical diagnosis and diagnosis as per the WHO guideline	59
4.20	Percentage of clinical and laboratory findings for the DF and DHF cases diagnosed as per the WHO guideline grouped by child and adult	61
4.21	Percentage of clinical and laboratory findings for the DF and DHF cases clinically diagnosed by the medical doctors grouped by child and adult	63
4.22	Percentage of clinical and laboratory findings for children and adults grouped by the diagnosis of DF and DHF as per the WHO guideline	65
5.1	Likelihood ratio test and odds ratio for the dengue serology test result of the adult patients	69
5.2	Estimated logistic regression model for the dengue serology test result of adult patients	71
5.3	Adjusted odds ratios and 95% confidence intervals for the estimated logistic model in Table 5.2	72

Table		Page
5.4	Estimated probability of positive dengue serology test under different conditions based on equation (2)	74
5.5	Classification performance of the estimated logistic model in Table 5.2	76
5.6	Measures of classification accuracy for the estimated logistic model in Table $5.2$	79
5.7	Summary of diagnostic tests for influential observations	81
6.1	Likelihood ratio test and odd ratio for the clinical dengue classification of child patients	85
6.2	Estimated logistic regression model for the clinical dengue classification in child patients	86
6.3	Adjusted odds ratio and 95% confidence intervals for the estimated logistic model in Table 6.2 $$	88
6.4	Classification performance of the estimated logistic model in Table 6.2	89
6.5	Measures of classification accuracy for the estimated logistic model in Table $6.2$	89
6.6	Likelihood ratio test and odds ratio for clinical dengue classification of adult patients	94
6.7	Estimated logistic regression model for the clinical dengue classification in adult patients	95
6.8	Adjusted odds ratio and 95% confidence interval for the estimated logistic model in Table $6.7$	96
6.9	Classification performance of the estimated logistic model in Table 6.7	97
6.10	Measures of classification accuracy for the estimated logistic model in Table 6.7	97

# LIST OF FIGURES

Figure		Page
1.1	The general distribution of dengue fever and/or dengue haemorrhagic fever, $1975-1996$	3
1.2	The spectrum of dengue virus infection	5
1.3	Incidence Rate (I.R. per 100,000 population) of reported dengue in Malaysia, 1980-2002	8
1.4	Case Fatality Rate according to DHF and DF $\&$ DHF in Malaysia, 1984-2002	11
4.1	Percentage distribution of cases by gender	37
4.2	Percentage distribution of cases by ethnicity	37
4.3	Frequency histogram of age group	39
4.4	Boxplot of age by gender of patients	40
4.5	Boxplot of age by ethnicity of patients	41
4.6	Percentage distribution of cases by child and adult patients	42
4.7	Percentage distribution of dengue patients by clinical diagnosis	44
4.8	Percentage distribution of child and adult dengue patients by clinical diagnosis	45
4.9	The percentage of clinical cases tested serologically and the result breakdown	52
4.10	Percentage of positive dengue serology result by clinical diagnosis	53
5.1	Estimated probability of positive dengue serology test for adult patients under different condition of platelet count at admission, skin rash and abdominal pain	75
5.2	Plot of sensitivity and specificity for all possible cut-points using the estimated logistic model in Table $5.2$	77
5.3	Receiver Operating Characteristic (ROC) curve for the estimated logistic model in Table 5.2	78

rigure		1 age
6.1	Plot of sensitivity and specificity for all possible cut-points using the estimated logistic model in Table $6.2$	90
6.2	Receiver Operating Characteristic (ROC) Curve for the estimated logistic model in Table 6.2	91
6.3	Estimated probability for the hematocrit change in adult patients for the classification of clinical dengue cases	96
6.4	Plot of sensitivity and specificity for all possible cut-points using the estimated logistic model in Table 6.7 $$	98
6.5	Receiver Operating Characteristic (ROC) Curve for the estimated logistic model in Table 6.7	99