

# REFERENCES

---

---

## REFERENCES

---

---

- Adams, A. and Lindahl, T. (1975) Epstein-Barr virus genome with properties of circular DNA molecules in carrier cells. *Proc. Natl. Acad. Sci. USA* **72**; 1477-1481
- Aldinger, H.K., Delius, H., Freffe, U.K., Clarke, J. and Bornkamm, G.W. (1985) A putative transforming gene of the Jijoye virus differs from that of Epstein-Barr virus prototypes. *Virology*; **141**: 221-234
- Amit, A.G., Mariuzza, R.A., Philips, S.E.V. and Poljaak, R.J. (1986) Three dimensional structure of an antigen-antibody complex at 2.8 Å resolution. *Science*; **233**: 747-753
- Alfieri, C.M., Birkenbach, M. and Kieff, E. (1991) Early events in Epstein-Barr virus infection of human B lymphocytes. *Virology*; **181**: 595-608
- Allday, M.J., Crawford, D.H. and Griffin, B.E. (1989) Epstein-Barr virus latent gene expression during the initiation of B cell immortalisation. *J Gen. Virol.*; **70**: 1755-1764
- Armstrong, R.W., Kannan-Kutty, M., Dharmalingan, S.K. and Ponnudurai, J.R. (1979) Incidence of nasopharyngeal carcinoma in Malaysia, 1968-1977. *Br. J. Cancer*; **40**: 557-567
- Armstrong, R.W., Armstrong, M.J., Yu, M.C. and Henderson, B.E. (1983) Salted fish and inhalants as risk factors for nasopharyngeal carcinoma in Malaysian Chinese. *Cancer Res.*; **43**: 2967-2970
- Arnon, R., Maron, E., Sela, M. and Anfinsen, C.B. (1971) Antibodies reactive with native lysozyme elicited by a completely synthetic antigens. *Proc. Natl. Acad. Sci. USA*; **68**: 1450-1455
- Arrand, J.R., Young, L.S. and Tugwood, J.D. (1989) Two families of sequences in the small RNA-encoding region of Epstein-Barr virus (EBV) correlate with EBV types A and B. *J. Virol.*; **63**: 983-986
- Atassi, M.Z. (1975) Antigenic structure of myoglobins: the complete immunochemical anatomy of a protein and conclusions relating to antigenic structures of proteins. *Immunochemistry*; **12**: 423-438

- Atassi, M.Z. (1980) Precise determination of protein antigenic structure has unravelled the molecular immune recognition of proteins and provides a prototype for synthetic mimicking of other protein binding site. *Mol. Cellular Biochem.*; **32**: 21
- Atassi, M.Z. (1984) Antigenic structures of proteins. Their determination has revealed important aspects of immune recognition and generated strategies for synthetic mimicking of protein binding sites. *Eur. J. Biochem.*; **145**: 1-20
- Atassi, M.Z. and Lee, C.G. (1978) The precise and entire antigenic structure of native lysozyme. *Biochem. J.*; **171**: 429
- Audibert, R., Jolivet, M., Chedid, L., Alouf, J.E., Boquet, P., Rivalle, P. and Sifert, D. (1981) Active antitoxic immunization by a diptheria toxin synthetic oligopeptide. *Nature*; **289**: 593-594
- Baer, R., Bankier, A.T., Biggin, M.D., Desinger, P.L., Farrel, P.J., Gibson, T.J., Halfull, G., Hudson, G., Satchwell, C., Sequin, C., Fuffnell, P. and Barrell, B. (1984) DNA sequence and expression of the B95-8 Epstein-Barr virus genome. *Nature (London)*; **310**: 207-211
- Bailar, J.C. (1967) Nasopharyngeal cancer in white populations - a worldwide survey In: **Cancer of Nasopharynx**, (Muir & Shnamugaratnam eds.), UICC Monograph Series, Vol I, Munksgaard, Copenhagen. pp 443-46
- Bayliss, G.J. and Nonoyama, M. (1978) Mechanisms of infection with Epstein-Barr virus, 3. The synthesis of proteins in superinfected Raji cells. *Virology*; **87**: 204-207
- Benjamin, D.C., Berzofsky, J.A., East, I.E., gurd, F.R.N., Hannum, C., Leach, S.J., Margoliash, E., Micheal, J.G., Miller, A., Prager, E.M., Reichlin, M., Sercarz, E.E., Smith-Gill, S.J., Todd, P.E. and Wilson, A.C. (1984) The antigenic structure of proteins: a reappraisal. *Ann. Rev. Immunol.*; **2**: 67-101
- Berzofsky, J.A. (1985) Intrinsic and extrinsic factors in protein antigenic structure. *Science*; **229**: 932-40
- Biggin, M., Bodescot, M., Perricaudet, M. and Farrel, P. (1987) Epsrein-Barr virus gene expression in P3HR1-superinfected Raji cells. *J. Virol.*; **61**: 3120-3132

- Boey, M.L., Wee, G.B., Mohan, C., Howe, H.S., Chan, S.H. and Feng, P.H. (1992) HLA in Singapore Chinese with rheumatoid arthritis. *J. Rheumatol*; **19**: 1517-1519
- Brooks, L., Yao, Q.Y., Rickinson, A.B. and Young, L.S. (1992) Epstein-Barr virus latent gene transcription in nasopharyngeal carcinoma cells: co-expression of EBNA1, LMP1 and LMP2 transcripts. *J. Virol.*; **66**: 2689-2697
- Buell, P. (1965) Nasopharynx cancer in Chinese of California. *Br. J. Cancer*; **19**: 459-470
- Burt, R.D., Vaughan, T.L., Nisperos, B., Swanson, M. and Berwick, M. (1994) A protective association between the HLA-A2 antigen and nasopharyngeal carcinoma in US caucasians. *Int. J. Cancer*; **56**: 465-4676
- Cameron, K.R., Staminger, T., Craxton, M., Bodmer, W., Honess, R.W. and Fleckstein, B. (1987) The 160,000 M<sub>r</sub> virion protein encoded at right end of the herpesvirus saimiri genome is homologous to the 140,000 M<sub>r</sub> membrane antigen encoded at the left end of the Epstein-Barr genome. *J. Virol.*; **61**: 2063-2070
- Cammoun, M., Hoener, G.V. and Mourali, N.L. (1974) Tumours of the nasopharynx in Tunisia: an anatomic and clinical study based on 143 cases. *Cancer*; **33**: 184-192
- Carter, J.M. (1994) Epitope prediction methods. In: **Methods in Molecular Biology, Vol. 36. Peptide Analysis Protocols** (Dunn, B.M. and Pennington, M.W. eds.), Humana Press, Totowa, New Jersey. pp 193-205
- Chan, S.H., Day, N.E., Kunaratnam, N., Chia, K.B. and Simons, M.J. (1983) HLA and nasopharyngeal carcinoma in Chinese - a further study. *Int. J. Cancer*; **32**: 171-176
- Chan, S.H., Chew, C.T., Prasad, U., Wee, G.B., Srinivasan, N. and Kunaratnam, N. (1985) HLA and nasopharyngeal carcinoma in Malays. *British J. Cancer*; **51**: 3892
- Chen, H.F., Sauter, M., Haiss, P. and Muller, L.N. (1991) Immunological characterization of the Epstein-Barr virus phosphoprotein PP58 and deoxyribonuclease expressed in the baculovirus expression system. *Int. J. Cancer*; **48**: 879-888

- Chen, X., deV Pepper, S. and Arrand, J.R. (1992) Prevalence of the A and B types of Epstein-Barr virus DNA in nasopharyngeal carcinoma biopsies from Southern China. *J. Gen. Virol.*; **73**: 463-466
- Chen, Y. and Chan, S.H. (1994) Polymorphism of T-cell receptor genes in nasopharyngeal carcinoma. *Int. J. Cancer*; **56**: 830-833
- Chen-Chuan, L., Yu, M.C. and Henderson, B.E. (1985) Some epidemiologic observations of nasopharyngeal carcinoma in Guangdong, People's Republic of China. *Natl. Cancer Inst. Monogr.*; **69**: 49-52
- Cheng, H.M., Foong, Y.T., Sam, C.K., and Prasad, U. and Dillner, J. (1991) Epstein-Barr virus nuclear antigen 1 linear epitopes that are reactive with immunoglobulin A (IgA) or IgG in sera from nasopharyngeal carcinoma patients or from healthy donors. *J. Clin. Microbiol.*; **29**: 2180-2186
- Cheng, H.M., Foong, Y.T., Mathew, A., Sam, C.K., Dillner, J. and Prasad, U. (1993) Screening for nasopharyngeal carcinoma with an ELISA using the Epstein-Barr virus nuclear antigen, EBNA1: a complementary test to the IgA/VCA immunofluorescence assay. *J. Virol. Methods*; **42**: 45-52
- Cheng, H.M., Foong, Y.T., AbuSamah, A.J., Dillner, J., Sam, C.K. and Prasad, U. (1995) Linear epitopes of the replication-activator protein of Epstein-Barr virus recognised by specific serum IgG in nasopharyngeal carcinoma. *Cancer Immunol. Immunother.*; **40**: 251-256
- Cheung, A. and Kieff, E. (1982) Long internal repeats in Epstein-Barr virus DNA. *J. Virol.*; **44**: 286-294
- Chevallier-Greco, A., Manet, E., Chavrier, P., Mosnier, C., Daillie, J. and Sergeant, A. (1986) Both Epstein-Barr virus (EBV)-encoded transacting factors, EB1 and EB2, are required to activate transcription from an EBV early promoter. *EMBO J.*; **5**: 3243-3249
- Chou, P.Y. and Fasman G.D. (1974) Prediction of protein conformation. *Biochem.*; **133**: 222-45
- Chou, P.Y. and Fasman, G.D. (1977)  $\beta$ -turns in proteins. *J. Mol. Biol.*; **115**: 135-175

- Chou, P.Y. and Fasman, G.D. (1978) Prediction of the secondary-structure of proteins from their amino acid sequence. *Adv. Enzymol.*; **47**: 45-37
- Cochet, C., Martel -Renoir, D., Grunewald, V., Bosq, J., Cochet, G., Schwaab, G., Bernaudin, J-F. and Joab, I. (1993) Expression of the Epstein-Barr virus immediate early gene, BZLF1, in nasopharyngeal carcinoma tumour cells. *Virology*; **197**: 358-365
- Collins, S.P., Ball, G., Vonarx, E., Hosking, C., Shelton, M., Hill, D. and Howden, M.E.H. (1996) Absence of continuous epitopes in the house dust mite major allergens *Der p 1* from *Dermatophagoides pteronyssinus* and *Der f 1* from *Dermatophagoides farinae*. *Clin. Exp. Allergy*; **26**: 36-42
- Countryman, J. and Miller, G. (1985) Activation of expression of latent Epstein-Barr herpes virus after gene transfer with a sma;; clones subfragment of heterogenous viral DNA. *Proc. Natl. Acad. Sci. USA*; **82**: 4085-4089
- Countryman, J., Jenson, H., Seibl, R., Wolf, H. and Miller, G. (1987) Polymorphic proteins encoded within BZLF1 of defective and standard Epstein-Barr viruses disrupt latency. *Virology*; **61**: 3672-3679
- Crofts, N., Mashill, W. and Gust, I.D. (1988) Evaluation of enzyme-linked immunosorbent assays: a method of data analysis. *J. Virol. Meth.*; **22**: 51-59
- Dambaugh, T. and Kieff, E. (1982) Identification and nucleotide sequences of two similar tandem direct repeats in Epstein-Barr virus DNA. *J. Virol.*; **44**: 823-833
- Dambaugh, T., Hennessy, K., Chamnankit, L. and Keiff, E. (1984) U2 region of Epstein-Barr virus DNA may encode Epstein-Barr nuclear antigen 2. *Proc. Natl. Acad. Sci. USA*; **81**: 7632-7636
- Davies, D.R., Sheriff, S. and Padlan, E.A. (1988) Antibody-antigen complexes. *J. Biol. Chem.*; **263**: 10541-10544
- Davison, A.J. and Wilkie, N.M. (1981) Nucleotide sequences of the point between L and S segments of herpes simplex virus type 1 and 2. *J. Gen Virol.*; **55**: 315-331
- Davison, A.J. and Scott, E. (1986) The complete DNA sequence of varicella-zoster virus. *J. Gen. Virol.*; **67**: 1759-1816

- Davison, A.J. and Taylor, P. (1987) Genetic relations between varicella-zoster virus and Epstein-Barr virus. *J. Gen. Virol.*; **68**: 1067-1079
- Degos, L., Lepage, V., de-The, G., Blanc, H., Ferngold, N., Lu, S.T. and Zeng, T. (1984) HLA genotypes, GM genotypes, IgA anti-EBV antibodies in multiple family cases of nasopharyngeal carcinoma in China. **9th. International Histocompatibility Workshop and Conference, IARC, Vienna.** pp59
- de-Schryver, A., Klein, G., Henle, W. and Henle, G. (1974) EB virus-associated antibodies in Caucasian patients with carcinoma of the nasopharynx and in long term survivors after treatment. *Int. J. Cancer*; **13**: 319-25
- de-The, G., Ho, J.H.C., Ablashi, D.V., Day, N.E., Macario, A.J.L., Martin-Berthelon, M.C., Pearson, G.R. and Sohler, R. (1974) Nasopharyngeal carcinoma. IX. Antibodies to EBNA and correlation with response to other EBV antigens in Chinese patients. *Int. J. Cancer*; **16**: 713-721
- de-The, G., Ho, J.H.C. and Muir, C.S. (1982) Nasopharyngeal carcinoma. In: **Viral Infections of Humans: Epidemiology and Control** (Evans, A.S. ed.) Plenum Publ. Co., New York. 2nd. ed. pp. 621-652
- Dickson, R.I. (1981) Nasopharyngeal carcinoma: an evaluation of 209 patients. *Laryngoscope*; **91**: 333-354
- Dillner, J., Sternas, L., Kallin, B., Alexander, H., Ehlin-Hendriksson, B., Jornvall, H., Klein, G. and Lerner, R. (1984) Antibodies against a synthetic peptide identify the Epstein-Barr virus-determined nuclear antigen. *Proc. Natl. Acad. Sci. USA*; **81**: 4652-4656
- Dillner, J., Kallin, B., Ehlin-Hendriksson, Timar, L., and Klein, G. (1985) Characterization of a second Epstein-Barr virus determined nuclear antigen associated with the Bam H1 WYH region of EBV DNA. *Int. J. Cancer*; **35**: 359-366
- Dillner, J., Kallin, B., Alexander, H., Ernberg, I., Uno, M., Ono, Y., Klein, G. and Lerner, R.A. (1986) An Epstein-Barr virus (EBV) determined nuclear antigen (EBNA5) partly encoded by the transformation associated Bam WYH region of EBV DNA: preferential expression in lymphoblastoid cell lines. *Proc. Natl. Acad. Sci. USA*; **83**: 6641-6645

- Dillner, J., Szigetti, R., Henle, W., Henle, G., Lerner, R.A. and Klein, G. (1987) Cellular and humoral immune responses to synthetic peptides deduced from the amino acid sequences of Epstein-Barr virus-encoded proteins in EBV-transformed cells. *Int. J. Cancer*; **40**: 455-460
- Dillner, J. and Kallin, B. (1988) The Epstein-Barr virus proteins. *Adv. Cancer Res.*; **50**: 95-158
- Dolyniuk, M., Pritchett, R. and Kieff, E. (1976) Proteins of Epstein-Barr virus. I. Analysis of the polypeptide of purified enveloped Epstein-Barr virus. *J. Virol.*; **17**: 935-949
- Doran, D.M. (1990) Prediction of peptide epitopes in proteins. In: **Modern Methods in Protein- and Nucleic Acid Research Review Articles**, (Tschesche, H. ed.), Walter de Gruyter, Berlin. pp 231-250
- Dyrberg, T. and Oldstone, M.B.A. (1986) Peptides as antigens. Importance of orientation. *J. Expt. Med.*; **164**: 1344-49
- Epstein, M.A., Henle, G., Achong, B.G. and Barr, Y.M. (1964) Virus particles in cultured lymphoblasts from Burkitt's lymphoma. *Lancet i*: 702-703
- Epstein, M.A., Henle, G., Achong, B.G. and Barr, Y.M. (1965) Morphological and biological studies on a virus in cultured lymphoblasts from Burkitt's lymphoma. *J. Exp. Med.*; **12**: 761-770
- Epstein, M.A. and Achong, B.G. (1979) Introduction: discovery and general biology of the virus. In: **The Epstein-Barr Virus** (Epstein, M.A. and Achong, B.G. eds.), Springer-Verlag, Berlin, Heidelberg, New York. pp 1-22
- Ernberg, I., Kallin, B., Dillner, J., Falk, K., Ehlin-Hendrikson, B., Hammarskjold, M-L and Klein, G. (1986) Lymphoblastoid cell lines and Burkitt's lymphoma-derived cell lines differ in the expression of a second Epstein-Barr virus encoded nuclear antigen. *Int. J. Cancer*; **38**: 729-737
- Fahraeus, R., Jansson, A., Ricksten, A., Sjoblom, A. and Rymo, L. (1990) Epstein-Barr virus encoded nuclear antigen 2 activates the viral latent membrane protein promoter by modulating the activity of a negative regulatory element. *Proc. Natl. Acad. Sci. USA*; **87**: 7390-7398

- Falk, K., Gratama, J.W., Rowe, M., Zou, J.Z., Khanim, F., Young, L.S., Oosterveer, M.A.P. and Ernberg, I. (1995) The role of repetitive DNA sequences in the size variation of Epstein-Barr virus (EBV) nuclear antigens and the identification of different EBV isolates using RFLP and PCR analysis. *J. Gen. Virol.*; **76**: 779-790
- Farrel, P.J. (1989) Epstein-Barr virus genome. *Adv. Viral. Oncol.*; **8**: 103-132
- Feighny, R.J., Henry, B.F. and Pagano, J.S. (1981) Epstein-Barr virus polypeptides: effect of inhibition of viral DNA replication on their synthesis. *J. Virol.*; **37**: 61-71
- Fields, B.N. Knipe, D.M., Howley, P.M., Chanock, R.M., Melnick, J.L., Monath, T.P. Rozman, B. and Straus, S.E. (1996) *Fields Virology*, 3rd. ed. Lippincott-Raven, Philadelphia. pp 2343-2396
- Fisher, A., Fischer, G.O. and Cooper, E. (1984) Familial nasopharyngeal carcinoma. *Pathology*; **16**: 23-24
- Flemington, E. and Speck, S.H. (1990a) Identification of phorbol ester-response elements in the promoter of Epstein-Barr virus putative lytic switch gene BZLF1. *J. Virol.*; **64**: 1217-1226
- Flemington, E. and Speck, S.H. (1990b) Autoregulation of Epstein-Barr virus putative lytic switch gene BZLF1. *J. Virol.*; **64**: 1227-1232
- Fong, Y.Y. and Walsh, E.O. (1971) Carcinogenic nitrosamines in Cantonese salt-dried fish. *Lancet ii*; 1032
- Fong, Y.Y., Ho, J.H.C. and Huang, D.P. (1979) Preserved food as possible cancer hazards: WA rats fed salted fish have mutagenic urine. *Int. J. Cancer*; **23**: 542-546
- Foong, Y.T., Cheng, H.M., Sam, C.K., Dillner, J., Hinderer, W. and Prasad, U. (1990) Serum and salivary IgA antibodies against a defined epitope of the Epstein-Barr virus nuclear antigen (EBNA) are elevated in nasopharyngeal carcinoma. *Int. J. Cancer*; **45**: 1061-1064
- Fraga, S. (1982).. Theoretical prediction of protein antigenic determinants from amino acid sequences. *Can. J. Chem.*; **60**: 2606

- Funkhouser, S.W., Concannon, P., Charmley, P., Vredevoe, D.L. and Hood, L. (1992) Differences in T cell receptor restriction fragments length polymorphisms in patients with rheumatoid arthritis. *Arthritis Rheum.*; **35**: 465-471
- Gan, Y.Y., Fones-Tan, A., Chan, S.H. and Gan, L.H. (1996) Epstein-Barr viral antigens used in the diagnosis of nasopharyngeal carcinoma. *J. Biomed. Sci.*; **3**: 159-169
- Gerber, P., Nonoyama, M., Lucas, S., Perein, E. and Goldstein, L.I. (1972) Oral excretion of Epstein-Barr virus by healthy subjects and patients with infectious mononucleosis. *Lancet ii*: 988-989
- Gergely, L., Klein, G. and Ernberg, I. (1971) Host cell macromolecular synthesis in cells containing EBV-induced early antigens studied by combined immunofluorescence and radioautography. *Virology*; **45**: 22-29
- Gergely, L., Sternas, L., Dillner, J. and Kelen, G. (1984) Molecular size variation of EBNA is determined by the Epstein-Barr virus-negative lymphomas converted with different EBV strains and somatic hybrids. *Intervirolog.*; **22**: 85-96
- Getzoff, E.D., Tainer, J.A. and Lerner, R.A. (1988) The chemistry and mechanism of antibody binding to protein antigens. *Adv. Immun.*; **43**: 1-98
- Geysen, H.M., Meloen, R.H. and Barteling, S.J. (1984) Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid. *Proc. Natl. Acad. Sci. USA*; **81**: 3998-4002
- Geysen, H.M. (1985) Antigen-antibody interactions at the molecular level: adventures in peptide synthesis. *Immunol. Today*; **6**: 364-69
- Geysen, H.M., Barteling, S.J. and Meloen, R.H. (1985) Small peptides induce antibodies with a sequence and structural requirement for binding antigen comparable to antibodies raised against the native protein. *Proc. Natl. Acad. Sci. USA*; **82**: 178-182
- Geysen, H.M., Rodda, S.J. and Mason, T.J. (1986) A priori delineation of a peptide which mimics a discontinuous antigenic determinant. *Mol. Immunol.*; **23**: 709-715

- Geysen, H.M., Rodda, S.J., Mason, T.J., Tribbick, G. and Schoofs, P.G. (1987a) Strategies for epitope analysis using peptide synthesis. *J. Immunol. Meth.*; **102**: 259-74
- Geysen, H.M., Tainer, J.A., Rodda, S.J., Masom, T.J., Alexander, H., Getzoff, E.D. and Lerner, R.A. (1987b) Chemistry of antibody binding to a protein. *Science*; **235**: 1184-1190
- Geysen, H.M., Mason, T.J. and Rodda, S.J. (1988) Cognitive features of continuous antigenic determinants. *J. Molecular Recognition*; **1**: 32-41
- Gnann, J.W., McCormick, J.B., Mitchell, S., Nelson, J.A. and Oldstone, M.B.A. (1987) Synthetic peptide immunoassay distinguishes HIV type 1 and HIV type 2 infections. *Science*; **237**: 1346-1349
- Gradoville, L., Grogan, E., Taylor, N. and Miller, G. (1990) Differences in the extent of activation of Epstein-Barr virus replicative gene expression among four non-producer cell lines stably transformed by OriP/BZLF1 plasmids. *Virology*; **178**: 345-354
- Guo, Y.Q., Liang, J.S., Zheng, S.A., Ou, X.T. and Zong, Y.S. (1996) Evaluation of EBV serological screening for detection of nasopharyngeal carcinoma (NPC) in Zhongshan city, southern China. Proceedings 3rd. Hong Kong International Cancer Congress and 7th. International EBV Meeting, Hong Kong. p177
- Hammerschmidt, W. and Sugden, B. (1989) Genetic analysis of immortalizing functions of Epstein-Barr virus in human B lymphocytes. *Nature*; **340**: 393-397
- Hampar, B., Derge, J., Martos, L., Tagemets, M., Chang, S. and Chakrabarty, M. (1973) Identification of a critical period during the S phase for activation of the Epstein-Barr virus by 5-iododeoxyuridine. *Nature (London) New Biol.*; **244**: 214-217
- Heller, M., Henderson, A. and Kieff, E. (1982) A repeat array in Epstein-Barr virus is related to cell DNA sequences interspersed on human chromosomes. *Proc. Natl. Acad. Sci. USA*; **79**: 5916-5920
- Henderson, B.E., Louie, E., Bugdanoff, E., Henle, W., Alenat, B. and Henle, G. (1974) Antibodies to herpes group viruses in patients with nasopharyngeal carcinoma and other head and neck cancers. *Cancer Res.*; **34**: 1207-1210

- Henderson, B.E. and Louie, E. (1978) Discussion of risk factors for nasopharyngeal carcinoma In: **Nasopharyngeal carcinoma: Etiology and control** (de-The, G. and Ito, Y. eds), IARC, Lyon. pp 251-55
- Henle, G. and Henle, W. (1966a) Studies on cell lines derived from Burkitt's lymphoma. *Trans. N. Y. Acad. Sci.*; 29: 71-79
- Henle, G. and Henle, W. (1966b) Immunofluorescence in cells derived from Burkitt's lymphoma. *J. Bacteriol.*; 91: 1248-1256
- Henle, G. and Henle, W. (1967) Immunofluorescence, interference and complement fixation technique in the detection of the herpes-type virus in Burkitt's tumour cell lines. *Cancer Res.*; 27: 2442-2446
- Henle, W., Henle, G. and Diehl, V. (1968) Relation of Burkitt's tumour-associated herpes-type virus to infectious mononucleosis. *Proc. Natl Acad. Sci. USA*; 59: 94-107
- Henle, W., Henle, G., Burtin, P., Cachin, Y., Clifford, P., de-Schryver, A., de-The, G., Diehl, V., Ho, H.C. and Klein, G. (1970) Antibodies to Epstein-Barr virus in nasopharyngeal carcinoma, other head and neck neoplasms and control groups. *J. Natl. Cancer Inst.*; 44: 225-31
- Henle, G., Henle, W. and Klein, G. (1971) Demonstration of two distinct components in the early antigen complex of Epstein-Barr virus-infected cells. *Int. J. Cancer*; 8: 272-278
- Henle, W., Ho, H.C., Henle, G. and Kwan, H.C. (1973) Antibodies to Epstein-Barr virus-related antigens in nasopharyngeal carcinoma. Comparison of active cases with long term survivors. *J. Natl. Cancer Inst.*; 51: 361-69
- Henle, G., Henle, W. and Horwitz, C.A. (1974) Antibodies to Epstein-Barr virus-associated nuclear antigen in infectious mononucleosis. *J. Infect. Dis.*; 130: 231-239
- Henle, G. and Henle, W. (1976) Epstein-Barr virus-specific IgA serum antibodies as an outstanding features of nasopharyngeal carcinoma. *Int. J. Cancer*; 17: 1-7

- Henle, W., Ho, J.H., Henle, G., Chau, J.C. and Kwan, H.C. (1977) Nasopharyngeal carcinoma: significance of changes in Epstein-Barr virus-related antibody patterns following therapy. *Int. J. Cancer*; **20**: 663-72
- Henle, G. and Henle, W. (1979) Seroepidemiology of the virus. In: **The Epstein-Barr Virus** (Epstein, M.A. and Achong, B.G. eds.), Springer, Berlin, Heidelberg and New York. pp 161-178
- Hennessy, K., Heller, M., van Santer, V. and Kieff, E. (1983) Simple repeat array in Epstein-Barr virus DNA encodes part of the Epstein-Barr nuclear antigen. *Science*; **220**: 1396-1398
- Hennessy, K. and Kieff, E. (1985) A second nuclear protein is encoded by Epstein-Barr virus in latent infection. *Science*; **227**: 1238-1240
- Hennessy, K., Fennewald, S. and Kieff, E. (1985) A third viral nucleoprotein in lymphoblasts immortalized by Epstein-Barr virus. *Proc. Natl. Acad. Sci. USA*; **82**: 5944-5948
- Hennessy, K., Wang, F., Bushman, E.W. and Kieff, E. (1986) Definitive identification of a member of the Epstein-Barr virus nuclear protein 3 family. *Proc. Natl. Acad. Sci. USA*; **83**: 5693-5697
- Hinderer, W., Gorgievski-Hrisoho, M., Nebel-Schikel, H., Horn, J., Vornhagen, R., Sonneborn, H.H., Sigel, G. and Wolf, H. (1990) Diagnosis of Epstein-Barr virus infection using recombinant viral proteins. Poster, VIIIth. Intl. Congress of Virology, Berlin
- Hinderer, W., Vornhagen, R., Sonneborn, H.H., Ng, M.H. and Wolf, H. (1996) Follow-up of EBV serology in NPC patients using recombinant EA, EBNA and VCA ELISAs. 3rd. Hong Kong International Cancer Congress and 7th. International EBV Meeting, Hong Kong. p180
- Ho, J.H.C. (1971) Genetic and environmental factors in nasopharyngeal carcinoma. In: **Recent Advances in Human Tumour Virology and Immunology** (Nakahara, W., Nishioka, K., Hirayama, T. and Ito, Y. eds.), Tokyo Univ. Press, Tokyo. pp 275-295
- Ho, J.H.C. (1972) Nasopharyngeal carcinoma (NPC). *Adv. Cancer Res.*; **15**: 57-92

- Ho, H.C. (1978) An epidemiologic and clinical study of nasopharyngeal carcinoma. *Int. J. Rad. Oncol. Biol. Phys.*; **4**: 181-98
- Ho, H.C., Ng, M.H., Kwan, H.C. and Chau, J.C.W. (1976) Epstein-Barr virus-specific IgA and IgG serum antibodies in nasopharyngeal carcinoma. *Br. J. Cancer*; **34**: 655-60
- Ho, H.C., Kwan, H.C., Wu, P., Chan, S.K., Ng, M.H. and Saw, D. (1978) Epstein-Barr antibodies in suspected nasopharyngeal carcinoma. *Lancet*; **2**: 1094-1095
- Ho, J.H.C., Lau, W.H., Kwan, H.C., Chan, C.L., Au, G.K.H., Sau, D. and de-The, G. (1981) Diagnostic and prognostic serological markers in NPC. In: **Cancer Campaign Vol. 1** (Grundmann, E., Krueger, G.R.F. and Ablashi, D.V. eds.), Gustav, Fisher Verlag Stuttgart, New York. pp 219-224
- Hopp, T.P. and Woods, K.R. (1981) Prediction of protein antigenic determinants from amino acid sequences. *Proc. Natl. Acad. Sci. USA*; **78**: 3824-28
- Hopp, T.P. and Woods, K.R. (1983) A computer program for predicting protein antigenic determinants and other interaction sites. *Mol. Immunol.*; **20**: 483-489
- Horal, P., Svennerholm, B., Jeansson, S., Rymo, L., Hall, W.W. and Vahine, A. (1991) Continuous epitopes of the human immunodeficiency virus type 1 (HIV-1) transmembrane glycoprotein and reactivity of human sera to synthetic peptides representing various HIV-1 isolates. *J. Virol.*; **65**: 2718-2723
- Houghten, R.A. (1985) General method for the rapid solid phase synthesis of large numbers of peptides. Specificity of antigen-antibody interaction at the level of individual amino acids. *Proc. Natl. Acad. Sci. USA*; **82**: 3824
- Houghten, R.A., DeGraw, S.T., Bray, M.K., Hoffman, S.R. and Frizzel, N.D. (1986) Simultaneous multiple peptide synthesis. The rapid preparation of large numbers of discrete peptides for biological, immunological and methodological studies. *Bio. Techniques*; **4**: 522
- Huang, D.P., Ho, J.H.C., Henle, W. and Henle, G. (1974) Demonstration of Epstein-Barr virus-associated nuclear antigen in nasopharyngeal carcinoma cells from fresh biopsies. *Int. J. Cancer*; **14**: 580-88

- Huang, D.P., Ho, J.H.C. and Gough, T.A. (1978) Analysis for volatile nitrosamines in salt-preserved food stuffs traditionally consumed by Southern Chinese. In: **Nasopharyngeal Carcinoma: Etiology and Control** (de-The, G. and Ito, Y. eds.), IARC, Lyon. pp 309-314
- Ito, Y., Yanase, S., Tokuda, H., Kishishita, M., Ohigashi, H., Hiroto, M. and Koshimizu, K. (1982) Epstein-Barr virus activator by tung oil, extracts of *Aleurites Fordii* and its diterpene ester 12-o-hexadecanoyl-16-hydroxyphorbol-13-acetate. **Cancer Letters**; **18**: 87-95
- Ito, Y., Tanimoto, M., Kamura, H., Yoneda, M., Morishima, Y., Takatsuki, K., Itatsu, T. and Saito, H. (1988) Association of HLA-DR phenotypes and T-lymphocyte receptor  $\beta$ -chain region RFLP with IDDM in Japanese. **Diabetes**; **37**: 1633-1636
- Joab, I., Rowe, D.T., Bodescot, M., Nicolas, J.C., Farrel, P.J. and Perricaudet, M. (1987) Mapping of the gene coding for Epstein-Barr virus-determined nuclear antigen EBNA3 and its transient over expression in an human cell line by using an Adenovirus expression vector. **J. Virol.**; **61**: 3340-3344
- Joab, I., Nicolas, J.C., Schwaab, G., de The, G., Clause, B., Perricaudet, M. and Zeng, Y. (1991) Detection of anti-Epstein-Barr virus transactivator, ZEBRA, antibodies in sera from patients with nasopharyngeal carcinoma. **Int. J. Cancer**; **48**: 647-649
- Jones, M.D., Foster, L., Sheddy, T. and Griffin, B. (1984) The EB virus genome in Daudi Burkitt's lymphoma cells has a deletion similar to that observed in a non-transforming strain (P3HR-1) of the virus. **EMBO J.**; **3**: 813-821
- Kabat, E.A. (1970) Heterogeneity and structure of antibody-combining sites. **Ann. N.Y. Acad. Sci.**; **169**: 43-54
- Kallin, B., Luka, J. and Klein, G. (1979) Immunochemical characterization of Epstein-Barr virus (EBV)-associated early and late antigens in n-butyrate treated P3HR-1 cells. **J. Virol.**; **32**: 710-716
- Karimi, L., Crawford, D.H., Speck, S. and Nicholson, L.J. (1995) Identification of an epithelial cell differentiation responsive region within the BZLF1 promoter of the Epstein-Barr virus. **J. Gen. Virol.**; **76**: 759-765

- Katz, B.Z., Raab-Traub, N. and Miller, G. (1989) Latent and replicating forms of Epstein-Barr virus DNA in lymphomas and lymphoproliferative disease. *J. Infect. Dis.*; **160**: 589-598
- Kemp, B., Rylatt, D., Bundesen, P., Doherty, R., McPhee, A., Stapleton, D., Cottis, L., Wilson, K., John, M., Kjan, J., Dihn, D., Miles, S. and Hillyard, C. (1988) *Science*; **241**: 1352-1354
- Kenny, S., Kamine, J., Holley-Guthrie, E., Lin, J.C., Mar, E.C. and Pagano, J. (1989) The Epstein-Barr virus (EBV) BZLF1 immediate-early gene product differentially affects latent versus productive EBV promoters. *J. Virol.*; **63**: 1729-1736
- Khanna, R., Burrows, S.R. and Moss, D.J. (1995) Immune regulation in Epstein-Barr virus-associated diseases. *Microbiological Reviews* **59**; (3): 387-405
- Kieff, E. and Liebowitz, D. (1990) Epstein-Barr virus and its replication. In: *Virology* (Fields, B.N. and Knipe, D.M. eds.), 2nd. ed. Raven Press, New York. pp1889-1919
- Kintner, C. and Sugden, B. (1981) Conservation and progressive methylation of Epstein-Barr virus DNA sequences in transformed cells. *J. Virol.*; **38**: 305-316
- Klein, G. (1983) The Epstein-Barr virus. In: *The Herpes Viruses* (Kaplan, A.S. ed.), Academic Press, New York. pp 521-555
- Kyte, J. and Doolittle, R.F. (1982) A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.*; **157**: 105-132
- Lanier, A.P., Henle, W., Bender, T.R., Henle, G. and Talbot, M.L. (1980) Epstein-Barr virus-specific antibody titers in seven Alaskan natives before and after diagnosis of nasopharyngeal carcinoma. *Int. J. Cancer*; **26**: 133-37
- Lau, R., Packham, G. and Farrel, P.J. (1992) Differential splicing of Epstein-Barr virus immediate-early RNA. *J. Virol.*; **66**: 6233-6236
- Lear, A.L., Rowe, M., Kurilla, M.G., Lee, S., Henderson, S.S., Kieff, E. and Rickinson, A.B. (1992) The Epstein-Barr virus (EBV) nuclear antigen 1 Bam H1 F promoter is activated on entry of EBV-transformed B cells into the lytic cycle. *J. Virol.*; **66**: 7461-7468

- Lee, H.P., Day, N.E. and Shanmugaratnem, K. (1988) Trends in Cancer Incidence in Singapore 1968-1982 (with attached appendix). IARC, Lyon.
- Lees, J.F., Arrand, J.E., Pepper, S., Stewart, J.P., Mackett, M. and Arrand, J.R. (1993) The Epstein-Barr virus candidate vaccine antigen gp340/220 is highly conserved between virus types A and B. *Virology*; **195**: 578-586
- Leinikki, P., Lehtinen, M., Hyoty, H., Parkkonen, P., Kantanen, M-L. and Hakulinen, J. (1993) Synthetic peptides as diagnostic tools in virology. *Adv. Cancer Res.*; **40**: 149-185
- Levine, P.H., Connely, R.R. and McKay F.W. (1983) The influence of residence, race and place of birth on the incidence and mortality in nasopharyngeal carcinoma. In: **Nasopharyngeal Carcinoma: Current Concepts** (Prasad, U., Ablashi, D.V., Levine, P.H. and Pearson, G.R. eds.), Univ. Malaya Press, Kuala Lumpur. pp143-156
- Levitt, M. (1976) A simplified representation of protein conformations for rapid simulation of protein folding. *J. Mol. Biol.*; **104**: 59-107
- Liang, J.S., Ou, X.T., Guo, Y.Q., Zheng, S.A. and Zong, Y.S. (1996) Comparative study of detection rates for nasopharyngeal carcinoma (NPC) between EBV seropositive and seronegative subjects. Proceedings 3rd. Hong Kong International Cancer Congress and 7th. International EBV Meeting, Hong Kong. p176
- Lin, T.M., Chen, K.P., Lin, C.C., Hsu, M.M., Chiang, T.C. Jung, P.F. and Hirayama, T. (1973) Retrospective study on nasopharyngeal carcinoma. *J. Natl. Cancer Inst.*; **51**: 1403-1408
- Lin, J.C., Shaw, J.E., Smith, M.C. and Pagano, J.S. (1979) Effect of 12-o-tetradecanoyl-phorbol-13-acetate on the replication of EBV. 1. Characterization of viral DNA. *Virology*; **99**: 183-187
- Lin, J.C., Etsuyo, I. and Pagano J.S. (1985) Qualitative and quantitative analyses of Epstein-Barr virus early antigen diffuse component by western blotting enzyme-linked immunosorbent assay with a monoclonal antibody. *J. Virol.*; **53**: 793-799

- Lindahl, T., Adams, A., Bjursell, G., Bornkamm, G.W., Kaschka-Dierich and Jehn, V. (1976) Covalently closed circular duplex DNA of Epstein-Barr virus in human lymphoid cell line. *J. Mol. Biol.*; **102**: 511-530
- Littler, E., Bayliss, S.A., Connolly, Y., Conway, M.J., Mackett, M., Chen, X., Zeng, Y. and Arrand, J.R. (1990) Diagnosis of nasopharyngeal carcinoma using Epstein-Barr virus coded alkaline DNase, membrane antigen and thymidine kinase. In: **Epstein-Barr Virus and Human Diseases** (Ablashi, D.V., Huang, A.T., Pagano, J.S., Pearson, G.R. and Young, C.S. eds.), Humana Press, Clifton, New Jersey. pp 353-356
- Lu, S.J., Day, N.E., Degos, L., Lepage, V., Wang, P.C., Chan, S.H., Simmons, M., McKnight, B., Easton, D., Zeng, Y. and de-The, G. (1990) Linkage of a nasopharyngeal carcinoma susceptibility locus to the HLA region. *Nature*; **346**: 470-71
- Luka, J., Kallin, B. and Klein, G. (1979) Induction of the Epstein-Barr virus (EBV) cycle in latently infected cells by n-butyrate. *Virology*; **94**: 228-231
- Maeji, N.J., Bray, A.M., Valerio, R.M. and Wang W. (1995) Larger scale multipin peptide synthesis. *Peptide Research*; **8**: 33-38
- Magrath, I. (1990) The pathogenesis of Burkitt's lymphoma. *Adv. Cancer Res.*; **55**: 133-270
- Martel-Lenoir, D., Grunewald, V., Touitou, R., Schwaab, G. and Joab, I. (1995) Qualitative analysis of the expression of Epstein-Barr virus lytic genes in nasopharyngeal carcinoma biopsies. *J. Gen. Virol.*; **76**: 1401-1408
- Mathew, A., Cheng, H.M., Sam, C.K., Joab, I., Prasad, U. and Cochet, C. (1994) A high incidence of serum IgG antibodies to the Epstein-Barr virus replication activator protein in nasopharyngeal carcinoma. *Cancer Immunol. Immunother.*; **38**: 68-70
- Maxfield, F.R. and Scheraga, H.A. (1976) Status of empirical methods for the prediction of protein backbone topography. *Biochem.*; **15**: 5138-5153
- Merrifield, R.B. (1963) Solid phase peptide synthesis. I. The synthesis of a tetrapeptide. *J. Am. Chem. Soc.*; **85**: 2149

- Middeldorp, J.M. and Meloen, R.H. (1988) Epitope-mapping of the Epstein-Barr virus major capsid protein using systematic synthesis of overlapping oligopeptides. *J. Virol. Methods*; **21**: 147-159
- Miller, G. (1990) Epstein-Barr virus. Biology, pathogenesis and medical aspects. In: *Virology* (Fields, B.N. and Knipe, D.M.M. eds.), 2nd. ed., Raven Press, New York. pp 1921-1958
- Montalvo, E.A., Shi, Y., Shenk, T.E. and Levine A.J. (1991) Negative regulation of the BZLF1 promoter of Epstein-Barr virus. *J. Virol.*; **65**: 3647-3655
- Morgan, D.G., Niederman, J.C., Miller, G., Smith, H.W. and Dowaliby, J.M. (1979) Site of Epstein-Barr virus replication in the oropharynx. *Lancet*; **II**: 1154-57
- Moss, D.J., Sculley, T.B. and Pope, J.H. (1986) Induction of Epstein-Barr virus antigens. *J. Virol.*; **58**: 988-90
- Muir, C.S. (1971) NPC in non-Chinese populations with special reference to South East Asia and Africa. *Int. J. Cancer*; **8**: 351-363
- Murray, R.J., Kurilla, M.G., Griffin, H.M., Brooks, J.M., Mackett, M., Arrand, J., Rowe, M., Burrows, S., Moss, D.J., Kieff, E. and Rickinson, A.B. (1990) Human cytotoxic T cell responses against Epstein-Barr virus nuclear antigens demonstrated using recombinant vaccinia virus. *Proc. Natl. Acad. Sci. USA*; **87**: 2906-2910
- Murray, R.J., Kurilla, M.G., Brooks, J.M., Thomas, W.A., Rowe, M., Kieff, M.E. and Rickinson, A.B. (1992) Identification of target antigens for the human cytotoxic T response to Epstein-Barr virus (EBV): implications for the immune control of EBV-positive malignancies. *J. Exp. Med.*; **176**: 157-168
- Neel, H.B. and Taylor, W.F. (1983) Clinical presentation and diagnosis of NPC: Current status In: *Nasopharyngeal Carcinoma - Current Concepts* (Prasad U, Ablashi DV, Levine PH & Pearson GR eds.), Univ Malaya Press, Kuala Lumpur. pp 1-10
- Norhanom, Abdul-Wahab, (1989) Epidemiology and Immunology of Nasopharyngeal Carcinoma. PhD. Thesis, Univ. Malaya, Kuala Lumpur

- Novotny, J., Handshumacher, M., Haaber, E., Bruccoleri, R.E., Carlson, W.B., Fanning, D.W., Smith, J.A. and Rose, G.D. (1986) Antigenic determinants in proteins coincide with surface regions accessible to large probes (antibody domains). *Proc. Natl. Acad. Sci. USA* **83**; 226-230
- Nyormoi, O., Thorley-Lawson, D.A., Elkington, J and Strominger, J.L. (1976) Differential effect of phosphonoacetic acid on the expression of Epstein-Barr viral antigens and virus production. *Proc. Natl. Acad. Sci. USA*; **73**: 1745-1748
- Old, L.J., Boyse, E.A., Oettgen, H.E., De Harven, E., Geering, G., Williamson, B. and Clifford, P. (1966) Precipitating antibody in human sera to an antigen present in cultured Burkitt's lymphoma. *Proc. Natl. Acad. Sci. USA*; **56**: 1699-1704
- Oshima, M. and Atassi, M.Z. (1989) Comparison of peptide-coating conditions in solid phase plate assays for detection of anti-peptide antibodies. *Immunological Investigations*; **18(7)**: 841-851
- Pallesen, G., Hamilton-Dutoit, S.J., Rowe, M., Lisse, I., Ralfkiaer, E., Sandvej, K. and Young, L. (1990) Expression of Epstein-Barr virus replicative protein in AIDS-related non-Hodgkin's lymphoma cells. *J. Pathol.*; **165**: 289-299
- Pallesen, G., Hamilton-Dutoit, S.J. and Zhou, X. (1993) The association of Epstein-Barr virus (EBV) with T cell lymphoproliferations and Hodgkin's disease: two new developments in the EBV field. *Adv. Cancer Res.*; **62**: 179-239
- Pathmanathan, R., Prasad, U., Sadler, R., Flynn, K. and Raab-Traub, N. (1995a) Clonal proliferations of cells infected with Epstein-Barr virus in preinvasive lesions related to nasopharyngeal carcinoma. *New. Engl. J. Med.*; **333**: 693-698
- Pathmanathan, R., Prasad, U., Chandrika, G., Sadler, R., Flynn, K. and Raab-Traub, N. (1995b) Undifferentiated, nonkeratinizing, ans squamous cell carcinoma of the nasopharynx. Variants of Epstein-Barr virus-infected neoplasia. *Am. J. Pathol.*; **146**: 1355-1367
- Pearson, G.R., Henle, G. and Henle, W. (1971) Production of antigens associated with Epstein-Barr virus in experimentally infected lymphoblastoid cell lines. *J. Natl. Cancer Inst.*; **46**: 1243-1250

- Pearson, G.R. (1980) Epstein-Barr virus immunology In: **Viral Oncology** (Klein, G. ed.), Raven Press, New York. pp 739-767
- Pearson, G.R., Vroman, B., Chase, B., Sculley, T., Hummel, M. and Kieff, E. (1983a) Identification of polypeptide components of the Epstein-Barr virus early antigen complex with monoclonal antibodies. *J. Virol.*; **47**: 193-201
- Pearson, G.R., Weiland, L.H., Neel, H.B., Taylor, W., Earle, J., Mulroney, S.E., Goepfert, H., Lanier, A., Talvot, M.L., Pilch, B., Goodman, M., Huang, A., Levine, P.H., Hyams, V., Moran, E., Henle, G. and Henle, W. (1983b) Application of Epstein-Barr virus (EBV) serology to the diagnosis of North American nasopharyngeal carcinoma. *Cancer*; **51**: 260-268
- Pearson, G.R. and Luka, J. (1986) Characterization of virus determined antigens. In: **The Epstein-Barr Virus: Recent Advances** (Epstein, M.A. and Achong, B.G. eds.), Heinemann, London. pp 48-73
- Permeen, A.M.Y., Sam, C.K., Paathmanathan, R., Prasad, U. and Wolf, H. (1990) Detection of Epstein-Barr virus DNA in nasopharyngeal carcinoma using non-radioactive digoxigenin-labelled probe. *J. Virol. Methods*; **27**: 261-268
- Petti, L. and Kieff, E. (1988) A sixth Epstein-Barr virus nuclear protein (EBNA 3B) is expressed in latently infected growth-transformed lymphocytes. *J. Virol.*; **62**: 2173-2178
- Petti, L., Sample, L., Wang, F. and Kieff, E. (1988) A fifth Epstein-Barr virus nuclear protein (EBNA 3C) is expressed in latently infected growth-transformed lymphocytes. *J. Virol.*; **62**: 1330-1338
- Poirier, S., Ohshima, H., de-The, G., Hunert, A., Bourgade, M.C. and Bartsch, H. (1987) Volatile nitrosamine levels in common food from Tunisia, South China and Greenland, high-risk areas for nasopharyngeal carcinoma (NPC). *Int. J. Cancer*; **39**: 293-296
- Pope, J.H., Horne, M.K. and Scott, W. (1968) Transformation of foetal human leukocytes in vitro by filtrates of a human leukaemic cell line containing herpes-like virus. *Int. J. Cancer*; **3**: 857-866
- Prasad, U. (1970) Problems in the early diagnosis of nasopharyngeal carcinoma. Proceedings 5th. Malaysia-Singapore Congress Medicine; **5**: 248-51

- Prasad, U. (1979) Fossa of Rosenmuller and nasopharyngeal carcinoma. *Med. J. Malaysia*; **33**: 222-25
- Prasad, U., Ablashi, D.V., Prathap, K., Yadav, M., Singaram, S.P., Singh, P. and Singh, J. (1983) Problem of occult primary in nasopharyngeal carcinoma. In: **Nasopharyngeal Carcinoma: Current Concepts** (Prasad, U., Ablashi, D.V., Levine, P.H. and Pearson, G.R. eds.), Univ. Malaya Press, Kuala Lumpur. pp 11-15
- Prathab, K., Prasad, U. and Ablashi, D.V. (1983) The pathology of nasopharyngeal carcinoma in Malaysians. In: **Nasopharyngeal Carcinoma: Current Concepts** (Prasad, U., Ablashi, D.V., Levine, P.H. and Pearson, G.R. eds.), Univ. Malaya Press, Kuala Lumpur. pp 55-63
- Prophet, D.N. (1984) Detection of immunoglobulin polymorphisms. In: **Detection of Immune-Associated Genetic Markers of Human Diseases** (Simon, M.J. and Tait, B.D. eds), Edinburgh, Churchill, Livingstone. pp 65-81
- Radford, A.J., Wood, P.R., Billman-Jacobe, H., Geysen, H.M., Mason, T.J. and Tribbick, G. (1990) Epitope mapping of the *Mycobacterium bovis* secretory protein MPB70 using overlapping peptide analysis. *J. Gen. Microbiol.*; **136**: 265-272
- Rawlins, D., Milman, G., Hayward, S. and Hayward, G. (1985) Sequence-specific DNA binding of the Epstein-Barr virus nuclear antigen EBNA1 to clustered site in the plasmid maintenance region. *Cell*; **42**: 859-868
- Reedman, B.M. and Klein, G. (1973) Cellular localization of an Epstein-Barr virus (EBV) associated complement-fixing antigen in producer and non-producer lymphoblastoid cell lines. *Int. J. Cancer*; **11**: 499-520
- Ren, E.C., Law, C.T. and Chan, S.H. (1995) HLA-A2 allelic microvariants in nasopharyngeal carcinoma. *Int. J. Cancer*; **63**: 213-215
- Rhodes, G., Houghten, R., Taulane, J.P., Carson, D. and Vaughan, J. (1984) The immune response to Epstein-Barr nuclear antigen: conformational and structural features of antibody binding to synthetic peptides. *Mol. Immunol.*; **21** (11): 1047-1054

- Rickinson, A.B. (1986) Cellular immunological response to the virus infection. In **Epstein-Barr Virus: Recent Advances** (Epstein, M.A. and Achong, B.G. eds.), Heinemann, London. pp75-125
- Rickinson, A.B., Jarvis, J.E., Crawford, D.H. and Epstein, M.A. (1974) Observation on the type of infection by Epstein-Barr virus in peripheral lymphoid cells of patients with infectious mononucleosis. *Int. J. Cancer*; **14**: 705-715
- Rickinson, A.B., Rowe, M., Hart, I.J., Yao, Q.Y., Henderson, L.E., Rabin, H. and Epstein, M.A. (1984) T-cell mediated regression of 'spontaneous' and of Epstein-Barr virus induced B-cell transformation in vitro: studies with cyclosporin A. *Cell. Immunol.*; **87**: 646-658
- Rickinson, A.B., Yao, Q.Y. and Wallace, L.E. (1985) The Epstein-Barr virus as a model of virus-host interactions. *British Med. Bull.*; **41**: 75-79
- Rickinson, A.B., Young, L.S. and Rowe, M. (1987) Influence of the Epstein-Barr virus nuclear antigen EBNA2 on the growth phenotype of virus-transformed B cells. *J Virol.*; **61**: 1310-1317
- Ricksten, A., Kallin, B., Alexander, H., Dillner, J., Fahraeus, R., Klein, G., Lerner, R. and Rymo, L. (1988) Bam H1E region of the Epstein-Barr virus genome encodes three transformation associated nuclear proteins. *Proc. Natl. Acad. Sci. USA*; **85**: 995-999
- Rogers, R.P., Strominger, J.L. and Speck, S.H. (1992) Epstein-Barr virus in B lymphocytes: viral gene expression and function in latency. *Adv. Cancer Res.*; **58**: 1-26
- Roivainen, M., Narvanen, A., Korkolainen, M., Huhtala, M.L. and Hovi, T. (1991) Antigenic region of poliovirus type 3/sabin capsid proteins recognized by human sera in the peptide scanning technique. *Virology*; **180**: 99-107
- Rose, G.D. and Roy, S. (1980) Hydrophobic basis of packing in globular proteins. *Proc. Natl. Acad. Sci. USA*; **77**: 4643-4647
- Rothwell, R.I. (1978) Carcinoma of the nasopharynx in Sabah. *South East Asian J. Surgery*; 88-95,

- Rowe, M., Finke, J., Szigeti, R. and Klein, G (1988) Characterization of the serological responses in man to the latent membrane protein and the six nuclear antigens encoded by Epstein-Barr virus. *J. Gen. Virol.*; **69**: 1217-1228
- Rymo, L., Klein, G. and Ricksten, A. (1985) Expression of a second Epstein-Barr virus nuclear antigen in mouse cells after gene transfer with a cloned fragment of the viral genome. *Proc. Natl. Acad. Sci. USA*; **82**: 3435-3439
- Saint-Clair, A.J., Brimmell, M. and Farrell, P.J. (1992) Reciprocal antagonism of steroid hormones and BZLF1 in switch between Epstein-Barr virus latent and productive cycle gene expression. *J. Virol.* **66**: 70-77
- Scopes, G.E., Watt, P.J. and Laambden, P.R. (1990) Identification of a linear epitope on the fusion glycoprotein of respiratory syncytial virus. *J. Gen. Virol.*; **71**: 53-59
- Sam, C.K., Prasad, U. and Pathmanathan, R. (1989) Serological markers in the diagnosis of histopathological types of nasopharyngeal carcinoma. *Eur. J. Surg. Oncol.*; **15**: 357-360
- Sam, C.K., Abu-Samah, A.J. and Prasad, U. (1994) IgAVCA as a follow-up marker in the monitoring of nasopharyngeal carcinoma. *Eur. J. Surg. Oncol.*; **20**: 261-564
- Sample, J., Young, L.S., Martin, B., Chatman, T., Rickinson, A.B. and Keiff E. (1990) Epstein-Barr types 1 and 2 differ in their EBNA-3A, EBNA- 3B and EBNA-3C genes. *J. Virol.*; **64**: 4084-4092
- Saul, A.J. and Geysen, H.M. (1990) Identification of epitopes through peptide technology. In: **New Generation Vaccines** (Woodrow, G.C. and Levine, M.M. eds.), Dekker, New York. pp 117-126
- Savoca, R., Schwaab, C. and Bosshard, H.R. (1991) Epitope mapping employing immobilized synthetic peptides. How specific is the reactivity of these peptides with antiserum raised against the parent protein? *J. Immunol. Methods*; **141**: 245-252
- Scanlon, P.W., Rhodes, R.E., Woolner, L.B. Levine, K.D. and McBean, J.B. (1967) Cancer of the nasopharynx: 142 patients treated in the 11 year period 1950-1960. *Am. J. Roentgenol.*; **99**: 313-325

- Schechter, I. (1971) Mapping of the combining sites of antibodies specific to polyalanine chains. *Ann. N.Y. Acad. Sci.*; **190**: 394-419
- Sculley, T.B., Walker, P.J., Moss, D.J. and Pope, J.H. (1984) Identification of multiple Epstein-Barr virus encoded nuclear antigens with sera from patients with rheumatoid arthritis. *J. Virol.*; **52**: 88-93
- Sculley, T.B., Apolloni, A., Moss, D.J., Mueller-Lantzech, N., Misko, I.S. and Cooper, D.A. (1989) Expression of Epstein-Barr virus nuclear antigens 3, 4 and 6 are altered in cell lines containing B-type virus. *Virology*; **171**: 410-408
- Se Thoe, S.Y. (1991) Antibody reactivity to Epstein-barr virus (EBV) encoded antigens and the detection of EBV DNA in nasopharyngeal and cervical carcinomas. M.Phil. thesis, Institute of Advanced Studies, Univ. Malaya, Kuala Lumpur.
- Shanmugaratnam, K. (1984) Histopathological aspects of nasopharyngeal carcinoma. In: **Cancer of the Liver, Esophagous and Nasopharynx**, (Wagner, G. and Zhang, Y.H. eds.), Springer-Verlag pp152-159.
- Shanmugaratnam, K. and Sobin, L.H. (eds.) (1978) Nasopharyngeal carcinoma. No. 19. In: **Intl. Histological Classification of Tumours**, WHO, Geneva. pp 13-15
- Shanmugaratnam, K., Chan, S.H., de The, G., Goh, J.E.H., Khor, T.H., Simon, M.J. and Tye, C.y. (1979) Histopathology of nasopharyngeal carcinoma: correlation with epidemiology, survival rate and other biological characteristics. *Cancer*; **44**: 1029-1043
- Sheriff, S., Silverton, E.W., Padlaan, E.A. aand Cohen, G.H. (1987) Three dimensional structure of an antigen-antibody complex. *Proc. Natl. Acad. Sci. USA*; **84**: 8075-8079
- Shimizu, N. and Takada, K. (1993) Analysis of the BZLF1 promoter of Epstein-Barr virus: identification of an anti-immunoglobulin response sequence. *J. Virol.*; **67**: 3240-3245
- Sigel, G., Schillinger, M., Henniger, K. and Bauer, G. (1994) IgA directed against early antigen of Epstein-Barr virus is no specific marker for the diagnosis of nasopharyngeal carcinoma. *J. Med. Virol.*; **43**: 222-227

- Sillins, S.L. and Sculley, T.B. (1995) Burkitt's lymphoma cells are resistant to programmed cell death in the presence of the Epstein-Barr virus latent antigen EBNA-4. *Int. J. Cancer*; **60**: 65-72
- Silvestre, D., Kourilsky, F.M., Klein, G., Yata, Y., Neauport-Sautes, C. and Levy, J.P. (1971) Relationship between EBV-associated membrane antigen on Burkitt's lymphoma cells and the viral envelop demonstrated by immunoferritin labelling. *Int. J. Cancer*; **8**: 222-233
- Simons, M.J., Day, N.E., Wee, B.G., Shanmugaratnam, K. and Ho, J.H.C. (1977) Nasopharyngeal carcinoma V. Immunogenetic studies of the Southeast Asian ethnic groups with high and low risk for the tumor. *Cancer Res.*; **34**: 1192-1195
- Simon, M.J., Wee, G.B., Day, N.E., Morris, P.J., Shanmugaratnam, K. and de-The, G. (1974) Immunogenetic aspects of nasopharyngeal carcinoma: I. Differences in HLA-antigen profiles between patients and comparison groups. *Int. J. Cancer*; **13**: 122-134
- Simon, M.J., Wee, G.B., Singh, D. Dharmalingam, S., Yong, N.K., Chai, J.C.W., Ho, J.H.C., Day, N.E. and de-The, G. (1977) Immunogenetic aspects of NPC. V. Conformation of a Chinese-related HLA profile (A2 Singapore 2) associated with an increased risk in Chinese for NPC. *Natl. Cancer Inst. Monograph No. 47*: 147-152
- Simon, M.J. and Shanmugaratnam, L. (1982) The biology of nasopharyngeal carcinoma. In: *UICC Technical Report Series, Vol. 7* (Simon, M.J. and Shanmugaratnam, K. eds.), Intl. Union Against Cancer. pp1-87
- Sinclair, A.J., Palmero, I., Peters, G and Farrel, P.J. (1994) EBNA-2 and EBNA-LP cooperate to cause G0 to G1 transition during immortalization of resting human B lymphocytes by Epstein-Barr virus. *EMBO J.*; **13**: 3321-3328
- Sixbey, J.W., Vesterinen, E.H., Nedrud, J.G., Raab-Traub, N., Walton, L.A. and Pagano, J. S. (1983) Replication of Epstein-Barr virus in human epithelial cells infected *in vitro*. *Nature*; **306**: 480-483
- Sixbey, J.W., Nedrud, J.G., Raab-Traub, N., Hanes, R.A. and Raagand, J.B. (1984) Epstein-Barr virus replication in oropharyngeal epithelial cells. *N. Engl. J. Med.*; **310**: 1225-1230

- Sixbey, J.W., Lemon, S.M. and Pagano, J. S. (1986) A second site for Epstein-Barr virus shedding: the uterine cervix. *Lancet ii*: 1122-1124
- Sixbey, J.W., Shirley, P., Chesney, P.J., Buntin, D.M. and Resnick L. (1989) Detection of a second widespread strain of Epstein-Barr virus. *Lancet ii*: 761-765
- Sixbey, J.W. and Yao, Q.Y. (1992) Immunoglobulin A-induced shift Epstein-Barr virus tropism. *Science*; 255: 1578-1580
- Social Statistics Bulletin (1980) Dept. of Statistics Malaysia, Kuala Lumpur. pp 51-53
- Strnad, B.C., Schuster, T.C., Hopkins, R.F., Neubauer, R.H. and Rabin, H. (1981) Identification of an Epstein-Barr virus nuclear antigen by fluoroimmuno-electrophoresis and radioimmuno-electrophoresis. *J. Virol.*; **38**: 996-1004
- Suhai, S. (1990) Modelling of Protein Structures on the Basis of Sequence Data In: **Modern Methods in Protein- and Nucleic Acid Research Review Articles**, (Tschesche, H. ed.), Walter de Gruyter, Berlin. pp 395-422
- Summers, W.C and Klein, G. (1976) Inhibition of EBV DNA synthesis and late gene expression by phosphonoacetic acid. *J. Virol.*; **18**: 151-155
- Takada, K., Shimizu, N., Sakuma, S. and Ono, Y. (1986) trans activation of the latent Epstein-Barr virus (EBV) genome after transfection of the EBV DNA fragment. *J. Virol.*; **57**: 1016-1022
- Takada, K. and Ono, Y. (1989) Synchronous and sequential activation of latently infected Epstein-Barr virus genome. *J. Virol.*; **63**: 445-449
- Takagi, S., Takada, K. and Sairenji, T. (1991) Formation of intranuclear replication compartments of Epstein-Barr virus with redistribution of BZLF1 and BMRF1 gene products. *Virology*; **185**: 309-315
- Tan, S.H., Chan, S.H., Lee, B.W., Wee, G.B. and Wong, H.B. (1988) HLA association in Singapore children with Grave's disease. *Metabolism*; **37**: 518-519

- Tarone, R.E., Levine, P.H., Yadav, M. and Pandey, J.P. (1990) relationship between immunoglobulin allotypes and susceptibility to nasopharyngeal carcinoma in Malaysia. *Cancer Res.*; **50**: 3186-3188
- Taylor, N., Countryman, J., Rooney, C., Katz, D. and Miller, G. (1989) Expression of the BZLF1 latency-disrupting gene differs in standard and defective Epstein-Barr viruses. *J. Virol.*; **63**: 1721-1728
- Tedeschi R., Foong, Y.T., Cheng, H.M., dePaoli, P. Lehtinen, T., Elfborg, T., and Dillner, J. (1995) The disease-associations of the antibody response against the Epstein-Barr virus transactivator protein ZEBRA can be separated into different epitopes. *J. Gen. Virol.*; **76**: 1393-1400
- Thorley-Lawson, D.A., Schooley, R.T., Bhan, A.K. and Nadler, L.M. (1982) Epstein-Barr virus superinduces a new human B cell differentiation antigen (Blast-1) expressed on transformed lymphocytes. *Cell*; **30**: 415-425
- Thornton, J.M., Edwards, M.S., Taylor, W.R. and Barlow, D.J. (1986) Location of 'continuous' antigenic determinants in the protruding regions of proteins. *EMBO J.*; **5**: 409-15
- Tribbick, G., Edmundser, A.B., Mason, T.J. and Geysen, H.M. (1989) Similar binding properties of peptide ligands for a human immunoglobulins and its light chain dimer. *Molec. Immunol.*; **26**: 625
- Uen, W.C., Luka, J. and Pearson, G.R. (1988) Development of an enzyme-linked immunosorbent assay (ELISA) for detecting IgA antibodies to the Epstein-Barr virus. *Int. J. Cancer*; **41**: 479-482
- Valerio, R.M., Benstead, M., Bray, A.M., Campbell, R.A. and Maeji, N.J. (1991) Synthesis of peptide analogues using the multipin peptide synthesis method. *Anal. Biochem.*; **197**: 168-177
- van Grunsven, W.M.J., Spaan, W.J.M. and Middeldorp, J.M. (1994) Localization and diagnostic application of immunodominant domains of the BKRF3-encoded Epstein-Barr virus capsid protein. *J. Infect. Dis.*; **170**: 13-19
- van Loghem, E. (1984) The immunoglobulin gene: genetic, biological and clinical significance. In: **Clinics in Immunology and Allergy: Immunogenetics** (van Rood, J.J. and de-Vries, R.R.P. eds), WB Saunders, Philadelphia. pp 641-645

- van Oss, C.J., Chaudhury, M.K., and Good, R.J. (1988) Interfacial Lifshitz-van der Waals and polar interactions in macroscopic systems. *Langmuir*; **88**: 927-941
- van Oss, C.J., and Good, R.J. (1991) Surface enthalpy and entropy and the physicochemical nature of hydrophobic and hydrophilic interactions. *J. Dispersion Sci. Technol.*; **12**: 273-287
- van Oss, C.J. (1994a) Nature of specific ligand-receptor bonds, in particular the antigen-antibody bond In: **Immunochemistry** (van Oss & van Regenmortel eds.), Marcel Dekker, New York. pp 581-613.
- van Oss, C.J. (1994b) Intefacial Forces in Aqueous Media. Marcel Dekker, New York
- van Oss, C.J. (1995) Hydrophibic, hydrophilic aand other interactions in epitope-paratope binding. *Molec. Immunol.*; **32(3)**: 199-211
- van Regenmortel, M.H.V. (1987) Antigenic cross-reactivity between proteins and peptides: New insights and applications. *Trends Biochem. Sci.*; **12**: 237-240
- Vroman, B., Luka, J., Rodrwquez, M. and Pearson, G.R. (1985) Characterisation of a major protein with a molecular weight of 160 000 associated with the viral capsid of Epstein-Barr virus. *J. Virol.*; **53**: 107-113
- Wallace, L.E. and Murray, R.J. (1989) Immunological aspects of the Epstein-Barr virus system. *Adv. Viral Oncol.*; **8**: 219-236
- Walter, G., Scheidtmann, K.H., Carbone, A., Laudano, A.P. and Doolittler, R.F. (1980) Antibodies specific for carboxy- and amino-terminal regions of simian virus 40 large tumor antigen. *Proc. Natl. Acad. Sci. USA*; **77**: 5197-5200
- Walter, G. (1986) Production and use of antibodies against synthetic peptides. *J. Immunol. Methods*; **88**: 149-161
- Wang, F., Petti, L., Braun, D., Seung, S. and Kieff, E. (1987) A bicistronic Epstein-Barr virus mRNA encodes two nuclear proteins in latently infected growth transformed lymphocytes. *J. Virol.*; **61**: 945-954

- Wang, F., Gregory, C.D., Sample, C., Rowe, M., Liebowitz, D., Murray, R., Rickinson, A.B. and Kieff, E. (1990) Epstein-Barr virus latent membrane protein (LMP1) and nuclear protein 2 and 3C are effectors of phenotypic changes in B lymphocytes: EBNA 2 and LMP1 cooperatively induce CD23. . *J. Virol.*; **64**: 2309-2318
- Waterhouse, J., Muir, C.S., Correa, P. and Powell, J. (1976) Cancer incidence in five continents Vol III. IARC Sci. Publ. No. 15, Lyon, France. pp 492
- Waterhouse, J., Muir, C.S., Shanmugaratnam, K. and Powell, J. (1982) Cancer incidence in five continents Vol IV. IARC Sci Publ No 42, Lyon, France. pp 294, 384, 388 & 436
- Weiland, L.H. (1978) The histopathological spectrum of nasopharyngeal carcinoma. In: **NPC: Etiology and Control** (de-The, G. and Ito, Y. eds.), IARC Sci. Publ. No. 20, Lyon, France. pp 41-50
- Westhof, E., Altschuh, D., Moras, D., Bloomer, A.C., Mondragon, A., Klug, A. and Van Rogenmortel, M.H.V. (1984) Correlation between segmental mobility and the location of antigenic determinants in proteins. *Nature*; **311**: 123-26
- Wolf, H., Hans, M. and Wilmes, E. (1984) Persistent of Epstein-Barr virus in the parotid gland. *J. Virol.* **51**: 795-798
- Yadav, M. (1987) Immunological, genetical and epidemiological studies of nasopharyngeal carcinoma. *Tropical Biomedicine*; **4**: 202-221
- Yadav, M., Tan, M.K., Singh, P. and Dharmalingam, S.K. (1984) Nasopharyngeal carcinoma in Malaysians under the age of 20 years. *Clin. Oncol.*; **10**: 353-361
- Yadav, M.S., Tan, M.K. and Singh, P. (1985) Nasopharyngeal carcinoma in Malaysia: Distribution by race for period 1981-1983. *J. Mal. Soc. Health*; **5**: 67-71
- Yadav, M.S., Malliga, N. and Ablashi, D.V. (1987) Development of immunity to Epstein-Barr virus in Malaysian children. *Microbiologica*; **10**: 29-35
- Yao, Q.Y., Richinson, A.B. and Epstein, M.A. (1985) A re-examination of the Epstein-Barr virus cancer state in healthy sero-positive individuals. *Int. J. Cancer*; **35**: 35-42

- Yates, J., Warren, N. and Sugden, D. (1985) Stable replication of plasmids derived from Epstein-Barr virus in various mammalian cells. *Nature*; **313**: 812-815
- Yeo, P.P.B., Chan, S.H., Thai, A.C., Ng, W.Y., Lui, K.F., Wee, G.B., Tan, S.H., Lee, B.W., Wong, H.B. and Cheah, J.S. (1989) HLA BW46 and DR9 associations in Graves' disease of Chinese patients are age- and sex-related. *Tiss. Antigens*; **34**: 179-184
- Young, L.S., Dawson, C.W., Clark, D., Rupani, H., Busson, P., Tursz, T., Johnson, A. and Rickinson, A.B. (1988) Epstein-Barr virus gene expression in nasopharyngeal carcinoma. *J. Gen Virol.*; **69**: 1051-1065
- Yu, M.C., Ho, J.H.C., Ross, R.K. and Henderson, B.E. (1981) Nasopharyngeal carcinoma in Chinese - salted fish or inhaled smoke? *Prev. Med.*; **10**: 15-24
- Yu, M.C., Ho, J.H.C., Henderson, B.E. and Armstrong, R.W. (1985) Epidemiology of nasopharyngeal carcinoma in Malaysia and Hong Kong. *Natl. Cancer Inst. Monogr.*; **69**: 203-207
- Yu, M.C., Ho, J.H.C., Lai, S.H. and Henderson, B.E. (1986) Cantonese-style salted fish as a cause of nasopharyngeal carcinoma: report of a case-control study in Hong Kong. *Cancer Res.*; **46**: 956-961
- Yu, M.C., Mo, C.C., Chong, W.X., Yeh, F.S. and Henderson, B.E. (1988) Preserved foods and nasopharyngeal carcinoma: a case-control study in Guangxi, China. *Cancer Res.*; **48**: 1954-59
- Yu, M.C., Garrabrant, D.H., Teng-Bo, H. and Henderson, B.E. (1990) Occupational and other non-dietary risk factors for nasopharyngeal carcinoma in Guangzhou, China. *Int. J. Cancer*; **45**: 1033-039
- Zeng, J., Gong, C.H., Jan, M.G., Fun, Z., Zhang, L.G. and Li, H.Y. (1983a) Detection of Epstein-Barr virus IgA/EA antibody for diagnosis of nasopharyngeal carcinoma by immunoradiography. *Int. J. Cancer*; **31**: 599-601
- Zeng, Y., Zhang, J.M., Li, L.Y., Wang, P.Z., Tang, H., Ma, Y.R., Zhu, Y.S., Pan, W.J., Liu, Y.X., Wei, J.N., Chen, J.Y., Mo, Y.K., Li, E.J. and Tan, B.F. (1983b) Follow-up studies on Epstein-Barr virus IgA/VCA antibody positive persons in Zangwu County, China. *Intervirol.*; **20**: 190-194

- Zeng, Y. (1985) Seroepidemiological studies on nasopharyngeal carcinoma in China. *Adv. Cancer Res.*; **44**: 121-138
- Zhang, Q., Gutsh, D. and Kenney, S. (1994) Functional and physical interaction between p53 and BZLF1: implications for Epstein-Barr virus latency. *Mol. Cell. Biol.*; **14**: 1929-1938
- Zimber, U., Adldinger, H.K., Lenoir, G.M., Vuillaume, M., Knebel-Doeberitz, M.V., Laux, G., Desgranges, C., Wittman, P., Freese, U.K., Schneider, U. and Bornkamm, G. (1986) Geographical prevalence of two types of Epstein-Barr virus. *Virology*; **154**: 56-66
- Zippin, C., Tekawa, I.S., Bragg, K.U., Watson, D.A. and Linden, G. (1962) Studies on heredity and environment in cancer of the nasopharynx. *J. Natl. Cancer Inst.*; **29**: 483-490
- zur Hausen, H., Schulte-Holthausen, H., Klein, G., Henle, W., Henle, G., Clifford, P. and Santesson, L. (1970) EBV DNA in biopsies of Burkitt's tumour and anaplastic carcinomas of the nasopharynx. *Nature*; **228**: 1056-1058
- zur Hausen, H., O'Neil, F.J. and Freese, V.K. (1978) Persisting oncogenic herpesvirus induced by the tumour promoter TPA. *Nature (London)*; **272**: 373-375

### **Additional References**

Mallen, R.W. and Sandro, W.G. (1974) Nasopharyngeal carcinoma in Eskimos. *Can. J. Otolaryngol.*; **3**: 157-179

Neilsen, N.H., Mikkelsen, F. and Hansen, J.P.H. (1977) Nasopharyngeal cancer in Greenland: the incidence in an Arctic Eskimo population. *Acta Pathol. Microbial. Scand.*; **85A**: 850-858

Prasad, U. and Rampal, L. (1992) Descriptive epidemiology of nasopharyngeal carcinoma in Peninsular Malaysia. *Cancer Causes and Control*; **3**: 179-182