

REFERENCES

Aminabhavi T.M, Balundgi R.H and Cassidy P.E. 1990 . A review on biodegradable plastics. Polymer Plastic Technology and Engineering, 29: 235-262

Annual Book of ASTM Standards D5209-91, Vol 8.03 .1992. Standard test method for determining the aerobic biodegradation of plastic materials in the presence of municipal sewage sludge.

Annual Book of ASTM Standards D5247-92, Vol 8.03. 1992. Standard test method for determining the aerobic biodegradability of degradable plastics by specific microorganisms.

Annual Book of ASTM Standards E252-84. 1992. Standard test method for thickness of thin foil and film by weighing.

Augusta J, Muller R.J, Widdecke H. 1993. A rapid evaluation plate-test for the biodegradability of plastics. Appl Microbiol Biotechnol, 39: 673 – 678

Baere L.D, Wilde B.D and Tillinger R. 1994. Standard test methods for polymer biodegradation in solid waste treatment systems. In: Doi Y. and Fukuda K. (Eds.). Biodegradable Plastics and Polymers. Elsevier Science: 217-227.

Baharuddin N.A. 17th November 1996. Plan to reduce use of plastic items. News Straits Times , pg :8.

Brock T.D and Brock K.M. 1973. Basic Microbiology with Applications. Prentice- Hall, Inc, Englewood Cliffs, New Jersey.

Bonthrone K.M, Clauss J, Horowitz D.M, Hunter B.K, Sanders J.K.M. 1992. The biological and physical chemistry of polyhydroxyalkanoates as seen by NMR spectroscopy. FEMS Microbiology Reviews, 103: 267-278

Brandl H and Puchner P. 1990. The degradation of shampoo bottles in a lake. In E.A.Dawes (Eds.). Novel Biodegradable Microbial Polymers. Kluwer Academic Publishers: 421-422

Braunegg C, Sonnleitner B, Lafferty R.M. 1978. A rapid gas chromatographic method for the determination of poly-β-hydroxybutyrate acid in microbiol biomass. Appl Microbiol Biotechnol, 6: 29-37

Briese B.H, Jendrossek D and Schlagel H.G. 1994. Degradation of poly(3-hydroxybutyrate-co-3-hydroxyvalerate) by aerobic sewage sludge. FEMS Microbiology Letters , 117:107-112

Cho K.S, Ryu H.W, Park C.H and Goodrich P.R. 1997.
Poly(hydroxybutyrate-co-hydroxyvalerate) from swine waste liquor by
Azobacter vinelandii UWD. Biotechnology Letters ,9(1): 7-10

Doi Y.1990. Biodegradation of microbial polyesters. In: Microbial Polyesters Chapter 8:135-153

Doi Y.1995.Microbial synthesis, physical properties and biodegradability of polyhydroxyalkanoates. Macromol Symp 98: 585-599

Doi Y, Kawaguchi Y, Nakamura Y, Kunioko M. 1989. Nuclear magnetic resonance studies of poly(3-hydroxybutyrate) and polyphosphate metabolism in *Alcaligenes eutrophus*. Appl Env Microbiol, 55: 2932-2938

Doi Y, Kanesawa Y, Kunioko M, Saito T. 1990a. Biodegradation of microbial copolyesters: poly(3-hydroxybutyrate-co-3-hydroxyvalerate) and poly(3-hydroxybutyrate-co-4-hydroxybutyrate). Macromolecules, 23:26-31

Doi Y, Mukai K, Kasuya K, Yamada K.1994. Biodegradation of biosynthetic and chemosynthetic polyhydroxyalkanoates. In :Doi Y. and Fukuda K.(Eds.)Biodegradable Plastics and Polymers. Elsevier Science :39-51

Doi Y, Kasuya K, Abe H, Koyama N, Ishiwatari S, Takagi K, Yoshida Y. 1996. Evaluation of biodegradabilities of biosynthetic and chemosynthetic polyesters in river water. Polymer Degradation and Stability, 51: 281-286

Donelly J.1990. Degradable Plastics. Garbage, 2:42-47

Eya H, Iwaki N, Otsuji Y. 1994. A study on methods of testing biodegradable plastics in soil: aerobic and anaerobic gas evolution test; petri dish test in various soils. In:Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science: 337-344

Foster L.J.R, Zervas S.J, Lenz R.W, Fuller R.C.1995. The biodegradation of poly-3-hydroxyalkanoates, PHAs with long alkyl substituents by *Pseudomonas maculicola*. Biodegradation,6:67-73

Furniss B.S, Hannaford A.J, Smith P.W.G and Tatchell A.R.(revisers). 1989. Vogel's Textbook of Practical Organic Chemistry (5th ed). Longman, London.

Griffin G.J.L. 1994. Chemistry and Technology of Biodegradable Polymers. Blackie Academic and Professional. Chapman and Hall.

Hammond T, Liggat J.J. 1994. Properties and applications of bacterially derived polyhydroxyalkanoates. In:Gerald Scott and Dan Gilead.(Eds.) Degradable Polymers. Chapter 5:89-109

Hobart H.W, Lynne L.M.Jr, John A.D, Frank A.S.Jr. 1981. Infrared Spectrophotometry. In: Instrumental Methods of Analysis 6th edition. Wadsworth Publishing company, Belmont, California.

Hori K, Soga K, Doi Y. 1994. Production of poly(3-hydroxyalkanoates-co-3-hydroxy-ω-fluoroalkanoates) by *Pseudomonas oleovorans* from 1-fluorononane and gluconate. Biotechnology Letters, 16(5):501-506

Ichikawa M, Nakamura K, Yoshie N, Asakawa N, Inoue Y. 1996. Morphological study of bacterial poly (3-hydroxybutyrate-co-3-hydroxypropionate). Macromol Chem Phys, 197:2467-2480

International Standard ISO 15986 (draft). 1998. Plastics Evaluation of compostability: Test scheme for final acceptance.

Iwata T, Doi Y, Kasuya K, Inoue Y. 1997. Visualization of enzymatic degradation of poly[(R)-3-hydroxybutyrate] single crystals by an extracellular PHB depolymerase. Macromolecules, 30: 833-839

Jendrossek D, Schirmer A, Schlegel H.G.1996. Biodegradation of polyhydroxyalkanoic acids. Appl Microbiol Biotechnol, 46: 451-463

Kanesawa Y, Tanahashi N, Doi Y, Saito T. 1994. Enzymatic degradation of microbial poly(3-hydroxyalkanoates). Polymer Degradation and Stability, 45:179-185

Kato M, Bao H.J, Kang C.K, Fukui T, Doi Y. 1996. Production of novel copolyesters of 3-hydroxybutyric acid and medium chain length 3-hydroxyalkanoic acids by *Pseudomonas* sp.61-3 from sugars. Appl Microbiol Biotechnol, 45: 363-370

Kawai F. 1994. Biodegradation of polyethers and polyacrylate. In:Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science: 24-38

Kim B.S, Chang H.N. 1998. Production of poly(3-hydroxybutyrate) from starch by Azobacter chroococcum. Biotechnology Letters, 20(2):109-112

Kimura M, Toyota K, Iwatsuki M, Sawade H. 1994. Effects of soil conditions on biodegradation of plastics and responsible microorganisms. In: Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science: 92-105

Kitamura S, Doi Y. 1994. Staining method of poly(3-hydroxyalkanoic acids) producing bacteria by nile blue. Biotechnology Techniques, 8(5):345-350

Kitano M, Yakabe Y. 1994. Strategy for biodegradability testing in OECD. In: Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science: 217-227

Koyama N, Doi Y. 1997. Effect of solid state structures on the enzymatic degradability of bacterial poly(hydroxyalkanoic acids). Macromolecules, 30:826-832

Kunioka M, Kawaguchi Y and Doi Y. 1989. Production of biodegradable copolyesters of 3-hydroxybutyrate and 4-hydroxybutyrate by *Alcaligenes eutrophus*. Appl Microbiol Biotechnol, 30:569-573

Lageveen R.G, Huisman G.W, Preusting H, Ketelaar P, Eggink G and Witholt B. 1988. Formation of polyesters by *Pseudomonas oleovorans*: Effect of substrates on formation and composition of poly-(R)-3-hydroxyalkanoates and poly-(R)-3-hydroxyalkanoates. Appl Environ Microbial. 54:2924-2932.

Matavulj M and Molitoris H.P. 1992. Fungal degradation of polyhydroxyalkanoates and a semiquantitative assay for screening their degradation by terrestrial fungi. FEMS Microbiology Reviews, 103:323-332

Mergaert J, Anderson C, Wouters A, Swings J and Kersters K. 1992. Biodegradation of polyhydroxyalkanoates. FEMS Microbiology Reviews, 103:317-322

Molitoris H.P, Moss S.T, de Koning G.J.M, Jendrossek D. 1996. Scanning electron microscopy of polyhydroxyalkanoate degradation by bacteria. Appl Microbiol Biotechnol, 46:570-579

Mukai K. and Doi Y. 1993. Enzymatic degradation of poly (3-hydroxybutyrate) by marine organisms. In: Proceedings of the International Symposium on Bacterial Polyhydroxyalkanoates. Schlegel H.G and Steinbuchel A. (Eds.) Goltze Druck, Gottingen, FRG:457-458.

Mukai K, Yamada K, Doi Y. 1994. Efficient hydrolysis of polyhydroxyalkanoates by *Pseudomonas stutzeri* YM1414 isolated from lake water. Polymer Degradation and Stability, 43:319-327

Muller R.J, Augusta J, Walter T, Widdecke H. 1994. The development and modification of some special test method and the progress in standardisation of test methods in Germany.In: Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science:237-249

Nakayama K., Saito T., Fukui T.,Shirakura Y., Tomita K. 1985.

Purification and properties of extracellular poly(3-hydroxybutyrate)depolymerases from *Pseudomonas lemoigne*.

Biochem.Biophys.Acta, 827:63-72

Narayan R. 1994. Impact of governmental policies, regulations and standards activities on an emerging biodegradable plastics industry. In: Doi Y and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science:261-272

Narayan R, Barenberg S.A, Brash J.L and Redpath A.E. 1990. Degradable materials: Perspectives, issues and opportunities. CRC Press, Fl.

Nishioka M, Tuzuki T, Wanajyo Y, Oonami H and Horiuchi T. 1994. Biodegradation of Bionolle. In: Doi Y and Fukuda K.(Eds.)Biodegradable Plastics and Polymers. Elsevier Science :584-590

OECD.1981. Guideline for testing of chemicals. Nos 301, A to E.

Pagga U. 1998. Biodegradability and compostability of polymeric materials in the context of the European packaging regulation. Polymer Degrad Stab, 59:371-376 **Richard P.W.** 1994. Perspectives on standards test methods for biodegradable plastics. In: Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science: 250-258

Saito Y. and Doi Y. 1994. Microbial synthesis and properties of poly(3-hydroxybutyrate-co-4-hydroxybutyrate) in *Comamonas acidovorans*. Int. J. Biol. Macromol, 16(2):99-104

Sawada H. 1994. Field testing of biodegradable plastics. In: Doi Y. and Fukuda K. (Eds.) Biodegradable Plastics and Polymers. Elsevier Science :298-312

Scott G. 1994. Environmental biodegradation of hydrocarbon polymers: Initiation and control. In: Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science:79-91

Seal K.J. 1994. Test methods and standards for biodegradable plastics. In: Griffin G.J.L.(Eds.) Chemistry and Technology of Biodegradable Polymers. Blackie Academic and Professional:117-134

Seal K.J. and Pantke M. 1986. An interlaboratory investigation into the biodeterioration of plastics with specific reference to polyurethane. Part 1: Petri dish test. Material und Organismen, 21(2): 151-164

Shimamura E. and Kasuya K, Kobayashi G, Shiotani T, Shima Y and Doi Y. 1994. Physical properties and biodegradability of microbial poly (3-hydroxybutyrate-co-3-hydroxyhexanoate). Macromolecules, 24: 878-880.

Shiraki M, Shimada T, Tatsumichi M, Saito T.1985. Purification and characterization of extracellular poly (3-hydroxybutyrate) depolymerases. J. Environ. Polym.Degrad, 3:13-21

Starnecker A. and Menner M. 1996. Assessment of biodegradability of plastics under simulated composting conditions in a laboratory test system. International Biodeterioration and Biodegradation: 85-92

Swift G. 1992. Biodegradability of polymers in the environment: complexities and significance of definitions and measurements. FEMS Microbiology Reviews ,103: 339-346

Swift G. 1994. Expectations for biodegradation testing methods. In: Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science:228-236

Tan I.K.P,Sudesh Kumar K, Theanmalar M,Gan S.N, Gordon III B. 1997. Saponified palm kernel oil and its major free fatty acids as carbon substrates for the production of polyhydroxyalkanoates in *Pseudomonas putida* PGA1. Appl Microbiol Biotechnol, 47: 207-211

Tanio T, Fukai T,Shirakura Y, Saito T, Tomita K, Kaino T, Masamune S.1982. An extracellular poly(3-hydroxybutyrate) depolymerase from Alcaligenes faecalis. Eur. J.Biochem, 124: 71-77

Theanmalar M. 1996. Studies of some cultivation conditions on polyhydroxyalkanoates (PHA) accumulation in *Pseudomonas putida* grown with palm kernel oil derivatives. M.Biotech thesis.

Tsuji M. and Omoda Y. 1994. A simple method for detection of polymeric degrading microorganisms on agar plates. In: Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science: 345-350

Uefuji M, Kasuya K, Doi Y. 1997. Enzymatic degradation of poly[(R)-3-hydroxybutyrate]:Secretions and properties of PHB depolymerase from *Pseudomonas stutzeri*. Polymer Degradation and Stability, 58:275-281

Urstadt S, Augusta J, Muller R.J, Deckwer W.D. 1995. Calculation of carbon balances for evaluation of the biodegradability of polymers. Journal of Environmental Polymer Degradation, 3(3):121-131

USEPA(U.S. Environmental Protection Agency). 1990. Characterization of municipal solid waste in the United States.Update. USEPA, Washington, D.C.

Vogel H.J. and Bonner D.M. 1956. Acetylornithunase of *Escherica coli*: partial purification and some properties. J. Biol. Chem, 218: 97-106

Witholt B, Huisman G.W and Preusting. 1990. Bacterial Poly(3-hydroxyalkanoates). In: E.A.Dawes.(Eds.) Novel Biodegradable Microbial Polymers. Kluwer Academic Publishers:161-173

Witt U, Muller R.J, Deckwer W.D. 1996. Studies on sequence distribution of aliphatic/aromatic copolyesters by high-resolution ¹³C nuclear magnetic resonance spectroscopy for evaluation of biodegradability. Macromol. Chem. Phys, 197: 1525-1535

Yabannavar A, Bartha R.1993. Biodegradability of some food packaging materials in soil. Soil. Biol. Biochem, 25(11): 1469-1475

Yabannavar A, Bartha R. 1994. Methods for assessment of biodegradability of plastic films in soil. Applied and Environmental Microbiology, 50(10):3608-3614

Yakabe Y. and Kitano M. 1994. Evaluation of biodegradability of biodegradable plastics in activated sludge. In: Doi Y. and Fukuda K.(Eds.)Biodegradable Plastics and Polymers. Elsevier Science: 331-336

Yokota Y, Ishioka R, Moteki Y, Watanabe N. 1994. Physical properties and processabilities of biodegradable plastics "Bionolle".In: Doi Y. and Fukuda K.(Eds.) Biodegradable Plastics and Polymers. Elsevier Science: 577-583