

DISTRIBUTION OF PETROGENIC
HYDROCARBONS IN WATER, SEDIMENT AND
COCKLES (*Anadara granosa*) FROM KUALA SUNGAI
SEPETANG AND KUALA SUNGAI SELANGOR

MOHD HILMI BIN JAAFAR

SUBMITTED AS PARTIAL FULFILLMENT FOR THE
DEGREE OF MASTER OF TECHNOLOGY
(ENVIRONMENTAL MANAGEMENT)

INSTITUTE OF POSTGRADUATE STUDIES AND
RESEARCH
UNIVERSITY OF MALAYA
2000

Perpustakaan Universiti Malaya



A510570859

OK

ABSTRACT

In order to assess the level of oil pollution in the cockle culture areas along the Straits of Malacca, water, sediment and cockle samples were collected from the culture beds of Kuala Sepetang and Kuala Selangor, two of the largest cockle producing areas of Malaysia. The samples were analysed for petrogenic hydrocarbons by employing a spectrofluorometric technique which measures the content of polycyclic aromatic hydrocarbons (PAH) in oil. The content of aliphatic straight chain hydrocarbons were measured by using gas chromatography-mass spectrometry (GCMS) which also allowed for the estimation of carbon preference indices and hence providing a means of estimating biogenic /petrogenic inputs.

Petrogenic hydrocarbons were observed in all the environmental samples collected from the two sites as evidenced by the spectrofluorometric measurements. The level of hydrocarbons in water at both sites ranged from 30.82 $\mu\text{g/L}$ to 92.59 $\mu\text{g/L}$ Seligi crude oil equivalents (1.195-3.075 chrysene equivalents). The highest level of hydrocarbons in Kuala Sepetang was recorded at Station 2: 92.59 $\mu\text{g/L}$ and 3.075 $\mu\text{g/L}$ expressed as Seligi crude oil and chrysene equivalents respectively. The levels of petrogenic hydrocarbons in surficial sediments were higher than those in the water column by three orders of magnitude, ranging from 2.42-45.50 $\mu\text{g/g}$ Seligi crude oil equivalents (0.092-1.614 $\mu\text{g/g}$ chrysene equivalents). Relatively higher levels of petrogenic hydrocarbons were found in Kuala Sepetang (average: 30.37 $\mu\text{g/g}$ Seligi crude oil equivalents) compared to Kuala Selangor (average: 12.49 $\mu\text{g/g}$ Seligi crude oil equivalents). Concentrations of petrogenic hydrocarbons in cockle tissues were noted to be in the same order of magnitude to those found in the sediments. Petrogenic hydrocarbons in cockle tissues ranged from 4.427 to 38.48 $\mu\text{g/g}$ Seligi crude oil equivalent (0.176-1.514 $\mu\text{g/g}$ chrysene equivalent) for the two sites. However, unlike the levels in the sediments there were no significant differences between the average concentration of petrogenic hydrocarbons in Kuala Sepetang (average: 17.75 $\mu\text{g/g}$ Seligi crude oil equivalent) and that found in Kuala Selangor (16.45 $\mu\text{g/g}$ Seligi crude oil equivalent).

There does not appear to be a correlation between hydrocarbon levels in sediments and those in the cockles. There was also an absence of correlation between hydrocarbon levels in tissues and those in the water column.

GCMS analysis of the extracts from water, sediments and cockles revealed the presence of hydrocarbons as low as C₁₄ up to C₃₀ based on available standards. Carbon preference indices (CPI) were also estimated using higher ranges of n-alkanes, as they are less susceptible to evaporation. For water samples from both sites, the CPIs were found to be close to unity, indicative of petrogenic hydrocarbon pollution. Extracts from sediment samples had CPI values exceeding 1, in many cases substantially so. This observation was attributed to the nature of the mud being highly biological and productive in nature characteristic of the mudflat ecosystem. CPI values in cockle tissues were also found to exceed 1 indicating the predominance of biogenic hydrocarbons in the tissues.

The present study also examined the uptake and depuration of oil by the cockles. In a laboratory-controlled experiment cockles were exposed to water-soluble fractions (WSF) of a crude oil and refined oil in a semi-static manner whereby the water was renewed every 24hrs. Analysis of aliquots of the WSFs gave initial concentrations of 1.32 µg/L and 1.71 µg/L for the Seligi crude oil and the refined oil respectively. The uptake of aromatic hydrocarbons as measured by fluorescence spectroscopy was clearly observed from the 1st day of exposure when compared to control animals. Cockles appear to take up greater amounts of crude oil compared to refined oil even though the initial concentration of the crude oil was lower than the refined oil. At the end of the 7th day exposure period the concentrations of petrogenic hydrocarbons in the cockles were 48 ng/g and 26 ng/g Seligi crude oil equivalents from crude oil and refined oil respectively. When removed into clean water, depuration was observed to take place. At the end of 12 days in clean water the levels of aromatic hydrocarbons in the tissues were still significantly higher than those of control animals. Crude oil-exposed animals had 38 ng/g Seligi crude oil equivalents of petrogenic hydrocarbons (78 % of the concentration following 1-day exposure) while refined oil-exposed animals had 18 ng/g Seligi crude oil equivalents (78 % of the concentration following 1-day exposure).

ACKNOWLEDGEMENTS

I would like to dedicate this work especially to my parents who have always been on my side during my ups and downs with their full support and encouragement throughout this course. They transmitted their courage to me to keep on going to complete this project. To them I owe a debt of gratitude that I can never entirely repay in my whole life.

My very special thanks to Associate Professor Dr. Abdul Rani Abdullah, whose patience is legendary in supervising, guiding and helping me in the completion of this project thesis. I am indebted to his enthusiasm, generosity and readiness to lay his helping hand even within the strictness of his hectic schedule. I wish him god bless and happiness.

I also wish to thank Miss Wong Kang Ying and Mr. Siew Yau Foo for their kindheartedness to help me in my laboratory work. Without their help, this thesis might not become a reality. Once again, I am indebted to many people in IPSP for their help and contribution. I particularly want to thank Mr. Zulkifli Abdullah, Mr. Mustafa Tahar, and Miss Zubaidah Yusop for their assistance. Unforgettable also are all IPSP librarian; Ms Molly Chuah, Ms Rafiati Shafie, Miss Zaini Abdul Wahab and Ms Rokiah Jaafar, who have always been kind and helpful to me. There are also many people who helped me during the course of this project which their names I do not know, who have given a substantial contribution. To all of them, I give my deepest appreciation.

Finally, I thank all my friends and my colleagues for their support and encouragement, especially Miss Marni Baharum, Ms Saraswathy and Miss Noor Zalida who have brought much joy during my lab work.

This little work is far from jewel. Neither it is a masterpiece. But, I hope it would provide some help and contribution to the development of knowledge, wisdom and virtue. Let's hold each other hands and together we strive for the betterment of tomorrow. May God save the Planet Earth. He sure will, only if we have the goodwill. By God willing.

CONTENTS

	Page
TITLE	i
ABSTRACT	ii
ACKNOWLEDGEMENTS	iv
CONTENTS	v
LIST OF TABLES	viii
LIST OF FIGURES	x
CHAPTER I: INTRODUCTION	1
1.1 Hydrocarbons in the Marine Environment	1
1.2 Crude Oils	2
1.3 Biogenic Hydrocarbons	3
1.4 Hydrocarbons in the Water Column	6
1.5 Effects of Weathering on Oil in the Marine Environment	8
1.6 Sources of Hydrocarbons in the Marine Environment	12
1.6.1 Natural Sources	13
1.6.2 Anthropogenic Sources	14
1.6.2.1 Oil exploration and production	15
1.6.2.2 Oil tanker and vessel operations	15
1.6.2.3 Oil spill accidents	16
1.6.2.4 Oil refinery discharges	17
1.6.2.5 Land-based Sources	18
1.7 Fate of Hydrocarbons in the Marine Environment	19
1.7.1 Evaporation	20
1.7.2 Dissolution	21
1.7.3 Emulsification	22
1.7.4 Adsorption and Sedimentation	23
1.7.5 Oxidation	24

1.7.6	Microbial Degradation	24
1.7.7	Entry into Marine Biota	26
1.8	Effects of Hydrocarbon Pollution	27
1.8.1	Effects on Marine Fauna	28
1.8.2	Effects on Benthic Invertebrate	30
1.8.3	Effects on Marine Plants	31
1.8.3.1	Mangroves	31
1.8.3.2	Marine algae	34
1.8.3.3	Seagrasses	35
1.8.3.4	Phytoplankton	36
1.8.4	Effects on Humans	37
 CHAPTER II: OIL POLLUTION IN MALAYSIA		 40
2.1	General	40
2.2	Sources of Oil Pollution	43
2.2.1	Sea-based Sources	43
2.2.2	Land-based Sources	47
2.3	Impacts of Hydrocarbons Pollution in Malaysia	48
 CHAPTER III: EXPERIMENTAL		 53
3.1	Introduction	53
3.2	Chemicals and Glassware	55
3.3	Sample Collection and Storage	56
3.4	Cleaning of Glassware and Other Materials	59
3.5	Sample Treatment and Extraction	60
3.6	Fractionation	61
3.7	Analysis	62
3.8	Uptake and Depuration of Petroleum Hydrocarbons in Cockles	63

CHAPTER IV: RESULTS AND DISCUSSION	66
4.1 Precision and Accuracy	66
4.2 Oil Pollution in the Mudflats of Kuala Sepetang and Kuala Selangor	69
4.2.1 Gas Chromatographic Analysis	76
4.2.2 Biogenic versus petrogenic hydrocarbons	85
4.3 Uptake and Depuration of Petrogenic Hydrocarbons by <i>Anadara granosa</i>	87
CHAPTER V: CONCLUSIONS	92
APPENDICES	94
Appendix A Representative Calibration Curves for Petrogenic Hydrocarbon Analysis Using UV-Fluorescence	94
Appendix B Results of Measurements of Petrogenic Hydrocarbon Using UV-Fluorescence	98
REFERENCES	101

LIST OF TABLES

Table		Page
1.1	Biogenic hydrocarbons and their origins	6
1.2	Representative room temperature solubility of hydrocarbons in distilled water	7
1.3	Estimates of annual petroleum hydrocarbons inputs into the world oceans	12
1.4	Tanker accidents resulting in oil loss (1989 – 1973)	17
1.5	Amounts of petroleum hydrocarbons in bivalve from contaminated areas	31
1.6	Effects of oil spill on mangroves	33
2.1	Forest cover in the Malacca Straits coastal zone	41
2.2	Types of shipping casualties in the Straits of Malacca, 1977 - 1993	45
2.3	Oil spill incidents in the Straits of Malacca from 1976 to 1997	45
3.1	Solvents and chemicals used in the present study	56
3.2	Eluents used for fractionation	62
4.1	Recovery and precision of the adopted methodology for Seligi crude oil added to cockle tissue as determined by fluorescence	68
4.2	Recovery and precision of the adopted methodology for Seligi crude oil added to artificial seawater as determined by fluorescence	68
4.3	Percentage recovery of internal standards	68
4.4	Petroleum hydrocarbons in estuarine water	70
4.5	Petroleum hydrocarbons in surficial sediment	71
4.6	Petroleum hydrocarbons in cockle tissue	72
4.7	Concentration of selective aliphatic hydrocarbons identified from water samples	82
4.8	Concentration of selective aliphatic hydrocarbons identified from sediment samples	83
4.9	Concentration of selective aliphatic hydrocarbons identified from tissue samples	84
4.10	CPI values for water sample extracts	85

Table	Page
4.11 CPI values for sediment sample extracts	86
4.12 CPI values for tissue sample extracts	86
4.13 Uptake and depuration of aromatic hydrocarbons by <i>Anadara granosa</i> exposed to Seligi crude oil (1.32 µg/L) and refined oil (1.71 µg/L)	88
B-1 Results of petrogenic hydrocarbon measurements for water samples using UV-Fluorescence	98
B-2 Results of petrogenic hydrocarbon measurements for sediment samples using UV-Fluorescence	99
B-3 Results of petrogenic hydrocarbon measurements for tissue samples using UV-Fluorescence	100

LIST OF FIGURES

Figure		Page
3.1	Sampling stations at Kuala Selangor	57
3.2	Sampling stations at Kuala Sepetang	58
4.1	Petroleum hydrocarbons in seawater for both sites	70
4.2	Petroleum hydrocarbons in sediment and tissue for Kuala Sepetang	73
4.3	Chromatogram of the aliphatic hydrocarbon standards	78
4.4	Representative chromatogram for the water samples	79
4.5	Representative chromatogram for the sediment samples	80
4.6	Representative chromatogram for the tissue samples	81
4.7	Uptake and depuration of aromatic hydrocarbons in <i>Anadara granosa</i> exposed to Seligi crude oil (1.32 µg/L) and marine engine oil (1.71 µg/L)	89
4.8	Gas chromatogram of cockle tissue extracts from test tank (crude oil exposed, 1.32 µg/L, 4 days)	90
4.9	Gas chromatogram of tissue extracts from control tank	91
A-1	Representative calibration curve for water samples using UVF analysis with Seligi crude oil as standard	94
A-2	Representative calibration curve for sediment samples using UVF analysis with Seligi crude oil as standard	95
A-3	Representative calibration curve for tissue samples using UVF analysis with Seligi crude oil as standard	96
A-4	Representative calibration curve using UVF analysis with chrysene as standard	97