

---

## CONTENTS

|  | <b>Page</b> |
|--|-------------|
| ACKNOWLEDGEMENT                                | i           |
| ABSTRACT                                       | ii          |
| CONTENTS                                       | iii         |
| LIST OF FIGURES                                | vi          |
| LIST OF PLATES                                 | viii        |
| LIST OF TABLES                                 | ix          |
| LIST OF ABBREVIATIONS                          | x           |
| <br>   |             |
| CHAPTER ONE                                    | 1 - 6       |
| INTRODUCTION                                   |             |
| <br>   |             |
| CHAPTER TWO                                    | 7 - 44      |
| LITERATURE REVIEW                              |             |
| <br>   |             |
| LIGNOCELLULOSE                                 | 7           |
| Lignin   |             |
| Cellulose                                      |             |
| Hemicellulose                                  |             |
| Lignocellulosic utilization                    |             |
| <br>   |             |
| MICROBIAL UTILIZATION OF LIGNOCELLULOSICS      | 9           |
| <i>Pycnoporus sanguineus</i>                   |             |
| <br>   |             |
| LIGNOCELLULOLYTIC ENZYMES                      | 19          |
| Lignin-degrading enzymes                       |             |
| Cellulose- degrading enzymes                   |             |
| Xylan- degrading enzymes                       |             |
| <br>   |             |
| SOME APPLICATIONS OF LIGNOCELLULOLYTIC SYSTEMS | 30          |
| <br>   |             |
| SOLID-STATE FERMENTATION                       | 32          |
| <br>   |             |
| SOLID-STATE FERMENTATION FOR ENZYME PRODUCTION | 41          |

---

|   |         |
|---|---------|
| <b>CHAPTER THREE</b>  | 45 - 65 |
| <b>ENZYME ACTIVITIES DURING LIGNOCELLULOSIC DEGRADATION</b>   |         |
| INTRODUCTION  | 45      |
| MATERIALS AND METHODS   | 46      |
| Substrate   |         |
| Inoculum  |         |
| Solid- state fermentation   |         |
| Extraction of crude extracellular enzymes   |         |
| Analysis  |         |
| RESULTS AND DISCUSSION  | 53      |
| pH variation  |         |
| Soluble protein   |         |
| Xylanase activity   |         |
| Lignin peroxidase activity  |         |
| Laccase activity  |         |
| CONCLUSION  | 65      |
| <b>CHAPTER FOUR</b>   | 66-89   |
| <b>EFFECT OF (a) NITROGEN CONCENTRATION AND (b) INOCULUM AGE AND DENSITY (W/W) ON LACCASE PRODUCTIVITY DURING SSF OF 'HAMPAS'</b> |         |
| INTRODUCTION  | 66      |
| MATERIALS AND METHODS   | 68      |
| Substrate   |         |
| Inoculum  |         |
| Culturing conditions  |         |
| Effect of nitrogen levels on enzyme production during SSF   |         |
| Effect of inoculum age and density on enzyme production in SSF  |         |
| Extraction of crude culture extract   |         |
| Analysis  |         |
| RESULTS AND DISCUSSION  | 71      |
| Effect of nitrogen  |         |
| pH variation  |         |
| Soluble protein   |         |
| Xylanase activity   |         |
| Lignin peroxidase activity  |         |
| Laccase activity  |         |

---

|   |                |
|---|----------------|
| Effect of inoculum age and density (w/w)                        |                |
| pH variation  |                |
| Soluble protein   |                |
| Xylanase activity   |                |
| Lignin peroxidase activity                                      |                |
| Laccase activity  |                |
| <b>CONCLUSION</b>   | <b>88</b>      |
| <b>CHAPTER FIVE</b>   |                |
| <b>ENZYME PRODUCTIVITIES DURING LIGNOCELLULOSIC DEGRADATION</b> |                |
| <b>UNDER OPTIMIZED PARAMETERS</b>                               | <b>90-102</b>  |
| <b>INTRODUCTION</b>   | <b>90</b>      |
| <b>MATERIALS AND METHODS</b>                                    | <b>91</b>      |
| Substrate   |                |
| Inoculum  |                |
| Fermentation  |                |
| Analysis  |                |
| <b>RESULTS AND DISCUSSION</b>                                   | <b>93</b>      |
| pH variation  |                |
| Soluble protein   |                |
| Xylanase activity   |                |
| Lignin peroxidase activity                                      |                |
| Laccase activity  |                |
| <b>CONCLUSION</b>   | <b>102</b>     |
| <b>CHAPTER SIX</b>  | <b>103-110</b> |
| <b>GENERAL DISCUSSION AND CONCLUSION</b>                        | <b>103</b>     |
| <b>RECOMMENDATIONS FOR FURTHER WORK</b>                         | <b>109</b>     |
| <b>CONCLUSION</b>   | <b>110</b>     |
| <b>REFERENCES</b>   | <b>111-125</b> |
| <b>APPENDIX A: ANALYTICAL TECHNIQUES</b>                        | <b>126</b>     |
| <b>APPENDIX B: BUFFERS AND MEDIA</b>                            | <b>134</b>     |
| <b>APPENDIX C: DATA AND STATISTICAL TABLES</b>                  | <b>135</b>     |

---

## LIST OF FIGURES

|            |   |    |
|------------|---|----|
| Figure 1.1 | Utilization of sago starch  | 2  |
| Figure 1.2 | Sago processing at a conventional factory and by-products generated                   | 4  |
| Figure 2.1 | Useful products derived from lignocellulosic materials                                | 10 |
| Figure 2.2 | Recycling of agro-residues  | 15 |
| Figure 2.3 | Principal steps in solid-state fermentation   | 33 |
| Figure 3.1 | Flow chart of the <i>P. sanguineus</i> Koji development using sterilized wheat grains | 49 |
| Figure 3.2 | Procedure for sampling analysis and enzyme assay during SSF of <i>Hampas</i>          | 52 |
| Figure 3.3 | Variation of pH in crude culture extract  | 56 |
| Figure 3.4 | Soluble protein content of crude culture extract                                      | 58 |
| Figure 3.5 | Xylanase activity of crude culture extract  | 60 |
| Figure 3.6 | Lignin peroxidase activity of crude culture extract                                   | 62 |
| Figure 3.7 | Laccase activity of crude culture extract   | 64 |
| Figure 4.1 | Variation of pH in crude culture extract  | 74 |
| Figure 4.2 | Soluble protein content of crude culture extract                                      | 74 |
| Figure 4.3 | Xylanase activity of crude culture extract  | 76 |
| Figure 4.4 | Lignin peroxidase activity of crude culture extract                                   | 78 |
| Figure 4.5 | Laccase activity of crude culture extract   | 78 |
| Figure 4.6 | Variation of pH in crude culture extract  | 80 |
| Figure 4.7 | Soluble protein content of crude culture extract                                      | 82 |
| Figure 4.8 | Xylanase activity of crude culture extract  | 84 |

---

|             |   |     |
|-------------|---|-----|
| Figure 4.9  | Lignin peroxidase activity of crude culture extract                                       | 85  |
| Figure 4.10 | Laccase activity of crude culture extract   | 87  |
| Figure 5.1  | Variation of pH in crude culture extract  | 94  |
| Figure 5.2  | Soluble protein content of crude culture extract  | 96  |
| Figure 5.3  | Xylanase activity of crude culture extract  | 98  |
| Figure 5.4  | Lignin peroxidase activity of crude culture extract                                       | 99  |
| Figure 5.5  | Laccase activity of crude culture extract   | 101 |
| Figure 6.1  | Proposed integrated system for utilization and treatment of sago starch processing wastes | 108 |

---

## LIST OF PLATES

|           |   |    |
|-----------|---|----|
| Plate 3.1 | Sago 'hampas' used in the experiment                            | 48 |
| Plate 3.2 | <i>Pycnopus sanguineus</i> on PDA plate (seven day old culture) | 48 |
| Plate 3.3 | Colonization of 'hampas' by <i>Pycnopus sanguineus</i>          | 54 |
| Plate 4.1 | Growth of <i>Pycnopus sanguineus</i> on 'hampas'                | 72 |

---

## LIST OF TABLES

|           |   |    |
|-----------|---|----|
| Table 2.1 | Constituents of <i>Pycnoporus sanguineus</i> and their approximate concentration                            | 17 |
| Table 2.2 | Extracellular enzymes of basidiomycetes and mode of action on lignocellulose                                | 21 |
| Table 2.3 | Enzymes involved in the conversion of biomass   | 31 |
| Table 2.4 | Industrial applications of enzymes produced by SSF  | 43 |
| Table 3.1 | Initial contents of flask   | 50 |
| Table 4.1 | Proximate analysis of sago 'hampas'   | 67 |
| Table 4.2 | Nitrogen levels tested in the experiment  | 69 |
| Table 4.3 | Experimental design for the SSF of <i>hampas</i> with <i>Pycnoporus sanguineus</i>                          | 70 |
| Table 5.1 | Variation in SSF of sago <i>hampas</i> with <i>P. sanguineus</i> using unoptimized and optimized parameters | 92 |