

References:

- Aggarwal, B.B., Sundaram, C., Malani, N., & Ichikawa, H. (2007). Curcumin: the Indian solid gold, *Medical Biology*, 595, 697-705.
- Arya, S.C., Kapoor, S., Agarwal, N., Bhasin, R., & George, S. (2004). Comments on use of a disc diffusion method with cefoxitin to detect of methicillin-resistant *Staphylococcus aureus*, *European Journal of Clinical Microbiol Infections Diseases*, 23, 867-868.
- Baldassarri, L., Bertuccini, L., Ammendolia, L.G., Arciola, C.R., & Montanaro, L. (2001). Effect of Iron limitation on slime production by *Staphylococcus aureus*, *European Journal of Clinical Microbiology*, 20, 343-345.
- Bauer, A.W., Kirby, W.M.M., Sherris, J.C., & Turck, J.C. (1966). Antibiotic susceptibility testing by a standardized single disc method, *Technical Bulletin of the Registry of Medical Technologists*, 36(3), 49-52.
- Barrueto, L.P., Machado, A.C.M.G., Carvalheira, S.B.R.C., & Brasileiro, A.C.M. (1999). Plant regeneration from seedling explants of *Eucalyptus grandis* x *E.urophylla*, *Plant Cell Tissue Organ Cultrve*, 56, 17-23.
- Boggetti, Jasik, J., & Mantell, S.H. (2001). *In vitro* root formation in *Anacardium occidentale* microshoots, *Biology Plant*, 44, 175-179.
- Butenk, R.G., Upsky, A.K.H., Cherinyak, N.D., & Arya, H.C. (1984). Changes in culture medium pH by cell suspension cultures of *Dioscorea deitoulea*, *Plant Science*, 35, 107-212.
- Chan, E.W.C., Lim, Y.Y., & Omar, M. (2007). Antioxidant and antibacterial activity of leaves of *Etlingera* species (Zingiberaceae) in Peninsular Malaysia, *Food Chemistry*, 104(4), 1586-1593.
- Chen, N.I., Chang, C.C., Wang, C.W., Shyu, Y.T., Chang, T.L., et al. (2008) Antioxidant and antimicrobial activity of Zingiberaceae plants in Taiwan, *Plant Foods for Human Nutrition*, 63, 15-20.
- Cousins, M., Adelberg, J., Chenb, F., & Rieck, J. (2007). Antioxidant capacity of fresh and dried rhizomes from four clones of turmeric (*Curcuma longa* L.) grown *in vitro*, *Industrial Crops and Products* 25(2), 129-135.



References

- Ebert, A., & Taylor, H.F. (1990). Assessment of the changes of 2, 4-dichlorophenoxyacetic acid concentrations in plant tissue culture media in the presence of activated charcoal, *Plant Cell Tissue Organ Culture*, 20, 165-172.
- Ebert, A., Taylor, F., & Blake, J. (1993). Changes of 6-benzylaminopurine and 2, 4-dichlorophenoxyacetic acid concentrations in plant tissue culture media in the presence of activated charcoal, *Plant Cell Tissue Organ Culture*, 33, 157-163.
- Ficker, C.E., Smith, M.L., Susiarti, S., Leaman, D.J., Irawati, C., & Arnason, J.T. (2003). Inhibition of human pathogenic fungi by members of Zingiberaceae used by the Kenyah in Indonesian Borneo, *Journal of Ethnopharmacology*, 85, 289-293.
- García-Sánchez, A., Sánchez, S., Rubio, S., Pereira, G., Alonso, J.M., Mendoza, J.H. & Rey, J. (2007). Presence of shiga toxin-producing *E.coli* O157:H7 in a survey of wild artiodactyls, *Veterinary Microbiology*, 121, 373-377.
- Granum, P.E. & Lund, T. (1997). *Bacillus cereus* and its food poisoning toxins, *FEMS Microbiology Letters*, 157, 223-228.
- George, E.F. (1993). Plant propagation by tissue culture, *The technology*, (2nd ed.), Part 1, 3-160.
- Govindarajan, V. S. (1980). Turmeric-Chemistry, Technology and Quality, *Food Science Nutrition*, 12, 257.
- Guillemot, D. (1999). Antibiotic use in humans and bacterial resistance, *Current Opinion in Microbiology*, 2, 494-498.
- Habsah, M., Amran, M., Mackeen, M.M., Lajis, N.H., Kikuzaki, H., Nakatani, N., et al. (2007). Screening of Zingiberaceae extracts for antimicrobial and antioxidant activities, *Journal of Ethnopharmacology*, 72, 403- 410.
- Heinrich, M., & Gibbons, S. (2004). *Fundamentals of Pharmacognosy and Phytotherapy*, Churchill Livingstone: Edinburgh, 245-252.
- Hennerty, M.J., Upton, M.E., Furlong, P.A., Harris, D.P. (1988). Microbial contamination of *in vitro* cultures of apple rootstocks, *Acta Horticulturae*, 225, 129-137.



References

- Ibrahim, B.J., Yassin, M.S.M., Chin, C.B., Chen, L.L., & Sim, N.E. (2003). Antifungal activity of the essential oils of nine Zingiberaceae species, *Pharmaceutical Biology*, 41(5), 392-397.
- Islam, M.A., Kloppstech, K., and Esch, E. (2005). Population genetic diversity of *Curcuma zedoaria Roscoe* – Conservation prioritised medicinal plant in Bangladesh, *Conservation Genetics*, 6, 1027-1033.
- Islam, M.A., Alam, M.M., Choudhur, M.E., Kobayashi, N., & Ahmed, M.U. (2008). Determination of minimum inhibitory concentration (MIC) of cloxacillin-Resistant *Staphylococcus aureus* with their antibiogram, *Bangladesh Society for Veterinary Medicine*, 6(1), 121-126.
- José, A., Rufián-Henares, A., & Morales, F.J. (2006). New application of a commercial microtiter plate-based assay for assessing the antimicrobial activity of Maillard reaction products, *Food Research International*, 39, 33-39.
- Kalemba, D. & Kunicka, A. (2003). Antibacterial and antifungal properties of essential oils, *Current Medicinal Chemistry*, 10, 813-829.
- Kapoor, R., Sharma, D., & Bhatnagar, A.K. (2008). Arbuscular mycorrhizae in micropropagation systems and their potential applications, *Scientia Horticulturae*, 116, 227-239.
- Keng, L.C., and Stanley, C. (2007). Micropropagation of *Curcuma roscoe* and *zingiber zerumbet smith*, *Biotechnology*, 6, 555-560.
- Komalavalli & Rao, M.V. (2000). *In vitro* micropropagation of *Gymnema sylvestre*-a multipurpose medicinal plant, *Plant Cell Tissue Organ Culture*, 61, 97-105.
- Kotiranta, A., Lounatmaa, k., & Haapasalo, M. (2000). Epidemiology and pathogenesis of *Bacillus cereus* infections, *Microbes and Infection*, 2, 189-198.
- Kintarak, S., Nair, S.P., Speight, P.M., & Whawell, S.A.A. (2004). Recombinant fragment of the fibronectin-binding protein of *Staphylococcus aureus* inhibits keratinocyte migration, *Arch Dermatol Res*, 296, 250-257.



References

- Kuetea, V., Fozing, D.C., Kapche, W.F.G.D., Mbaveng, A.T., Kuatea J.R., Ngadjui, B.T., et al. (2009). Antimicrobial activity of the methanolic extract and compounds from *Morus mesozygia* stem bark, *Journal of Ethnopharmacology*, 124, 551-555.
- Loc, N.H., Duc, D.T., Kwon, T.H., & Yang M.S. (2005). Micropropagation of zedoary (*Curcuma zedoaria* Roscoe) - a valuable medicinal plant, *Plant Cell Tissue Organ Culture*, 81, 119-122.
- Loc, N.H., Ha, T.T.T., & Hirata, Z. (2008). Isolation and characterization of antioxidation enzymes from cells of Zedoary (*Curcuma zedoaria* Roscoe) cultured in a 5-l bioreactor, *Molecular Biotechnology*, 38, 81-87.
- Marcus, Y. (1999). The properties of solvents, *Wiley series in solution chemistry*, 4, 1-32.
- Maua, J.L., Eric, Y.C., Laib, Wangc, N.P., Chena, C.C., Changd, C.H., et al. (2003). Composition and antioxidant activity of the essential oil from *Curcuma zedoaria*, *Food Chemistry*, 82, 583-591.
- Miachir, J.I., Romani, V.L.M., Amaral, A.F.D.C., Mello, M.O., Crocomo, O.J., & Melo, M. (2004). Micropropagation and callogenesis of *Curcuma zedoaria* Roscoe, *Sci. Agric. (Piracicaba, Braz.)*, 61, 42Z-432.
- Murashige, T., & Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures, *Plant physiology*, 15, 473-497.
- Naylor, S.W., Gallyb, D.L. & Low, J.C. (2005). Entero-haemorrhagic *E. coli* in veterinary medicine, *International Journal of Medical Microbiology*, 295, 419-441.
- Niamsa, N., & Sittiwet, C. (2009). Antibacterial activity of *Curcuma longa* aqueous extract, *Journal of Pharmacology and Toxicology*, 4, 173-179.
- Nissen, S.J., & Sutter, E.G. (1990). Stability of IAA and IBA in nutrient medium to several tissue culture procedures, *HortScience*, 25, 800-802.
- Owen, H.V.L., Wengerd, D., & Miller, A.R. (1991). Culture medium pH influenced by basal medium, carbohydrate source, gelling agent, activated charcoal, and medium storage method, *Plant Cell Reports*, 10, 583-586.
- Pan, M.J., & Van Staden, J. (1998). The use of Charcoal in *in vitro* culture- a review, *Plant Growth Regulation*, 26, 155-163.



References

- Rasko, D.A., Altherr, M.R., Han, C.S., & Ravel, J. (2005). Genomics of the *Bacillus cereus* group of organisms, *FEMS Microbiology Reviews*, 29, 303-329.
- Raven, P.H., Evert, R.F., & Eichhorn, S.E. (1999). Micropropagation of orchids, *Biology of Plants* (6th ed.), 794-798.
- Ríos, J.L., & Recio, M.C. (2005). Medicinal plants and antimicrobial activity, *Journal of Ethnopharmacology*, 100, 80-84.
- Ripa, F.A.L., Nahar, M.H., & Islam, M.M. (2009). Antibacterial, cytotoxic and antioxidant activity of crude extract of *Marsilea Quadrifolia*, *European Journal of Scientific Research*, 33(1), 123-129.
- Rojas, R., Bustamante, B., Bauer, J., Fernández, I., Albán, J., & Locka O. (2003). Antimicrobial activity of selected Peruvian medicinal plants, *Journal of Ethnopharmacology*, 88, 199-204.
- Rout, G.A., Samantaray, S., & Das, P. (2000). *In vitro* manipulation and propagation of medicinal plants *Biotechnology Advances*, 18, 91-120.
- Schaffera, A.C., & Lee, J.C. (2008). Vaccination and passive immunisation against *Staphylococcus aureus*, *International Journal of Antimicrobial Agents*, 32(1), 71-78.
- Sharma, R.A., Gescher, A.J., & Steward, W.P. (2005). Curcumin: The story so far, *European Journal of Cancer*, 41, 1955-1968.
- Singh, R.K., Rai, D., Yadav, D., Bhargava, A., Balzarini, J., & Clercq, E.D. (2010). Synthesis, antibacterial and antiviral properties of Curcumin bio-conjugates bearing dipeptide, fatty acids and folic acid, *European Journal of Medicinal Chemistry*, 1-9.
- Singila, S. (1984). Influence of two commercial agars on *in vitro* shoot proliferation of “Almey” crabapple and “Secke” pear. *HonScience*, 19, 227-228.
- Skirvin, R.M., Chu, M.C., Mann, M.L., Young, H., Sullivan, J., & Fermanian, T. (1986). Stability of tissue culture medium pH as a function of autoclaving, time and cultured plant material. *Plant Cell Reports*, 5, 292-294.



References

- Smith, S.M., Andrew, J., Beattie, Gillings, M.R., Marita, P., Holley, et al. (2008). An enhanced miniaturized assay for antimicrobial prospecting, *Journal of Microbiological Methods*, 72, 103-106.
- Stanislavskya, E.S., & Lam, J.S. (1997). *Pseudomonas aeruginosa* antigens as potential vaccines, *FEMS Microbiology Reviews*, 21, 243-277.
- Stickel and Schuppan, D. (2007). Herbal medicine in the treatment of liver diseases, *Digestive and Liver Diseases*, 39, 293-304.
- Tewtrakul, S., & Subhadhirasakul, S. (2007). Anti-allergic activity of some selected plants in the Zingiberaceae family, *Journal of Ethnopharmacology*, 109, 535-538.
- Tiphara, P., Sangvanich, P., Macth, M., & Petsom, A. (2007). Mannose-binding Lectin from *Curcuma zedoaria* Rosc., *Journal of Plant Biology*, 50(2), 167-173.
- Tkenheuer, G.F., Preuss, M., Salzberger, B., Schmeiber, N., Cornely, O.A., Wisplinghoff, H., et al. (2004). Long-term outcome and quality of care of patients with *Staphylococcus aureus* bacteremia, *European journal of clinical microbiology*, 23, 157-162.
- Thomas, D.T. (2008). The role of activated charcoal in plant tissue culture, *Biotechnology Advances*, 26(6), 618-631.
- Tivarekar, S., & Eapen, S. (2001). High frequency plant regeneration from immature cotyledons of mungbean, *Plant Cell Tissue Organ Culture*, 66, 227-230.
- Toit, E.A.D., & Rautenbach, M. (2000). A sensitive standardised micro-gel well diffusion assay for the determination of antimicrobial activity, *Journal of Microbiological Methods*, 42, 159-165.
- Unda, F., Kalynyak, P., & Risem, A. (2007). Organogenesis plant regeneration from leaf explants of *Exacum styer* group, *Plant Cell Tissue Organ Culture*, 89, 105-111.
- Ursic, V., Tomic, V., & Kosnik, M. (2008). Effect of different incubation atmospheres on the production of biofilm in Methicillin-resistant *Staphylococcus aureus* (MRSA) grown in Nutrient-limited medium, *Current Microbiology*, 57, 386-390.



References

- Van, S., Winkle, Johnson, S., & Pullman, G.S. (2003). The impact of gel-rite and activated carbon on the elemental composition of plant tissue culture media, *Plant Cell Reports*, 21, 1175–1182.
- Volk, R.B. (2008). A newly developed assay for the quantitative determination of antimicrobial activity of both hydrophilic and lipophilic test compounds without any restriction, *Microbiological Research*, 163, 161-167.
- Weatherhead, M.A., Burdon, J., & Henshaw, G.G. (1978). Some effects of activated charcoal as an additive to plant tissue culture media, *Z Pflanzenphysiologie*, 89, 141-147.
- Weatherhead, M.A., Burdon, J., & Henshaw, G.G. (1979). Effects of activated charcoal as an additive plant tissue culture media, *Z Pflanzenphysiologie*, 94, 399-405.
- Wilson, B., Abraham, G., Manju, V.S., Mathew, M., Vimala, B., Sundaresan, S., & Nambisan, B. (2005). Antimicrobial activity of *Curcuma zedoaria* and *Curcuma malzbarica* extracts, *Journal of Ethnopharmacology*, 99, 147-151.
- Yam, T.Y., Ernst, R., Arditti, J., Nair, H., & Weatherhead, M.A. (1990). Charcoal in orchid seed germination and tissue culture media: a review, *Lindleyana*, 5, 256-265.
- Yokoigawa, K., Takikawa, A., & HKawai, H. (1999). Difference between *E.coli* 0157:H7 and Non-pathogenic *E. coli*: Survival and growth in seasonings, *Journal of Bioscience and Bioengineering*, 88, 574-576.