

## References

- Ahn, J. H., Pan, J. G. and Rhee, J. S. (2001). Homologous expression of the lipase and ABC transporter gene cluster, tliDEFA, enhances lipase secretion in *Pseudomonas* spp. *Applied and Environmental Microbiology*, 67:5506-5511.
- Amara, A. A. and Rehm, B. H. A. (2003). Replacement of the catalytic nucleophile Cys-296 by serine in Class II polyhydroxyalkanoate synthase from *Pseudomonas aeruginosa* - mediated synthesis of a new polyester: Identification of catalytic residues. *Biochemical Journal*, 374:413-421.
- Anderson, A. J. and Dawes, E. A. (1990). Occurrence, metabolism, metabolic role, and industrial uses of bacterial polyhydroxyalkanoates. *Microbiological Reviews*, 54:450-472.
- Aneja, K. K., Ashby, R. D. and Solaiman, D. K. Y. (2009). Altered composition of *Ralstonia eutropha* poly(hydroxyalkanoate) through expression of PHA synthase from *Allochromatium vinosum* ATCC 35206. *Biotechnology Letters*, 31:1601-1612.
- Albertsson, A. C., Barenstedt, C. and Karlsson, S. (1994). Abiotic degradation products from enhanced environmentally degradable polyethylene. *Acta Polymerica*, 45:97-103.
- Antonio, R. V., Steinbüchel, A. and Rehm, B. H. A. (2000). Analysis of in vivo substrate specificity of the PHA synthase from *Ralstonia eutropha*: formation of novel copolyesters in recombinant *Escherichia coli*. *FEMS Microbiology Letters*, 182:111-117.
- Ashby, R. D. and Foglia, T. A. (1998). Poly(hydroxyalkanoate) biosynthesis from triglyceride substrate. *Applied Microbiology and Biotechnology*, 49:431-437.
- Binet, R., Létoffé, S., Ghigo, J. M., Delepelaire, P. Wandersman, C. (1997). Protein secretion by Gram-negative bacterial ABC exporters-a review. *Gene*, 192:7-11.
- Bitter, W., Koster, M., Latijnhouwers, M., de Cock, H. and Tommassen, J. (1998). Formation of oligomeric rings by XcpQ and PilQ, which are involved in protein transport across the outer membrane of *Pseudomonas aeruginosa*. *Molecular Microbiology*, 27:209-219.
- Brandl, H., Gross, R. A., Lenz, R. W. and Fuller, R. C. (1988). *Pseudomonas oleovorans* as a source of poly(beta-hydroxyalkanoates) for potential applications as biodegradable polyesters. *Applied and Environmental Microbiology*, 54:1977-1982.
- Brandl, H., Gross, R. A., Lenz, R. W. and Fuller, R. C. (1990). Plastics from bacteria and for bacteria: Poly( $\beta$ -Hydroxyalkanoates) as natural, biocompatible, and biodegradable polyesters. *Advances in Biochemical Engineering/ Biotechnology*, 41:77-93.
- Blum, P., Ory, J., Bauernfeind, J. and Krska, J. (1992). Physiological consequences of DNAK and DNAJ overproduction in *Escherichia coli*. *Journal of Bacteriology*, 174:7436-7444.

Brown, T. A. (2006). The application of Gene cloning and DNA analysis in research. *Gene cloning and DNA analysis: An introduction* (5th edition) (pp. 163-221). Blackwell Publishing Ltd., United kingdom.

Byrom, D. (1987). Polymer synthesis by microorganisms: technology and economics. *Trends in Biotechnology*, 5:246-250.

Cai, L., Tuan, M. Q., Liu, F., Jian, J. and Chan, G. Q. (2009). Enhanced production of medium-chain-length polyhydroxyalkanoates (PHA) by PHA depolymerase knockout mutant of *Pseudomonas putida* KT2442. *Bioresource Technology*, 100:2265-2270.

Chen, G. Q. (2001). Introduction of bacterial plastics PHA, PLA, PBS, PE, PTT, and PPP. In Chen, G. Q. (ed.), *Plastics from bacteria, natural functions and applications* (pp. 1-16). Springer-verlag, Berlin Heidelberg, Germany.

Chen, G. Q., Xu, J. Wu, Q., Zhang, Z. M. and Ho, K. P. (2001). Synthesis of co-polyesters consisting of medium-chain-length  $\beta$ -hydroxyalkanoates by *Pseudomonas stutzeri* 1317. *Reactive and Functional Polymers*, 48:107-121.

Chihara-Siami, M., Yoshikawa, K., Oshima-Hirayama, N., Yamamoto, K., Sogabe, Y., Nakatani, T., Nishioka, T. and Oda, Junichi. (1992). Purification, molecular cloning, and expression of lipase from *Pseudomonas aeruginosa*. *Archives of Biochemistry and Biophysics*, 296:505-513.

Cooper, M. L., Hansbrough, J. F., Spievogel, R. L., Cohen, R., Bartel, R. L. and Naughton, G. (1991). *In vivo* optimization of a living dermal substitute employing cultured human fibroblasts on a biodegradable polyglycolic acid or polyglactin mesh. *Biomaterials*, 12:243-248.

Cruz-Pinto, J. J. C., Carvalho, M. E. S., Ferreira, J. F. A. (1994). The kinetics and mechanism of polyethylene photo-oxidation. *Die Angewandte Makromolekulare Chemie*, 216:113-133.

De Eugenio, L. I., García, P., Luengo, J. M., Sanz, J. M., Román, J. S., García, J. L. and Prieto, M. A. (2007). Biochemical evidence that *phaZ* encodes a specific intracellular medium chain length polyhydroxyalkanoate depolymerase in *Pseudomonas putida* KT2442. *The Journal of Biological Chemistry*, 282:4951-4962.

Deretic, V., Konyecsni, W. M., Mohr, C. D., Martin, D. W. and Hibler, N. S. (1989). Common denominators of promoter control *Pseudomonas* and other bacteria. *Biotechnology*, 7:1249-1254.

Doi, Y., Kitamura, S. and Abe, H. (1995). Microbial synthesis and characterization of poly(hydroxybutyrate-co-hydroxyhexanoate). *Macromolecules*, 23:4822-4828.

- Dong, H., Nilsson, L. and Kurland, C. G. (1995). Gratuitous overexpression of genes in *Escherichia coli* leads to growth inhibition and ribosome destruction. *Journal of Bacteriology*, 177:1497-1504.
- Duong, F., Soscia, C., Lazdunski, A. and Murgier, M. (1994). The *Pseudomonas fluorescens* lipase has a C-terminal secretion signal and is secreted by a three-component bacterial ACB-exporter system. *Molecular Microbiology*, 11:1117-1126.
- Elvin, C. M., Thompsan, P. R., Argall, M. E., Hendry, P., Stamford, N. P., Lilley, P. E. and Dixon, N. E. (1990). Modified bacteriophage lambda promoter vectors for overproduction of proteins in *Escherichia coli*. *Gene*, 87:123-126.
- Eggink, G., de Waard, P. and Huilberts, G. N. M. (1992). The role of fatty acid biosynthesis and degradation in the supply of substrates for poly(3-hydroxyalkanoate) formation in *Pseudomonas putida*. *FEMS Microbiology Reviews*, 103:159-164.
- Eggink, G., Waard, P. de and Huijberts, G. N. M. (1995). Formation of novel poly(hydroxyalkanoates) from long-chain fatty acids. *Canadian Journal of Microbiology*, 41:14-21.
- Fakirov, S. and Bhattacharyya, D. (2007). Plastics, utilization and environmental impact. In Fakirov, S., and Bhattacharyya, D. (eds.), *Handbook of engineering biopolymers: Homopolymers, blends and composites* (pp. 109-150). Carl Hanser Verlag, Munich, Germany.
- Fiedler, S., Steinbüchel, A. and Rehm., B. H. A. (2000). PhaG mediated synthesis of poly(3-hydroxyalkanoates) consisting of medium-chain-length constituents from nonrelated carbon sources in recombinant *Pseudomonas fragi*. *Applied and Environment Microbiology*, 66:2117-2124.
- Fiedler, S., Steinbüchel, A. and Rehm, B. H. A. (2002). The role of the fatty acid beta-oxidation multi-enzyme complex from *Pseudomonas oleovorans* in polyhydroxyalkanoate biosynthesis: molecular characterization of the *fadBA* operon from *P. oleovorans* and of the enoyl-CoA hydratase genes *phaJ* from *P. oleovorans* and *Pseudomonas putida*. *Archives of Microbiology*, 178:149-60.
- Füchtenbusch, B., Wullbrandt, D. and Steinbüchel, A. (2000). Production of polyhydroxyalkanoic acids by *Ralstonia eutropha* and *Pseudomonas oleovorans* from oil remaining from biotechnological rhamnase production. *Applied Microbiology and Biotechnology*, 53:167-172.
- Fukui, T. and Doi, Y. (1998). Efficient production of polyhydroxyalkanoates from plant oils by *Alcaligenes eutrophus* and its recombinant strain. *Applied Microbiology and Biotechnology*, 49:333-336.

- Galegoa, N., Rozsaa, C., Sánchez, R., Fung, J., Vázquez, A. and Tomás, J. S. (2000). Characterization and application of poly(b-hydroxyalkanoates) family as composite biomaterials. *Polymer Testing*, 19:485-492.
- Guzman, L., Belin, D., Carson, M. J. and Beckwith, J. (1995). Tight regulation, modulation, and high-level expression by vectors containing the arabinose pBAD promoter. *American Society for Microbiology*, 117:4121-4130.
- Hasan, F., Shah, A. A. and Hameed, A. (2006). Industrial applications of microbial lipases. *Enzyme and Microbial Technology*, 39:235-251.
- Henderson, I. R., Navarro-Garcia, F. and Nataro, J. P. (1998). The great escape: structure and function of the autotransporter proteins. *Trends in Microbiology*, 6:370-378.
- Higgins, C. F. (1992). ABC Transporters: From Microorganism to Man. *Annual Reviews Cell and Developmental Biology*, 8:67-113.
- Hoffmann, N., Steinbüchel, A. and Rehm, B. H. A. (2000). The *Pseudomonas aeruginosa* *phaG* gene product is involved in the synthesis of polyhydroxyalkanoic acid consisting of medium-chain-length constituents from non-related carbon sources. *FEMS Microbiology Letters*, 184:253-259.
- Huijberts, G. N. M., Eggink, G., de Waard, P., Huisman, G. W. and Witholt, B. (1992). *Pseudomonas putida* KT2442 cultivated on glucose accumulates poly(3-hydroxyalkanoates) consisting of saturated and unsaturated monomers. *Applied and Environmental Microbiology*, 58:536-544.
- Huisman, G.W., Leeuw, O. D., Enggink, G. and Witholt, B. (1989). Synthesis of poly-3-hydroxyalkanoates is a common feature of fluorescent pseudomonads. *Applied and Environmental Microbiology*, 55:1949-1954.
- Ishuang, S. L., Crane, G. M., Miller, M. J. and Yasko, A. W. (1997). Bone formation by three-dimensional stromal osteoblast culture in biodegradable polymer scaffolds. *Journal of Biomedical Materials Research*, 36:17-28.
- Jaeger, K. E., Ransac, S., Dijkstra, B. W., Colson, C., Van Heuvel, M. and Misset, O. (1994). Bacterial lipase. *FEMS Microbiology Reviews*, 15:29-63.
- Jaeger, K. E. and Reetz, M. T. (1998). Microbial lipases form versatile tools for biotechnology. *Trends in Biotechnology*, 16:396-403.
- Jendrossek, D., and Handrick, R. (2002). Microbial degradation of polyhydroxyalkanoates. *Annual Review of Microbiology*, 56:403-432.
- Jia, Y., Yuan, W., Wodzinska, J., Park, C., Sinskey, A. J. and Stubbe, J. (2001). Mechanistic studies on class I polyhydroxybutyrate (PHB) synthase from *Ralstonia*

*eutropha*: Class I and III synthases share a similar catalytic mechanism. *Biochemistry*, 40:1011-1019.

Khanna, S. and Srivastava, A. K. (2005). Recent advances in microbial polyhydroxyalkanoates. *Process Biochemistry*, 40:607-619.

Kessler, B. and Witholt, B. (2001). Factors involved in the regulatory network of polyhydroxyalkanoate metabolism. *Journal of Biotechnology*, 86:97-104.

Kim, D. Y., Kim, Y. B. and Rhee, Y. H. (2000). Evolution of various carbon substrates for the biosynthesis of polyhydroxyalkanoates bearing functional groups by *Pseudomonas putida*. *International Journal of Biological Macromolecules*, 28:23-29.

Kinoshita, Y., Kirigakubo, M., Kobayashi, M., Tabata, T., Shimura, K. and Ikada, Y. (1993). Study on the efficacy of biodegradable poly(L-lactide) mesh for supporting transplanted particulate cancellous bone and marrow: Experiment involving subcutaneous implantation in dogs. *Biomaterials*, 14:729-736.

Kolibachuk, D., Miller, A. and Dennis, D. (1999). Cloning, molecular analysis, and expression of the polyhydroxyalkanoic acid synthase (*phaC*) gene from *Chromobacterium violaceum*. *Applied and Environmental Microbiology*, 65:3561-3565.

Kouker, G. and Jaeger, K. E. (1987). Specific and sensitive plate assay for bacterial lipases. *Applied and Environmental Microbiology*, 53:211-213.

Lageveen, R. G., Huiman, G. W., Preusting, H., Ketelaar, P., Enggink, G. and Witholt, B. (1988). Formation of polyesters by *Pseudomonas oleovorans*: Effect of substrates on the formation and composition of poly-(R)-3-hydroxyalkanoates and poly-(R)-3-hydroxyalkenoates. *Applied and Environmental Microbiology*, 54:2924-2932.

Langenbach, S., Rehm, B. H. A. and Steinbüchel, A. (1997). Functional expression of the PHA synthase gene (*phaC1*) from *Pseudomonas aeruginosa* in *Escherichia coli* results in poly(3-hydroxyalkanoate) synthesis. *FEMS Microbiology Letters*, 150:303-309.

Larsen, J. E. L., Gerdes, K., Light, J. and Molin, S. (1984). Low copynumber plasmid cloning vectors amplifiable by de-repression of an inserted foreign promoter. *Gene*, 28:45-54.

Ljungberg, N. and Wesslén, B. (2005). Preparation and properties of plasticized poly(lactic acid) films. *Biomacromolecules*, 6: 1789-1796.

Lee S. Y. (1996). Plastic bacteria? Progress and prospects for polyhydroxyalkanoate production in bacteria. *Trends in Biotechnology*, 14:431-438.

Loss, M., Wedler, V., Künzi, W., Simmen, C. M. and Meyer, V. E. (2000). Artificial skin, split-thickness auto-graft and cultured autologous keratinocytes combined to treat a severe burn injury of 93% of TBSA. *Burns*, 26:644-652.

- Koning, G. D. and Witholt, B. (1996). A biodegradable rubber from bacteria, poly(hydroxyalkanoate) from *Pseudomonas*. *Materials Science and Engineering C*, 4:121-124.
- Luengo, J. M., García, B., Sandoval, A., Naharro, G. and Olivera, E. R. (2003). Bioplastics from microorganisms. *Current Opinion in Microbiology*, 6:251-260.
- Lu, X. Y., Zhang, J. Y., Wu, Q. and Chen, G. Q. (2003). Enhanced production of poly (3-hydroxybutyrate-co-3-hydroxyalkanoates) via manipulating the fatty acid  $\beta$ -oxidation pathway in *E.coli*. *FEMS Microbiology Letters*, 221:97-101.
- Ma, P. X. and Lui, X. H. (2003). Polymeric scaffolds for bone tissue engineering. *Annals of Biomedical Engineering*, 32:477-486.
- Madison, L. L. and Huisman, G. W. (1999). Metabolic engineering of poly (3-hydroxyalkanoates): From DNA to plastic. *Microbiology and Molecular Biology Reviews*, 63:21-53.
- Marsudi, S., Tan, I. K. P., Gan, S. N. and Ramachandran, K. B. (2007). Production of medium chain length polyhydroxyalkanoates from oleic acid using *Pseudomonas putida* PGA1 by fed batch culture. *Makara Tecknologi*, 11:1-4.
- Miyada, C. G., Stoltzfus, L. and Wilcox, G. (1984). Regulation of the *araC* gene of *Escherichia coli*: Catabolite repression, auto-regulation, and effect on *araBAD* expression. *Proceedings of the National Academy of Sciences*, 81:4120-4124.
- Nebe, B., Forster, C., Pommerenke, H., Fulda, G., Behrend, D., Bernewski, U., Schmitz, K. P. and Rychly, J. (2001). Structural alterations of adhesion mediating components in cells cultured on poly beta-hydroxybutyric acid. *Biomaterials*, 22:2425-34.
- Ostle, A. G. and Holt, J. G. (1982). Nile Blue A as a fluorescent stain for poly-beta-hydroxybutyrate. *Applied and Environmental Microbiology*, 44:238-241.
- Park, S. J., Chaoi, J. I. and Lee, S. Y. (2005). Engineering of *Escherichia coli* fatty acid metabolism for the production of polyhydroxyalkanoates. *Enzymes and Microbial Technology*, 36:579-588.
- Poirier, Y., Nawrath, C. and Somerville, C. (1995). Production of polyhydroxyalkanoates, a family of biodegradable plastics and elastomers, in bacteria and plants. *Biotechnology*, 13:142-150.
- Pugsley A.P. (1993). The complete secretory pathway in Gram-negative bacteria, *Microbiological Reviews*, 57:50-108.
- Qi, Q., Rehm, B. H. and Steinbüchel, A. (1997). Synthesis of poly(3-hydroxyalkanoates) in *Escherichia coli* expressing the PHA synthase gene *phaC2* from *Pseudomonas aeruginosa*: comparison of PhaC1 and PhaC2. *FEMS Microbiology Letters*, 157:155-62.

- Qi, Q., Steinbüchel, A. and Rehm, B. H. A. (2000). *In vitro* synthesis of poly(3-hydroxydecanoate): Purification and enzymatic characterization of type II polyhydroxyalkanoate synthases *phaC1* and *phaC2* from *Pseudomonas aeruginosa*. *Applied Microbiology and Biotechnology*, 54:37-43.
- Reddy, C. S. K., Ghai, R. and Kalai, V. C. (2003). Polyhydroxyalkanoates: an overview. *Bioresource Technology*, 87:137-146.
- Rehm, B. H. A. (2003). Polyester synthase: Natural catalysts for plastics. *Biochemical Journal*, 376:15-33.
- Rehm, B. H. A., Antonio, R. V., Spiekermann, P., Amara, A. A. and Steinbüchel, A. (2002). Molecular characterization of the poly (3-hydroxybutyrate) (PHB) synthase from *Ralstonia eutropha*: *In vitro* evolution, site specific mutagenesis and development of a PHB synthase protein model. *Biochimica et Biophysica Acta*, 1549:178-190.
- Rehm, B. H. A., Kruger, N. and Steinbüchel, A. (1998). A new metabolic link between fatty acid de novo synthesis and polyhydroxyalkanoic acid synthesis. *The Journal of Biological Chemistry*, 273:24044-24051.
- Rehm, B. H. and Steinbüchel, A. (1999). Biochemical and genetic analysis of PHA synthases and other proteins required for PHA synthesis. *International Journal of Biological Macromolecules*, 25:3-19.
- Ren, Q., Roo, G. D., Beilen, J. B. V., Zinn, M., Kessler, B. and Witholt, B. (2005). Poly(3-hydroxyalkanoate) polymerase synthesis and *in vitro* activity in recombinant *Escherichia coli* and *Pseudomonas putida*. *Applied Microbiology and Biotechnology*, 69:286-292.
- Ren, Q., Sierro, N., Kellerhals, M., Kessler, B. and Witholt, B. (2000). Properties of engineered poly-3-hydroxyalkanoates produced in recombinant *Escherichia coli* strains. *Applied and Environmental Microbiology*, 66:1311-1320.
- Ren, Q., Van Beilen, J. B., Zinn, M., Kessler, B. and Witholy, B. (2005). Expression of PHA polymer genes of *Pseudomonas putida* in *Escherichia coli* and its effect on PHA formation. *Antonie van Leeuwenhoek*, 87:91-100.
- Rosenau, F. and Jaeger, K. E. (2000). Bacterial lipases from *Pseudomonas*: Regulation of gene expression and mechanisms of secretion. *Biochimie*, 82:1023-1032.
- Sambrook, J. and Russel, D. W. (2001). Plasmids and their usefulness in molecular cloning. *Molecular cloning: A laboratory manual* (2nd ed.) (pp. 107-111). Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, United State.
- Saxena, R. K., Ghosh, P. K., Gupta, R., Davidson, W. S., Bradoo, S. and Gulati, R. (1999). Microbial lipases: Potential biocatalysts for the future industry. *Current Science*, 77:101-115.

- Schleif, R. (2000). Regulation of the L-arabinose operon of *Escherichia coli*. *Trends in Genetics*, 16:559-565.
- Solaiman, D. K. Y., Ashby, R. D. and Foglia T. A. (1999). Medium-chain-length poly( $\beta$ -hydroxyalkanoate) synthesis from triacylglycerols by *Pseudomonas saccharophila*. *Current Microbiology*, 38:151-154.
- Solaiman, D. K. Y., Ashby, R. D. and Foglia, T. A. (2001). Production of polyhydroxyalkanoates from intact triacylglycerols by genetically engineered *Pseudomonas*. *Applied Microbiology and Biotechnology*, 156:664-669.
- Spyros, A. and Marchessault, R. H. (1996). Segmental dynamics in poly(3-hydroxybutyrate-co-4-hydroxybutyrate)s above the glass transition temperature:  $^{13}\text{C}$  nuclear magnetic relaxation in the amorphous phase. *Macromolecules*, 29:2479-2486.
- Steinbüchel, A. (1991). Polyhydroxyalkanoic acids. In D. Byrom (ed.), *Biomaterials, novel materials from biological sources* (pp. 123-213). Macmillan Publishers Ltd., Basingstoke, England.
- Steinbüchel, A., Fuchtenbush, B., Gorenflo, V., Hein, S., Jossek, R., Langenbach, S. and Rehm, B. H. A. (1998). Biosynthesis of polyester in bacteria and recombinant organisms. *Polymer Degradation and Stability*, 59:177-182.
- Steinbüchel, A. and Valentin, H. E. (1995). Diversity of bacterial polyhydroxyalkanoic acids. *FEMS Microbiology Letters*, 129:219-228.
- Stubbe, J. and Tian, J. (2003). Polyhydroxyalkanoate (PHA) homeostasis: The role of PHA synthase. *Natural Product Reports*, 20:445-57.
- Sudesh, K., Abe, H. and Doi, Y. (2000). Synthesis, structure and properties of polyhydroxyalkanoates: Biological polyesters. *Progress in Polymer Science*, 25:1503-1555.
- Suyatma, N. E., Copinet, A., Tighzert, L. and Coma, V. (2004). Mechanical and barrier properties of biodegradable films made from chitosan and poly(lactic acid) blends. *Journal of Polymers and the Environment*, 12:1-6.
- Taguchi, K., Aoyagi, Y., Matsusaki, H., Fukui, T. and Doi, Y. (1999). Co-expression of 3-ketoacyl-ACp reductase and polyhydroxyalkanoate synthase genes induces PHA production in *Escherichia coli* HB101 strain. *FEMS Microbiology Letters*, 176:183-190.
- Tajima, K., Igari, T., Nishimura, D., Nakamura, M., Satoh, Y. and Munekata, M. (2003). Isolation and characterization of *Bacillus* sp. INT005 accumulating polyhydroxyalkanoate (PHA) from gas field soil. *Journal of Bioscience and Bioengineering*, 95:77-81.
- Tamura, K., Dudley, J., Nei, M. and Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0.



- Tan, I. K. P., Sudesh, K., Theanmalar, M., Gan, S. N. and Gordon, III B. (1997). Saponified palm kernel oil and its major free fatty acids as carbon substrates for the production of polyhydroxyalkanoates in *Pseudomonas putida* PGA1, *Applied Microbiology and Biotechnology*, 47:207-211.
- Tan, Y. P. and Miller, K. J. (1992). Cloning, expression, and nucleotide sequence of a lipase gene from *Pseudomonas fluorescens* B52. *Applied and Environmental Microbiology*, 52:1402-1407.
- Trainer, M. A. and Charles, T. C. (2006). The role of PHA metabolism in the symbiosis of rhizobia with legumes. *Applied Microbiology and Biotechnology*, 71:377-386.
- Timm, R. and Steinbüchel, A. (1990). Formation of polyester consisting of medium-chain-length 3-hydroxyalkanoic acids from gluconate by *Pseudomonas aeruginosa* and other fluorescent pseudomonads. *Applied and Environmental Microbiology*, 56:3360-3367.
- Ueda, H. and Tabata, Y. (2003). Polyhydroxyalkanoate derivation in current clinical application and trails. *Advanced Drug Delivery Reviews*, 55:501-518.
- Van der Veen, V. C., Van der Wal, M. B. A., Van Leeuwen, M. C. E., Ulrich, M. M. W. and Middelkoop, E. (2010). Biological background of dermal substitutes. *Burns*, 36:305-321.
- Wang, H. H., Li, X. T. and Chen, G. Q. (2009). Production and characterization of homopolymer polyhydroxyheptanoate (P3HHp) by a *fadBA* knockout mutant *Pseudomonas putida* KTOY06 derived from *P. putida* KT2442. *Process Biochemistry*, 44:106-111.
- Wang, J. G. and Bakken, L. R. (1998). Screening of soil bacteria for poly-beta-hydroxybutyric acid production and its role in the survival of starvation. *Microbial Ecology*, 35:94-101.
- Witholt, B. and Kessler, B. (1999). Perspectives of medium chain length poly(hydroxyalkanoates), a versatile set of bacterial bio-plastics. *Current Opinion in Biotechnology*, 10:279-285.
- Zinn, M., Witholt, B. and Egli, T. (2001). Occurrence, synthesis and medical application of bacterial polyhydroxyalkanoate. *Advanced Drug Delivery Reviews*, 53:5-21.
- Zhang, J. F. and Sun, X. J. (2004). Mechanical properties of poly(lactic acid)/starch composites compatibilized by maleic anhydride. *Biomacromolecules*, 5:1446-1451.
- Zhao, H. Y., Li, H. M., Qin, L. F., Wang, H. H. and Chen, G. Q. (2007). Disruption of the polyhydroxyalkanoate synthase gene in *Aeromonas hydrophila* reduces its survival ability under stress conditions. *FEMS Microbiology Letters*, 276:34-41.
- Zhao, K., Deng, Y., Chen, J. C. and Chen, G. Q. (2003). Polyhydroxyalkanoate (PHA) scaffolds with good mechanical properties and biocompatibility. *Biomaterials*, 24:1041-1045.

