

Chapter Twelve

References

- Aanensen, D. M., & Spratt, B. G. (2005). The multilocus sequence typing network: mlst.net. *Nucleic Acids Res*, 33, 728-733.
- Achtman, M. (2001). A phylogenetic perspective on molecular epidemiology. In M. Sussmann (ed.), *Molecular medical microbiology*. Academic Press, San Diego, Calif. pp. 485-509.
- Alam, M., Sadique, A., Nur, A. H., Bhuiyan, N. A., Nair, G. B., Siddique, A. K., *et al.* (2006). Effect of transport at ambient temperature on detection and isolation of *Vibrio cholerae* from environmental samples. *Appl Environ Microbiol*, 72(3), 2185-2190.
- Angelichio, M. J., Spector, J., Waldor, M. K., & Camilli, A. (1999). *Vibrio cholerae* intestinal population dynamics in the suckling mouse model of infection. *Infect Immun*, 67(8), 3733-3739.
- Ansaruzzaman, M., Chowdhury, A., Bhuiyan, N. A., Sultana, M., Safa, A., Lucas, M., *et al.* (2008). Characteristics of a pandemic clone of O3 : K6 and O4 : K68 *Vibrio parahaemolyticus* isolated in Beira, Mozambique. *J Med Microbiol*, 57(Pt 12), 1502-1507.
- Armstrong, G. L., Hollingsworth, J., & Morris, J. G. Jr. (1998). Bacterial foodborne disease. In *Bacterial infectious of humans: Epidemiology and control* (3rd Ed), by Evan, A. S., & Brachman, P. S. (eds). p 123-125.
- Bina, J., Zhu, J., Dziejman, M., Faruque, S., Calderwood, S., & Mekalanos, J. (2003). ToxR regulon of *Vibrio cholerae* and its expression in vibrios shed by cholera patients. *Proc Natl Acad Sci U S A*, 100(5), 2801-2806.
- Bisharat, N., Cohen, D. I., Harding, R. M., Falush, D., Crook, D. W., Peto, T., *et al.* (2005). Hybrid *Vibrio vulnificus*. *Emerg Infect Dis*, 11(1), 30-35.
- Boyce, J. D., Cullen, P. A., & Adler, B. (2004). Genomic-scale analysis of bacterial gene and protein expression in the host. *Emerg Infect Dis*, 10(8), 1357-1362.
- Castaneda, N. C., Pichel, M., Orman, B., Binsztein, N., Roy, P. H., & Centron, D. (2005). Genetic characterization of *Vibrio cholerae* isolates from Argentina by *V. cholerae* repeated sequences-polymerase chain reaction. *Diagn Microbiol Infect Dis*, 53(3), 175-183.
- Chakraborty, S., Mukhopadhyay, A. K., Bhadra, R. K., Ghosh, A. N., Mitra, R., Shimada, T., *et al.* (2000). Virulence genes in environmental strains of *Vibrio cholerae*. *Appl Environ Microbiol*, 66(9), 4022-4028.

- Chakraborty, S., Garg, P., Ramamurthy, T., Thungapathra, M., Gautam, J. K., Kumar, C., *et al.* (2001). Comparison of antibiogram, virulence genes, ribotypes and DNA fingerprints of *Vibrio cholerae* of matching serogroups isolated from hospitalised diarrhoea cases and from the environment during 1997-1998 in Calcutta, India. *J Med Microbiol*, 50(10), 879-888.
- Chaparro, A. P., Ali, S. K., & Klose, K. E. (2010). The ToxT-dependent methyl-accepting chemoreceptors AcfB and TcpI contribute to *Vibrio cholerae* intestinal colonization. *FEMS Microbiol Lett*, 302(2), 99-105.
- Bustin, S. A. (2000). Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol*, 25(2), 169-193.
- Caipang, C. M., Brinchmann, M. F., & Kiron, V. (2009). Profiling gene expression in the spleen of Atlantic cod, *Gadus morhua* upon vaccination with *Vibrio anguillarum* antigen. *Comp Biochem Physiol B Biochem Mol Biol*, 153(3), 261- 267.
- Cam, D. T. V., Hao, N. V., Dierckens, K., Defoirdt, T., Boon, N., & Sorgeloos, *et al.* (2009). Novel approach of using homoserine lactone-degrading and poly-b-hydroxy-butyrate-accumulating bacteria to protect *Artemia* from the pathogenic effects of *Vibrio harveyi*. *Aquaculture*, 291, 23–30.
- Canigral, I., Moreno, Y., Alonso, J. L., Gonzalez, A., & Ferrus, M. A. (2010). Detection of *Vibrio vulnificus* in seafood, seawater and wastewater samples from a Mediterranean coastal area. *Microbiol Res*, 165(8), 657-664.
- CDC, PulseNet USA. (2006). The national molecular subtyping network for foodborne disease surveillance: Rapid standard laboratory protocol for molecular subtyping of *Vibrio cholerae* by Pulse-Field Gel Electrophoresis (PFGE).
- Chakraborty, R., Sinha, S., Mukhopadhyay, A. K., Asakura, M., Yamasaki, S., Bhattacharya, S. K., Nair, G. B., & Ramamurthy, T. (2006). Species-specific identification of *Vibrio fluvialis* by PCR targeted to the conserved transcriptional activation and variable membrane and other regions of the *toxR* gene. *J Med Microbiol*, 55, 805-808.
- Charlesworth, D., & Charlesworth, B. (1995). Transposable elements in inbreeding and outbreeding populations. *Genetics*, 140(1), 415-417.
- Chen, C. H., Shimada, T., Elhadi, N., Radu, S., & Nishibuchi, M. (2004). Phenotypic and genotypic characteristics and epidemiological significance of ctx+ strains of *Vibrio cholerae* isolated from seafood in Malaysia. *Appl Environ Microbiol*, 70(4), 1964-1972.
- Chokesajjawatee, N., Zo, Y. G., & Colwell, R. R. (2008). Determination of clonality and relatedness of *Vibrio cholerae* isolates by genomic fingerprinting, using long-range repetitive element sequence-based PCR. *Appl Environ Microbiol*, 74(17), 5392-5401.

- Choi, S. Y., Lee, J. H., Jeon, Y.-S., Lee, H. R., Kim, E. J., Ansaruzzaman, M., *et al.* (2010). Multilocus variable-number tandem repeat analysis of *Vibrio cholerae* O1 El Tor strains harbouring classical toxin B. *J Med Microbiol*, 59(7), 763-769.
- Chow, K. H., Ng, T. K., Yuen, K. Y., & Yam, W. C. (2001). Detection of RTX toxin gene in *Vibrio cholerae* by PCR. *J Clin Microbiol*, 39(7), 2594-2597.
- Choopun, N., Louis, V., Huq, A., & Colwell, R. R. (2002). Simple procedure for rapid identification of *Vibrio cholerae* from the aquatic environment. *Appl Environ Microbiol*, 68(2), 995-998.
- Colwell, R. R., Huq, A., Chowdhury, M. A., Brayton, P. R., & Xu, B. (1995). Serogroup conversion of *Vibrio cholerae*. *Can J Microbiol*, 41(10), 946-950.
- Colwell, R. R., & Huq, A. (1994). Environmental reservoir of *Vibrio cholerae*. The causative agent of cholera. *Ann N Y Acad Sci*, 740, 44-54.
- Cooper, J. E., & Feil, E. J. (2004). Multilocus sequence typing--what is resolved? *Trends Microbiol*, 12(8), 373-377.
- Cooper, K. L., Luey, C. K., Bird, M., Terajima, J., Nair, G. B., Kam, K. M., *et al.* (2006). Development and validation of a PulseNet standardized pulsed-field gel electrophoresis protocol for subtyping of *Vibrio cholerae*. *Foodborne Pathog Dis*, 3(1), 51-58.
- Dalmasso, A., Civera, T., & Bottero, M. T. (2009). Multiplex primer-extension assay for identification of six pathogenic vibrios. *Int J Food Microbiol*, 129(1), 21-25.
- Dalsgaard, A., Forslund, A., Mortensen, H. F., & Shimada, T. (1998). Ribotypes of clinical *Vibrio cholerae* non-O1 non-O139 strains in relation to O-serotypes. *Epidemiol Infect*, 121(3), 535-545.
- Dalsgaard, A., Serichantalergs, O., Forslund, A., Lin, W., Mekalanos, J., Mintz, E., *et al.* (2001). Clinical and environmental isolates of *Vibrio cholerae* serogroup O141 carry the CTX phage and the genes encoding the toxin-coregulated pili. *J Clin Microbiol*, 39(11), 4086-4092.
- Danin-Poleg, Y., Cohen, L. A., Gancz, H., Broza, Y. Y., Goldshmidt, H., Malul, E., *et al.* (2007). *Vibrio cholerae* strain typing and phylogeny study based on simple sequence repeats. *J Clin Microbiol*, 45(3), 736-746.
- Dumontet, S., Krovacek, K., Svenson, S. B., Pasquale, V., Baloda, S. B., & Figliuolo, G. (2000). Prevalence and diversity of *Aeromonas* and *Vibrio* spp. in coastal waters of Southern Italy. *Comp Immunol Microbiol Infect Dis*, 23(1), 53-72.
- Elhadi, N., Radu, S., Chen, C. H., & Nishibuchi, M. (2004). Prevalence of potentially pathogenic *Vibrio* species in the seafood marketed in Malaysia. *J Food Prot*, 67(7), 1469-1475.

- Elizabeth, J., Kelkar, P. N., & Weishali, G. (1995). Yellow phosphorus poisoning - an unusual presentation. *J Assoc Physicians India*, 43(5), 371-372.
- Elliot, E. L., Kaysner, A. C., Tamplin, M. L. (2001). *V. cholerae*, *V. parahaemolyticus*, *V. vulnificus* and other *Vibrio* species In: Bacteriological analytical manual online, Chapter 9 (9th Ed). US FDA Centre for Food Safety and Applied Nutrition. Retrieved from: [http:// vm.cfsan.fda.gov/ebam/bam-toc-html](http://vm.cfsan.fda.gov/ebam/bam-toc-html).
- Faruque, S. M., Albert, M. J., & Mekalanos, J. J. (1998). Epidemiology, genetics, and ecology of toxigenic *Vibrio cholerae*. *Microbiol Mol Biol Rev*, 62(4), 1301-1314.
- Faruque, S. M., Kamruzzaman, M., Meraj, I. M., Chowdhury, N., Nair, G. B., Sack, R. B., *et al.* (2003). Pathogenic potential of environmental *Vibrio cholerae* strains carrying genetic variants of the toxin-coregulated pilus pathogenicity island. *Infect Immun*, 71(2), 1020-1025.
- Faruque, S. M., Chowdhury, N., Kamruzzaman, M., Dziejman, M., Rahman, M. H., Sack, D. A., *et al.* (2004). Genetic diversity and virulence potential of environmental *Vibrio cholerae* population in a cholera-endemic area. *Proc Natl Acad Sci U S A*, 101(7), 2123-2128.
- Faruque, S. M., Islam, M. J., Ahmad, Q. S., Biswas, K., Faruque, A. S., Nair, G. B., *et al.* (2006). An improved technique for isolation of environmental *Vibrio cholerae* with epidemic potential: monitoring the emergence of a multiple-antibiotic-resistant epidemic strain in Bangladesh. *J Infect Dis*, 193(7), 1029-1036.
- Fields, P. I., Popovic, T., Wachsmuth, K., & Olsvik, O. (1992). Use of polymerase chain reaction for detection of toxigenic *Vibrio cholerae* O1 strains from the Latin American cholera epidemic. *J Clin Microbiol*, 30(8), 2118-2121.
- Fullner, K. J., Boucher, J. C., Hanes, M. A., Haines, G. K., 3rd, Meehan, B. M., Walchle, C., *et al.* (2002). The contribution of accessory toxins of *Vibrio cholerae* O1 El Tor to the proinflammatory response in a murine pulmonary cholera model. *J Exp Med*, 195(11), 1455-1462.
- Garg, P., Aydanian, A., Smith, D., J. Glenn M, J., Nair, G. B., & Stine, O. C. (2003). Molecular epidemiology of O139 *Vibrio cholerae*: mutation, lateral gene transfer, and founder flush. *Emerg Infect Dis*, 9(7), 810-814.
- Gonzalez, S. F., Krug, M. J., Nielsen, M. E., Santos, Y., & Call, D. R. (2004). Simultaneous Detection of Marine Fish Pathogens by Using Multiplex PCR and a DNA Microarray. *J. Clin. Microbiol.*, 42(4), 1414-1419.
- Harkey, C. W., Everiss, K. D., & Peterson, K. M. (1994). The *Vibrio cholerae* toxin-coregulated-pilus gene *tcpI* encodes a homolog of methyl-accepting chemotaxis proteins. *Infect Immun*, 62(7), 2669-2678.

- Harwood, V. J., Gandhi, J. P., & Wright, A. C. (2004). Methods for isolation and confirmation of *Vibrio vulnificus* from oysters and environmental sources: a review. *J Microbiol Methods*, 59(3), 301-316.
- Hase, C. C., & Mekalanos, J. J. (1998). TcpP protein is a positive regulator of virulence gene expression in *Vibrio cholerae*. *Proc Natl Acad Sci U S A*, 95(2), 730-734.
- Hazen, T. H., Kennedy, K. D., Chen, S., Yi, S. V., & Sobecky, P. A. (2009). Inactivation of mismatch repair increases the diversity of *Vibrio parahaemolyticus*. *Environ Microbiol*, 11(5), 1254-1266.
- Holmes, B., Willcox, W. R., & Lapage, S. P. (1978). Identification of Enterobacteriaceae by the API 20E system. *J Clin Pathol*, 31(1), 22-30.
- Hinton, J. C., Hautefort, I., Eriksson, S., Thompson, A., & Rhen, M. (2004). Benefits and pitfalls of using microarrays to monitor bacterial gene expression during infection. *Curr Opin Microbiol*, 7(3), 277-282.
- Hoshino, K., Yamasaki, S., Mukhopadhyay, A. K., Chakraborty, S., Basu, A., Bhattacharya, S. K., *et al.* (1998). Development and evaluation of a multiplex PCR assay for rapid detection of toxigenic *Vibrio cholerae* O1 and O139. *FEMS Immunol Med Microbiol*, 20(3), 201-207.
- Hunter, P. R., & Gaston, M. A. (1988). Numerical index of the discriminatory ability of typing systems: an application of Simpson's index of diversity. *J Clin Microbiol*, 26(11), 2465-2466.
- Kaper, J. B., Morris, J. G., Jr., & Levine, M. M. (1995). Cholera. *Clin Microbiol Rev*, 8(1), 48-86.
- Kay, B. A., Bopp, C. A., Wells, J. G. (1994). Isolation and identification of *Vibrio cholerae* O1 from fecal specimens. In: *Vibrio cholerae* and cholera: molecular to global prospective, by Wachsmuth, K., Blake, P. A., Olsvik, O. p 3-25.
- Keymer, D. P., Miller, M. C., Schoolnik, G. K., & Boehm, A. B. (2007). Genomic and phenotypic diversity of coastal *Vibrio cholerae* strains is linked to environmental factors. *Appl Environ Microbiol*, 73(11), 3705-3714.
- Kaysner, C. A., Depaola, A.J. (2004). *Vibrio* bacteriological analytical manual online. U.S. Food and Drug Administration, centre for Food Safety and Applied Nutrition. Retrieved from: <http://www.cfsan.fda.gov/~ebam/bam-9.html>.
- Kita-Tsukamoto, K., Oyaizu, H., Nanba, K., & Simidu, U. (1993). Phylogenetic Relationships of Marine Bacteria, Mainly Members of the Family Vibrionaceae, Determined on the Basis of 16S rRNA Sequences. *Int J Syst Bacteriol*, 43(1), 8-19.

- Klein, D. (2002). Quantification using real-time PCR technology: applications and limitations. *Trends Mol Med*, 8(6), 257-260.
- Kotetishvili, M., Stine, O. C., Chen, Y., Kreger, A., Sulakvelidze, A., Sozhamannan, S., *et al.* (2003). Multilocus sequence typing has better discriminatory ability for typing *Vibrio cholerae* than does pulsed-field gel electrophoresis and provides a measure of phylogenetic relatedness. *J Clin Microbiol*, 41(5), 2191-2196.
- Ravi Kumar, A., Sathish, V., Balakrish Nair, G., & Nagaraju, J. (2007). Genetic characterization of *Vibrio cholerae* strains by inter simple sequence repeat-PCR. *FEMS Microbiol Lett*, 272(2), 251-258.
- Lee, S. H., Hava, D. L., Waldor, M. K., & Camilli, A. (1999). Regulation and temporal expression patterns of *Vibrio cholerae* virulence genes during infection. *Cell*, 99(6), 625-634.
- Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods*, 25(4), 402-408.
- Li, M., Shimada, T., Morris, J. G., Jr., Sulakvelidze, A., & Sozhamannan, S. (2002). Evidence for the emergence of non-O1 and non-O139 *Vibrio cholerae* strains with pathogenic potential by exchange of O-antigen biosynthesis regions. *Infect Immun*, 70(5), 2441-2453.
- Liu, Z., Miyashiro, T., Tsou, A., Hsiao, A., Goulian, M., & Zhu, J. (2008). Mucosal penetration primes *Vibrio cholerae* for host colonization by repressing quorum sensing. *Proc Natl Acad Sci U S A*, 105(28), 9769-9774.
- Maiden, M. C., Bygraves, J. A., Feil, E., Morelli, G., Russell, J. E., Urwin, R., *et al.* (1998). Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc Natl Acad Sci U S A*, 95(6), 3140-3145.
- Mantri, C. K., Mohapatra, S. S., Ramamurthy, T., Ghosh, R., Colwell, R. R., & Singh, D. V. (2006). Septaplex PCR assay for rapid identification of *Vibrio cholerae* including detection of virulence and int SXT genes. *FEMS Microbiol Lett*, 265(2), 208-214.
- Matson, J. S., Withey, J. H., & DiRita, V. J. (2007). Regulatory networks controlling *Vibrio cholerae* virulence gene expression. *Infect Immun*, 75(12), 5542-5549.
- Merrell, D. S., Butler, S. M., Qadri, F., Dolganov, N. A., Alam, A., Cohen, M. B., *et al.* (2002). Host-induced epidemic spread of the cholera bacterium. *Nature*, 417(6889), 642-645.
- Mohapatra, S. S., Ramachandran, D., Mantri, C. K., Colwell, R. R., & Singh, D. V. (2009). Determination of relationships among non-toxicogenic *Vibrio cholerae* O1

biotype El Tor strains from housekeeping gene sequences and ribotype patterns. *Res Microbiol*, 160(1), 57-62.

- Nandi, B., Nandy, R. K., Mukhopadhyay, S., Nair, G. B., Shimada, T., & Ghose, A. C. (2000). Rapid Method for Species-Specific Identification of *Vibrio cholerae* Using Primers Targeted to the Gene of Outer Membrane Protein OmpW. *J. Clin. Microbiol.*, 38(11), 4145-4151.
- Nair, G. B., Shimada, T., Kurazono, H., Okuda, J., Pal, A., Karasawa, T., et al. (1994). Characterization of phenotypic, serological, and toxigenic traits of *Vibrio cholerae* O139 bengal. *J Clin Microbiol*, 32(11), 2775-2779.
- Nair, G. B., Faruque, S. M., Bhuiyan, N. A., Kamruzzaman, M., Siddique, A. K., & Sack, D. A. (2002). New variants of *Vibrio cholerae* O1 biotype El Tor with attributes of the classical biotype from hospitalized patients with acute diarrhea in Bangladesh. *J Clin Microbiol*, 40(9), 3296-3299.
- Nair, G. B., Qadri, F., Holmgren, J., Svennerholm, A. M., Safa, A., Bhuiyan, N. A., et al. (2006). Cholera due to altered El Tor strains of *Vibrio cholerae* O1 in Bangladesh. *J Clin Microbiol*, 44(11), 4211-4213.
- Navia, M. M., Capitano, L., Ruiz, J., Vargas, M., Urassa, H., Schelleberg, D., et al. (1999). Typing and characterization of mechanisms of resistance of *Shigella* spp. isolated from feces of children under 5 years of age from Ifakara, Tanzania. *J Clin Microbiol*, 37(10), 3113-3117.
- Nhung, P. H., Shah, M. M., Ohkusu, K., Noda, M., Hata, H., Sun, X. S., et al. (2007). The *dnaJ* gene as a novel phylogenetic marker for identification of *Vibrio* species. *Syst Appl Microbiol*, 30(4), 309-315.
- Noriega-Orozco, L., Acedo-Felix, E., Higuera-Ciapara, I., Jimenez-Flores, R., Cano, R. (2007). Pathogenic and non-pathogenic *Vibrio* species in aquaculture shrimp ponds. *Rev Latinoam Microbiol*, 49, 60-67.
- Nucera, D. M., Maddox, C. W., Hoiem-Dalen, P., & Weigel, R. M. (2006). Comparison of API 20E and *invA* PCR for identification of *Salmonella enterica* isolates from swine production units. *J Clin Microbiol*, 44(9), 3388-3390.
- O'Hara, C. M., Sowers, E. G., Bopp, C. A., Duda, S. B., & Strockbine, N. A. (2003). Accuracy of six commercially available systems for identification of members of the family vibriaceae. *J Clin Microbiol*, 41(12), 5654-5659.
- Olivier, V., Salzman, N. H., & Satchell, K. J. (2007). Prolonged colonization of mice by *Vibrio cholerae* El Tor O1 depends on accessory toxins. *Infect Immun*, 75(10), 5043-5051.
- Olsen, J. S., Aarskaug, T., Skogan, G., Fykse, E. M., Ellingsen, A. B., & Blatny, J. M. (2009). Evaluation of a highly discriminating multiplex multi-locus

variable-number of tandem-repeats (MLVA) analysis for *Vibrio cholerae*. *J Microbiol Methods*, 78(3), 271-285.

- Olsvik, O., Wahlberg, J., Petterson, B., Uhlen, M., Popovic, T., Wachsmuth, I. K., et al. (1993). Use of automated sequencing of polymerase chain reaction-generated amplicons to identify three types of cholera toxin subunit B in *Vibrio cholerae* O1 strains. *J Clin Microbiol*, 31(1), 22-25.
- Overman, T. L., Kessler, J. F., & Seabolt, J. P. (1985). Comparison of API 20E, API rapid E, and API rapid NFT for identification of members of the family Vibrionaceae. *J Clin Microbiol*, 22(5), 778-781.
- Ottaviani, D., Leoni, F., Rocchegiani, E., Santarelli, S., Masini, L., Di Trani, V., et al. (2009). Prevalence and virulence properties of non-O1 non-O139 *Vibrio cholerae* strains from seafood and clinical samples collected in Italy. *Int J Food Microbiol*, 132(1), 47-53.
- Panicker, G., Call, D. R., Krug, M. J., & Bej, A. K. (2004). Detection of Pathogenic *Vibrio* spp. in Shellfish by Using Multiplex PCR and DNA Microarrays. *Appl Environ Microbiol*, 70(12), 7436-7444.
- Di Pinto, A., Ciccarese, G., Tantillo, G., Catalano, D., & Forte, V. T. (2005). A collagenase-targeted multiplex PCR assay for identification of *Vibrio alginolyticus*, *Vibrio cholerae*, and *Vibrio parahaemolyticus*. *J Food Prot*, 68(1), 150-153.
- Radu, S., Vincent, M., Apun, K., Abdul-Rahim, R., Benjamin, P. G., Yuherman, et al. (2002). Molecular characterization of *Vibrio cholerae* O1 outbreak strains in Miri, Sarawak (Malaysia). *Acta Trop*, 83(2), 169-176.
- Rivera, I. N., Chun, J., Huq, A., Sack, R. B., & Colwell, R. R. (2001). Genotypes associated with virulence in environmental isolates of *Vibrio cholerae*. *Appl Environ Microbiol*, 67(6), 2421-2429.
- Rivera, I. G., Chowdhury, M. A., Huq, A., Jacobs, D., Martins, M. T., & Colwell, R. R. (1995). Enterobacterial repetitive intergenic consensus sequences and the PCR to generate fingerprints of genomic DNAs from *Vibrio cholerae* O1, O139, and non-O1 strains. *Appl Environ Microbiol*, 61(8), 2898-2904.
- Ruimy, R., Breittmayer, V., Elbaze, P., Lafay, B., Boussemart, O., Gauthier, M., et al. (1994). Phylogenetic Analysis and Assessment of the Genera *Vibrio*, *Photobacterium*, *Aeromonas*, and *Plesiomonas* Deduced from Small-Subunit rRNA Sequences. *Int J Syst Bacteriol*, 44(3), 416-426.
- Le Roux, F., Gay, M., Lambert, C., Nicolas, J. L., Gouy, M., & Berthe, F. (2004). Phylogenetic study and identification of *Vibrio splendidus*-related strains based on *gyrB* gene sequences. *Dis Aquat Organ*, 58(2-3), 143-150.

- Rozen, S., & Skaletsky, H. J. (2000). Primer 3 on the WWW for general users and for biologist programmers,” In: Krawetz S, Misener S (Eds), Bioinformatics methods and protocols: methods in molecular biology. Humana Press, Totowa, N.J., pp. 223-233.
- Rader, A. E., & Murphy, J. R. (1988). Nucleotide sequences and comparison of the hemolysin determinants of *Vibrio cholerae* El Tor RV79(Hly+) and RV79(Hly-) and classical 569B(Hly-). *Infect Immun*, 56(6), 1414-1419.
- Rahman, M. H., Biswas, K., Hossain, M. A., Sack, R. B., Mekalanos, J. J., & Faruque, S. M. (2008). Distribution of genes for virulence and ecological fitness among diverse *Vibrio cholerae* population in a cholera endemic area: tracking the evolution of pathogenic strains. *DNA Cell Biol*, 27(7), 347-355.
- Reidl, J., & Klose, K. E. (2002). *Vibrio cholerae* and cholera: out of the water and into the host. *FEMS Microbiol Rev*, 26(2), 125-139.
- Raychoudhuri, A., Patra, T., Ghosh, K., Ramamurthy, T., Nandy, R. K., Takeda, Y., et al. (2009). Classical ctxB in *Vibrio cholerae* O1, Kolkata, India. *Emerg Infect Dis*, 15(1), 131-132.
- Sack, D. A., Lyke, C., McLaughlin, C., & Suwanvanichkij, V. (2001). Antimicrobial resistance in shigellosis, cholera and campylobacteriosis. *World Health Org* ,8, 8-20.
- Safa, A., Bhuyian, N. A., Nusrin, S., Ansaruzzaman, M., Alam, M., Hamabata, T., et al. (2006). Genetic characteristics of Matlab variants of *Vibrio cholerae* O1 that are hybrids between classical and El Tor biotypes. *J Med Microbiol*, 55(11), 1563-1569.
- Salim, A., Lan, R., & Reeves, P. R. (2005). *Vibrio cholerae* pathogenic clones. *Emerg Infect Dis*, 11(11), 1758-1760.
- [Salyers](#), A. A. & [Whitt](#), D. D. (2002). Bacterial pathogenesis: A molecular approach. ASM Press, Wahington,D.C.
- Shelburne, S. A., & Musser, J. M. (2004). Virulence gene expression *in vivo*. *Curr Opin Microbiol*, 7(3), 283-289.
- Singh, D. V., Matte, M. H., Matte, G. R., Jiang, S., Sabeena, F., Shukla, B. N., et al. (2001). Molecular analysis of *Vibrio cholerae* O1, O139, non-O1, and non-O139 strains: clonal relationships between clinical and environmental isolates. *Appl Environ Microbiol*, 67(2), 910-921.
- Steffen, R., Acar, J., Walker, E., & Zuckerman, J. (2003). Cholera: assessing the risk to travellers and identifying methods of protection. *Travel Med Infect Dis*, 1(2), 80-88.

- Syed, K. A., Beyhan, S., Correa, N., Queen, J., Liu, J., Peng, F., *et al.* (2009). The *Vibrio cholerae* flagellar regulatory hierarchy controls expression of virulence factors. *J Bacteriol*, 191(21), 6555-6570.
- Tamura, K., Dudley, J., Nei, M., & Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol*, 24(8), 1596-1599.
- Tarr, C. L., Patel, J. S., Puhr, N. D., Sowers, E. G., Bopp, C. A., & Strockbine, N. A. (2007). Identification of *Vibrio* isolates by a multiplex PCR assay and *rpoB* sequence determination. *J Clin Microbiol*, 45(1), 134-140.
- Teh, C. S., Chua, K. H., Puthuchery, S. D., & Thong, K. L. (2008). Further evaluation of a multiplex PCR for differentiation of *Salmonella* paratyphi A from other salmonellae. *Jpn J Infect Dis*, 61(4), 313-314.
- Thompson, F. L., Iida, T., & Swings, J. (2004). Biodiversity of vibrios. *Microbiol Mol Biol Rev*, 68(3), 403-431, table of contents.
- Thompson, F. L., Gevers, D., Thompson, C. C., Dawyndt, P., Naser, S., Hoste, B., *et al.* (2005). Phylogeny and molecular identification of vibrios on the basis of multilocus sequence analysis. *Appl Environ Microbiol*, 71(9), 5107-5115.
- Thong, K. L., Goh, Y. L., Cam, P. C. (2002). Molecular subtyping of clinical isolates *Vibrio cholerae* from Malaysia and Vietnam by pulsed-field gel electrophoresis. *Asia Pac J Mol Biol Biotechnol*, 10, 103-108.
- Tison, D. L. (1999). *Vibrio*. Bergeys Manual of Microbiology. 17th Edition. In Murray PR (ed). Washington DC, ASM Press. 497-506
- Tokunaga, A., Yamaguchi, H., Morita, M., Arakawa, E., Izumiya, H., Watanabe, H., *et al.* (2010). Novel PCR-based genotyping method, using genomic variability between repetitive sequences of toxigenic *Vibrio cholerae* O1 El Tor and O139. *Mol Cell Probes*, 24(2), 99-103.
- Tracz, D. M., Backhouse, P. G., Olson, A. B., McCrea, J. K., Walsh, J. A., Ng, L. K., *et al.* (2007). Rapid detection of *Vibrio* species using liquid microsphere arrays and real-time PCR targeting the *ftsZ* locus. *J Med Microbiol*, 56(Pt 1), 56-65.
- Uma, G., Chandrasekaran, M., Takeda, Y., & Nair, G. B. (2003). Recent advances in cholera genetics. *Current Science*, 85(11), 1538—1545.
- World Health Organization. (2002) Cholera annual report 2001. *Weekly Epidemiological Record*, 77(31), 257-268.
- Wakeley, J. (1996). The variance of pairwise nucleotide differences in two populations with migration. *Theor Popul Biol*, 49(1), 39-57.

- Vadivelu, J., Iyer, L., Kshatriya, B. M., & Puthucheary, S. D. (2000). Molecular evidence of clonality amongst *Vibrio cholerae* O1 biotype El Tor during an outbreak in Malaysia. *Epidemiol Infect*, *124*(1), 25-30.
- van Belkum, A., Struelens, M., de Visser, A., Verbrugh, H., & Tibayrenc, M. (2001). Role of genomic typing in taxonomy, evolutionary genetics, and microbial epidemiology. *Clin Microbiol Rev*, *14*(3), 547-560.
- van Belkum, A. (2007). Tracing isolates of bacterial species by multilocus variable number of tandem repeat analysis (MLVA). *FEMS Immunol Med Microbiol*, *49*(1), 22-27.
- van Belkum, A., Tassios, P. T., Dijkshoorn, L., Haeggman, S., Cookson, B., Fry, N. K., *et al.* (2007). Guidelines for the validation and application of typing methods for use in bacterial epidemiology. *Clin Microbiol Infect*, *13 Suppl 3*, 1-46.
- Versalovic, J., Koeuth, T., Lupski, J. R. (1991). Distribution of repetitive DNA sequences in eubacteria and applications to fingerprinting of bacterial genomes. *Nucleic Acids Res*, *19*, 6823-6831.
- Virdi, J. S. & Sachdeva, P. (2005). Genetic diversity of pathogenic microorganisms: Basic insights, public health implications and the Indian initiatives. *Current Science*, *89*(1), 113-123.
- Xu, Q., Dziejman, M., & Mekalanos, J. J. (2003). Determination of the transcriptome of *Vibrio cholerae* during intrainestinal growth and midexponential phase in vitro. *Proc Natl Acad Sci U S A*, *100*(3), 1286-1291.
- Zhang, W., Jayarao, B. M., & Knabel, S. J. (2004). Multi-virulence-locus sequence typing of *Listeria monocytogenes*. *Appl Environ Microbiol*, *70*(2), 913-920.
- Zulkifli, Y., Alitheen, N. B., Son, R., Raha, A.R., Samuel, L., Yeap, S.K., *et al.* (2009) Random amplified polymorphic DNA-PCR and ERIC PCR analysis on *Vibrio parahaemolyticus* isolated from cockles in Padang, Indonesia. *Int Food Res J* *16*, 141-150.