

## REFERENCES

- Abbott, I. A., Fisher, J. & McDermid, K. J. (2002). New reported and revised marine algae from the vicinity of Nha Trang, Vietnam. In: *Taxonomy of Economic Seaweeds with reference to some Pacific species. Vol. VIII.* (Abbott, I.A. & Mcdermid, K.J. Eds) Vol.8, pp. 291-321.
- Agardh, C. A., (1817). Synopsis algarum Scandinaviae, adjecta dispositione universali algarum. *Lund.* 1: 135.
- Agardh, C. A., (1823). Species algarum rite cognitae cum synonymis, differentis specificis et descriptionibus succinctis. *Lund.* 1: 169-521.
- Agardh, C. A., (1849). Caulerpearum sciagraphia. *Linnaea*, 22: 129-144.
- Agardh, J. G. (1872). Till algernes systematik. I. Acta Univ. *Lund.* 9:1–136.
- Agardh, J.G. (1837). Novae species algarum, quas in itinere ad oras maris rubri collegit Eduardus Rüppell; cum observationibus nonnullis in species rariores antea cognitas. *Museum Senckenbergianum* 2: 169-174.
- Aleem, A. A. (1948). The recent migration of certain Indopacific algae from the Red Sea into the Mediterranean. *New Phytol.* 47: 88–94.
- Amico, V., Oriente, G., Piatelli. M., Tringali, C.. Fattorusso, E., Magno. S., & *et al.* (1978). Caulerpenyne, an unusual sesquiterpenoid from the green alga *Caulerpa prolifera*. *Tetrahedron Lett.* 3593-3596.
- Andrade, L. R., Salgado, L. T., Farina, M., Pereira, M. S., Mourão, P. A. S., & Amado Filho, G. M. (2004). Ultrastructure of acidic polysaccharides from the cell walls of brown algae. *J. Struct. Biol.* 145: 216–25.
- Arasaki, S., & Arasaki, T. (1983). Low Calorie, High Nutrition Vegetables from the Sea to Help You Look and Feel Better. Japan Publications, Tokyo, 196 pp.
- Argyrou, M., Demetropoulos, A., & Hadjichristophorou, M. (1999). Expansion of the macroalga *Caulerpa racemosa* and changes in softbottom macrofaunal assemblages in Moni Bay. *Cyprus. Oceanol. Acta* 22 (5): 517– 528.
- Ayyad, S. E. N., & Badria, F.A. (1994). Caulerpin, an antitumor indole alkaloid from *Caulerpa racemosa*, Alex. *J. Pharm. Sci.* 8: 217.
- Baba, M., Snoeck, R., Pauwels, R., & De Clercq, E. (1988). *Antimicrob. Agents Chemother.* 32: 1742.

- Baldauf, S. L., Manhart, J. R., & Palmer, J. D. (1990). Different fates of the chloroplast *tufA* gene following the transfer to the nucleus in green algae. *Proc. Natl. Acad. Sci. USA* 87: 5317-5321.
- Baldauf, S. L., Palmer, J. D., & Doolittle, W. F. (1996). The root of the universal tree and the origin of eukaryotes based on elongation factor phylogeny. *Proc. Natl. Acad. Sci. USA* 93:7749–54.
- Barbier, P., Guise, S., Huitorel, P., Amade, P., Pesando, D., Briand, C., & Peyrot, V. (2001). Caulerpenyne from *Caulerpa taxifolia* has an antiproliferative activity on tumor cell line SK-N-SH and modifies the microtubule network. *Life. Sci.* 70: 415–429.
- Belsher, T., & Meinesz, A. (1995). Deep-water dispersion of the alga *Caulerpa taxifolia* introduced into the Mediterranean. *Aquat. Bot.* 51: 163–169.
- Benzie, J. A. H., Price, I. R., & Ballment, E. (1997). Population genetics and taxonomy of *Caulerpa* (Chlorophyta) from the Great Barrier Reef, Australia. *J. Phycol.* 33, 491–504.
- Bhattacharya, D., & Druehl, L. D. (1990). Restriction enzyme analysis of variation and taxonomy in the kelp genus *Laminaria* (Laminariales, Phaeophyta), *Hydrobiologia*, 204-205 (1): 105-110.
- Bird, K. T., Chynoweth, D. P., & Jerger, D. E. (1990). Effects of marine algal proximate composition on methane yields. *J. Appl. Phycol.* 2: 207-213.
- Biren, K., Young, J., Buttenshaw, R., Searle, J., Cooksley, G., Xu, D. B., & et al. (1997). Frequency of mutation and deletion of the tumor suppressor gene CDKN2A (MTS1/p16) in hepatocellular carcinoma from an Australian population. *Hepatology* 25: 593–597.
- Blunden, G. (1977). Cytokinin activity of seaweed extracts. In Faulkner DL, Fenical WH (eds), *Marine Natural Products Chemistry*. Plenum Publishing Corporation, New York, 337-344.
- Bold, H. C., & Wynne, M. J. (1985). *Introduction to the Algae. Structure and Reproduction.* 2nd ed. Prentice Hall, Englewood Cliffs, NJ, 1706 pp.
- Bonny, C., & Stutz, E. (1993). Soybean (*Glycine max* L.) nuclear DNA contains four *tufA* genes coding for the chloroplastspecific translation elongation factor EF-Tu. *Chimia* 47: 247-249.
- BØrgesen, F. (1907). An ecological and systematic account of the Caulerpas of the Danish West Indies. K.danske Vidensk. Selsk. SKR., Ser VII. Naturvid. *Og Math. Afd.*, 4: 337-392.
- BØrgesen, F. (1913). The marine algae of the Danish West Indies. Part 1. Chlorophyceae. *Dansk Botanisk Arkiv* 1(4): 1–158.

- Botstein, D., Raymond, L. W., Skolnick, M., & Davis, R. W. (1980). Construction of a Genetic Linkage Map in Man Using Restriction Fragment Length Polymorphisms. *Am. J. Hum. Genet.* 32: 314-331.
- Brodie, J., & Lewis, J. (2007). Unravelling the Algae: the Past, Present and Future of Algal Systematics. CRC Press, Boca Raton, USA.
- Brown, J. R., Douady, C. J., Italia, M. J., Marshall, W. E., & Stanhope, M. J. (2001). Universal trees based on large combined protein sequence data sets. *Nature Genet.* 28, 281–285.
- Buschmann, A. H., Mora, O. A., Gómez, P., Böttger, M., Buitano, S., Retamales, C., & et al. (1994). *Gracilaria* tank cultivation in Chile: use of land based salmon culture effluents. *Aquacult. Eng.* 13: 283–300.
- Calvert, H. (1976). Culture studies on some Florida species of *Caulerpa*: morphological responses to reduced illumination. *Br. Phycol. J.* 11:203–14.
- Calvert, H. E., Dawes, C. J., & Borowitzka, M. A. (1976). Phylogenetic relationships of *Caulerpa* (Chlorophyta) based on comparative chloroplast ultrastructure. *J. Phycol.* 12: 149-162.
- Carruthers, T. J. B., Walker, D. I., & Huisman, J. M. (1993). Culture studies on two morphological types of *Caulerpa* (Chlorophyta) from Perth, Western Australia, with a description of a new species. *Bot. Mar.* 36: 589–96.
- Ceccherelli, G., & Cinelli, F. (1999a). The role of vegetative spreading in dispersal of the invasive alga *Caulerpa taxifolia* in the Mediterranean. *Mar. Ecol. Prog. Ser.* 182: 299–303.
- Chamberlain, A. (1998). Culture and shelf-life extension of *Caulerpa racemosa*. *S. Pac. J. Nat. Sci.* 16: 11–19.
- Chan, H. T. (2003). Use of morphological and molecular analyses to understand the relationship between *Gracilaria tenuistipitata* and its closely related species. Bsc. Thesis, University of Malaya, Kuala Lumpur.
- Chynoweth, D. P., Kerby, F., & Vipul, J. (1987). Biological gasification of marine algae. University of Florida, Gainesville, FL, 32611 and Institute of Gas Technology, Chicago, IL, 60616, U. S. A.
- Clavijo, R. E., Weiss-Lopez, B., & Rivera, P. (1996). Vibrational spectra of Caulerpin, a green algae pigment. *Bol. Soc. Chil. Quim.* 41: 153.

- Cohen, I., & Neori, A. (1991). *Ulva lactuca* biofilters for marine fish pond effluent. I. Ammonia uptake kinetics and nitrogen content. *Bot. mar.* 34: 475–482.
- Coleman, A. W., & Goff, L. J. (1988). The use of plastid DNA restriction endonuclease patterns in delineating red algal species and populations. *J. Phycol.* 24: 357–368.
- Coleman, A. W., & Goff, L. J. (1991). DNA analysis of Eukaryotic algal species. *J. Phycol.* 27: 463-473.
- Collado-Vides, L. (2002). Morphological plasticity of *Caulerpa prolifera* (Caulerpales, Chlorophyta) in relation to growth form in a coral reef lagoon. *Bot. Mar.* 45: 123–9.
- Collado-Vides, L., & Robledo, D. (1999). Morphology and photosynthesis of *Caulerpa* (Chlorophyta) in relation to growth form. *J. Phycol.* 35: 325–30.
- Coppejans, E. (1992). Marine algae of Papua New Guinea (Madang Prov.). 2. A revised and completed list of *Caulerpa* (Chlorophyta- Caulerpales). *Blumea* 35: 383–410.
- Coppejans, E., & Beeckman, T. (1990). *Caulerpa* (Chlorophyta, Caulerpales) from the Kenyan coast. *J. Nov. Hedw.* 49: 381-393.
- Coppejans, E., & Meinesz, A. (1988). Marine algae of Papua New Guinea (Madang Prov.) Caulerpaceae (Chlorophyta-Caulerpales). *Blumea* 33:181–96.
- Coppejans, E., & Prud'homme van Reine, W. F. (1992). Seaweeds of the Snellius-II Expedition (E. Indonesia): the genus *Caulerpa* (Chlorophyta- Caulerpales). *Bull. Séanc. Acad. Sci. Outre-Mer.* 37: 667–712.
- Coppejans, E., & Van Den Heede, C. (1996). The Codiales (excluding *Codium*) (Chlorophyta) from Kenya, Tanzania and the Seychelles. *Belg. J. Bot.* 129 (1): 47–65.
- COST 48. (1987). Aquatic Primary Biomass (Marine Macroalgae): Biomass conversion, removal and use of nutrients. I. Proceedings of the First Workshop of subgroup 3. Crema- L'Houmeau, CNRS-IFREMER, L'Homeau, France. Commission of European Communities, Brussels, p. 140.
- Cribb, A.B. (1996). Seaweeds of Queensland. A naturalist's guide. pp. 130. Brisbane: The Queensland Naturalists' Club Inc.
- Dalton, R. (2000). Researchers criticize response to killer algae. *Nature* 406: 447.
- De senerpont domis, L. N., Fama, P., Bartlett, A. J., Prud'homme reine, W. F., Espinosa C. A., & Trono, G.C. (2003). Defining taxon boundaries in members of the morphologically and genetically plastic genus *Caulerpa* (Caulerpales, Chlorophyta). *J. Phycol.* 39: 1019-1037.

- Delwiche, C. F., Kuhsel, M., & Palmer, J. D. (1995). Phylogenetic analysis of *tufA* sequences indicates a cyanobacterial origin of all plastids. *Mol. Phyl. Evol.* 4:110–28.
- Dharmananda, S. (2007). The nutritional and medicinal value of seaweeds used in Chinese medicine. itmonline.org Available from: <http://www.itmonline.org/arts/seaweed.htm>
- Djellouli, A., Langar, H., & El abed, A. (1998). *Caulerpa racemosa* (Forsskaô l) J. Agardh: bio-geographie et écologie. In: Third International Workshop on *Caulerpa Taxifolia* (C. F. Boudouresque, V. Gravez, A. Meinesz & F. Palluy, eds), pp. 233-238. GIS Posidonie Publ., Marseilles.
- Domis, L., Fama, P., Bartlett, A. J., Prud'homme van Reine, W. F., Espinosa, C. A., & Trono, G. C. (2003). Defining taxon boundaries in members of the morphologically and genetically plastic genus *Caulerpa* (Caulerpales, Chlorophyta). *J. Phycol.* 39: 1019–1037.
- Donoghue, M. J., & Sanderson, M. J. (1992). The suitability of molecular and morphological evidence in reconstructing plant phylogeny. 340- 368.
- Doty, M., & Aguilar-Santos, G. (1970). Transfer of toxic algal substances in marine food. *Pac. Sci.* 24: 351-355.
- Durand, C., Manuel, M., Boudouresque, C. F., Mienesz, A., Verlaque, M. Le Parco, M., & et al. (2002). Molecular data suggest a hybrid origin for the invasive *Caulerpa racemosa* (Caulerpales, Chlorophyta) in the Mediterranean Sea. *J. Evol. Biol.* 15: 122-133.
- Dutcher, J. A., & Kapraun, D. F. (1994). Random amplified polymorphic DNA (RAPD) identification of genetic variation in three species of *Porphyra* (Bangiales, Rhodophyta). *Appl. Phycol.* 6: 267-273.
- Eubank, L. L. (1946). Hawaiian representatives of the genus *Caulerpa*. *Univ. Publ. Bot.* 18: 409- 432.
- Eyras, M. C., Rostagno, C. M., & Defossé, G.E. (1998). Biological evaluation of seaweed composting. *Compost Sci. Util.* 6: 74–81.
- Fama, P., Wysor, B., Kooistra, W. H., & Zuccarello, G. C., (2002). Molecular phycology of the genus *Caulerpa* (Caulerpales, Chlorophyta) inferred from chloroplast *tufA* gene. *J. phycol.* 38: 1040-1050.
- Fischel, J. L., Lem'ee, R., Formento, P., Caldani, C., Moll, J. L., Pesando, D., & et al. (1995). Cell growth inhibitory effects of caulerpenyne, a sesquiterpenoid from the marine alga *Caulerpa taxifolia*. *Anticancer Res.* 15: 2155–2160.

- Fleurence, J. (1999). Seaweed proteins: biochemical, nutritional aspects and potential uses. *Trends Food Sci. Tech.* 10: 25-28.
- Flores-Nava, A. (1995). Some considerations on sustainable shrimp mariculture. *Jaina* 6: 8.
- Flowers, A. B., & Bird, K. T. (1990). Methane production from seaweeds. In Akatsuka I (ed.), *Introduction to Applied Phycology*. SPB Academic Publishing by, The Hague, 575-587.
- Forsskål, P. (1775). *Flora Aegyptiaca-Arabica sive descriptiones plantarum, quas per Aegyptum inferiorem et Arabium delicem detexit illustravit Petrus Forskål. Prof. Haun. Post mortem auctoris edidit Carsten Niebuhr.* pp. [1]-32, [i]- xxxvi, ... [1]-219, [220, err.], map. Hauniæ
- Gacia, E., Littler, M. M., & Littler, D.S., (1996). The relationships between morphology and photosynthetic parameters within the polymorphic genus *Caulerpa*. *J. Exp. Mar. Biol. Ecol.* 204: 209–224.
- Gambi, M. C., & Terlizzi, A. (1998). Record of large population of *Caulerpa racemosa* (Forsska<sup>o</sup>l) J. Agardh (Chlorophyceae) in the Gulf of Salerno (Southern Tyrrhenian Sea, Italy). *Biol. Mar. Mediterr.* 5 (1): 553– 556.
- Gao, K., & McKinley, K. (1994). Use of macroalgae for marine biomass production and CO<sub>2</sub> remediation: a review. *J. Appl. Phycol.* 6: 45–60.
- Gilbert, W. J. (1941). Notes on *Caulerpa* from Java and the Philippines. *Pap. Mich. Acad. Sci. Arts. Lett.* 27: 7–26.
- Goff, L. J., & Coleman, A. W. (1988). The use of plastid DNA restriction endonuclease patterns in delineating red algal species and populations. *J. Phycol.* 24: 357–368.
- Goncalves, A. G., Ducatti, D. R., Duarte, M. E., & Noseada, M. D. (2002). Sulfated and pyruvylated disaccharide alditoles obtained from a red seaweed galactan: ESIMS and NMR approaches. *Carbohydr. Res.* 337: 2443–2453.
- González, M. A., Montoya, R., Candia, A., Gómez, P., & Cisternas, M. (1996). Organellar DNA restriction fragment length polymorphism (RFLP) and nuclear random amplified polymorphic DNA (RAPD) analyses of morphotypes of *Gracilaria* (Gracilariales, Rhodophyta) from Chile. *Hydrobiologia* 326/327: 229–234.
- Goodwin, T. W. (1974). Carotenoids and biliproteins. In WDP Steward, ed, *Algal Physiology and Biochemistry*. Blackwell, Oxford, pp: 176-205.
- Graham, L. E., & Wilcox, L. W. (2000). *Algae*. Upper Saddle River, NJ: Prentice Hall.

- Haglund, K., & Pedersén, M. (1993). Outdoor pond cultivation of the subtropical marine red alga *Gracilaria tenuistipitata* in brackish water in Sweden. Growth, nutrient uptake, co-cultivation with rainbow trout and epiphyte control. *J. Appl. Phycol.* 5: 271–284.
- Haines, K. C. (1975). Growth of the carrageenan-producing tropical red seaweed *Hypnea musciformis* in surface water, 870 m deep water effluent from a clam mariculture system, and in deep water enriched with artificial fertilizers or domestic sewage. In Persson G, Jaspers E (eds), Proc. 10th Eur. Symp. Mar. Biol., Ostend, Belgium, Sep. 17–23, 1975. Universa Press, Wetteren. 1: 207–220.
- Hall, T. A. (1999). BioEdit: a user friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids Symp. Series*. 41: 95-98.
- Hamel, G. (1926). Quelques algues rares ou nouvelles pour la flore méditerranéenne. *Bull. Mus. Natl. Hist. Nat. Paris* 32 (6): 420.
- Hamel, G. (1930). Les caulerpes méditerranées. *Rev. Algol.* 5: 229.
- Hamel, G. (1931a). Chlorophyceæ de France (suite). *Rev. Algol.* 5: 383-430.
- Hardy, S., Kitamura, M., Harris-Stansil, T., Dai, Y., & Phipps, M. L. (1997). Construction of adenovirus vectors through Cre-lox recombination. *J. Virol.* 71: 1842–1849.
- Harlin, M. M., Thorne-Miller, B., & Thursby, G.H. (1979). Ammonium uptake by *Gracilaria* sp. (Florideophyceae) and *Ulva lactuca* (Chlorophyceae) in closed system fish culture. *Proc. int. Seaweed Symp.* 9: 258–292.
- Haslin, C., Lahaye, M., Pellegrini, M., & Chermann, J. C. (2001). In vitro anti-HIV activity of sulfated cell-wall polysaccharides from gametic, carposporic and tetrasporic stages of the Mediterranean red alga *Asparagopsis armata*. *Planta Med.* 67: 301–305.
- Hillis, D. M., Mable, B. K., & Moritz, C. (1996). Applications of molecular systematics: the state of the field and a look to the future. In Hillis, D. M., Montz, C. and Mable, B. K. (eds.). Molecular Systematics, Sinauer Associates, Inc., Sunderland.
- Hillis, D. M., Moritz, C., & Mable, B. K. (1990). Molecular Systematics. Sinauer Associates, Inc. Publishers. Sunderland, Massachusetts U.S.A.
- Ho, C. L., Phang, S. M., & Pang, T. (1995a). Molecular characterization of *Sargassum polycystum* and *S. siliquosum* (Phaeophyta) by polymerase chain reaction (PCR) using random amplified polymorphic DNA (RAPD) primers. *J. Appl. Phycol.* 7: 33–41.

- Ho, C. L., Phang, S. M., & Pang, T. (1995b). Application of polymerase chain reaction (PCR) using random amplified polymorphic DNA (RAPD) primers in the molecular identification of selected *Sargassum* species (Phaeophyta, Fucales). *Eur. J. Phyol.* 30: 273–280.
- Hodgson, L. M. Pham Huu Tri, Lewmanomont, K. & McDermid, K.J. (2004). Annotated checklist of species of *Caulerpa* and *Caulerpella* (Bryopsidales, Caulerpaceae) from Vietnam, Thailand and the Hawaiian Islands. In: *Taxonomy of Economic Seaweeds with reference to the Pacific and other locations*. 9: 21-38.
- Hodgson, L. M. (1984). Antimicrobial and antineoplastic activity in some South Florida seaweeds. *Bot. Mar.* 27: 387–390.
- Hoek, C., Mann, D. G., & Jahns, H. M. (1995). Algae, an introduction to phycology. Cambridge University Press. pp. 15-17.
- Holdsworth, R. H. (1971). The isolation and partial characterization of the pyrenoid protein of *Eremosphaera viridis*. *J. Cell. Biol.* 51: 499-513.
- Hoppe, H. A. (1979). Marine algae and their products and constituents in pharmacy. In: Marine Algae in Pharmaceutical Science, edited by H. A. Hoppe, T. Levring, and Y. Tanaka. Walter de Gruyter, Berlin: 25-119.
- Huelsenbeck, J. P., Ronquist, F., Nielsen, R., & Bollback, J. P. (2001). Bayesian inference of phylogeny and its impact on evolutionary biology. *J. Science* 294: 2310–2314.
- Indergaard, M. (1983). The aquatic resource. In Cote WA (ed.), Biomass Utilization. Plenum Press, New York: 137-168.
- Ismail, A. (1995). Rumpai Laut Malaysia. Kuala Lumpur, Dewan Bahasa & Pustaka.
- Janson , S., & Hayes , P. K. ( 2006 ). Molecular taxonomy of harmful algae . In *Ecological Studies*, Vol. 189. Exology of Harmful Algae. Edited by Granéil, E., & Turner, J.T. pp. 9 – 21. Springer-Verlag , Berlin.
- Jiménez del Rio, M., Ramazanov, Z., & García-Reina, G. (1994). Optimizationof yield and biofiltering efficiencies of *Ulva rigida* C. Ag. cultivated with *Sparus aurata* L. waste waters. *Sci. Mar.* 58: 329–335.
- John, M. E. (1992). An efficient method for isolation of RNA and DNA from plants containing polyphenolics. *Nucleic Acids Res.* 20: 2381.
- Jousson, O., Pawłowski, J., Zaninetti, L., Meinesz, A., & Boudouresque, C. F. (1998). Molecular evidence for the aquarium origin of the green alga *Caulerpa taxifolia* introduced to the Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 172: 275–280.

Kahl, G., & Winter, P. (1995). Molecular marker technologies for. Plant improvement. *World. J. Microbiol. Biotechnol.* 11: 438-448.

Kain, J. M., & Destombe, C. (1995). A review of the life history, reproductionnand phenology of *Gracilaria* and *Gracilaropsis*. *J. Appl. Phycol.* 7: 269-281.

Kaiser, J. (2000). California algae may be feared European species. *J. Science* 289: 222–223.

Karina, H. M., Cardozo, M. P., Barros, V. R., Falcão, A. P., Tonon, N. P., Lopes, S. C., & et al. (2006). Metabolites from algae with economical impact. *Comp. Biochem. Phys. C.* 146 (2007) 60–78.

Karp, A., Seberg, O., & Buiatti, M. (1996). Molecular Techniques in the Assessment of Botanical Diversity. *Ann. Bot.* 78: 143-149.

Keeton, W. T., & Gould, J. L. (1986). Biological science 4th ed.W. W. Norton & Company. New York.

Kerby, N. W., & Evans, L. V. (1987). Isolation and partial characterization of pyrenoids from the brown alga *pillayella littoralis*. (L). Kjellm. *Planta*. 142: 91-95.

King, R., & Stansfield, W. D. (1985). A dictionary of genetics, third edition. Oxford university press.

Kraft, G.T. (2007). Algae of Australia. Marine benthic algae of Lord Howe Island and the southern Great Barrier Reef, 1. Green algae. pp. [i-iv], v-vi, 1-347, 110 text-figs; 11 pls. Canberra & Melbourne: Australian Biological Resources Study & CSIRO Publishing.

Krom, M. D, Ellner, S., van Rijn, J., & Neori, A. (1995) Nitrogen and phosphorus cycling and transformations in a prototype ‘nonpolluting’ integrated mariculture system, Eilat, Israel. *Mar. Ecol. Prog. Ser.* 118: 25–36.

Kuchitsu, K., Tsuzuki, M., & Miyachi, S. (1988). Characterization of the pyrenoid isolated from unicellular green alga *Chlamydomonas reinhardtii*: particulate form of Rubisco protein. *J. Protoplasma*. 144: 17-24.

Kützing, F. T. (1849). Species algarum. F. A. Brockhaus, Leipzig, 922 pp.

Lamouroux, J. V. F. (1809). Memoires sur les *Caulerpas*. Nouveau genre de la famille des algues marines. *J. Bot. (Paris)* 2: 136–42.

Langton, R. W., Haines, K. C., & Lyon, R. E. (1977). Ammonia-nitrogen production by the bivalve mollusc *Tapes japonica* and its recovery by the red seaweed *Hypnea musciformis* in a tropical mariculture system. *Helgoländer wiss. Meeresun.* 30: 217–229.

Lee, R. E. (1999). Phycology. Third ed. Cambridge University Press, New York, pp. 230.

Leme'e, R., Pesando, D., Issanchou, C., & Amade, P. (1997). Microalgae: a model to investigate the ecotoxicology of the green alga *Caulerpa taxifolia* from the Mediterranean Sea. *J. Mar. Environ. Res.* Lond. 44: 13–25.

Lewin, B. (1997). Genes. Oxford University Press, Oxford, 6: 1260.

Lewis, J. A. (1987). Checklist and bibliography of benthic marine macroalgae recorded from northern Australia III. Chlorophyta. pp. 1- 56.: Department of Defense. Defense Science and Technology Organisation. Materials Research Laboratories, Melbourne, Victoria, Report MRL-R-1063.

Lewis, L. A., & McCourt, R. M. (2004). Green algae and the origin of land plants. *American J. Bot.* 91(10): 1535–1556.

Lewis, S. M. (1985). Herbivory on coral reefs: algal susceptibility to herbivorous fishes. *Oecologia*. 65: 370-375.

Lin, C. K. (1995). Progression of intensive marine shrimp culture in Thailand. In Browdy CL, Hopkins SJ (eds), Swimming Through Troubled Water. Proceedings of the special session on shrimp farming, Aquaculture '95. World Aquaculture Society, Baton Rouge, Louisiana, U.S.A. 13–23.

Lipkin, Y. (1972). Contribution to the knowledge of Suez canal migration. Marine algal and sea-grass flora of the Suez canal. *J. Zool.* 21: 405– 446.

Lipkin, Y., & Safriel, U. (1971). Intertidal zonationon rocky shore at Mikhmoret (Israel). *J. Ecol.*, 59 (1).

Lisette, N., Fama, P., Bartlett, A. J., Prud, W. F., Van Reine, H., Espinosa, C. A., & et al. (2003). Defining taxon boundaries in members of the morphologically and genetically plastic genus *Caulerpa* (Caulerpales, Chlorophyta). *J. Phycol.* 39: 1019-1037.

Littler, D. S., & Littler, M. M. (2000). Caribbean Reef Plants. Offshore Graphics, Washington, D. C.

Lu'scher-Mattli, M. (2000). Polyanions-a lost chance in the fight against HIV and other virus diseases. *Antivir. Chem. Chemoth.* 11: 249–259.

Ludwig, W., Weizenegger, M., Betzl, D., Leidel, E., Lenz, T., Ludvigsen, A., & et al. (1990). Complete nucleotide sequences of seven eubacterial genes coding for the elongation factor Tu: functional, structural, and phylogenetic evaluations. *Arch. Microbiol.* 153:241–247.

- Madigan, M. T., Martinko, J. M., & Parker, J. (2003). Brock Biology of Microorganisms, 10th edn. Pearson Education, Inc., NJ, U. S. A, pp. 151–156.
- Madlener, J. C. (1977). The Seavegetable Book. Crown Publishers, New York, pp. 288.
- Maiti, B. C., Thomson, R. H., & Mahendran, M. (1978). The structure of caulerpin, a pigment from *Caulerpa* algae. *J. Chem. Res. (S)*: 126-127.
- Malea, P., & Haritonidis, S. (1995). Local distribution and seasonal variation of Fe, Pb, Zn, Cu, Cd, Na, K, Ca and Mg concentrations in the seagrass *Cymodocea nodosa* (Ucria) Aschers. in the Antikyra Gulf, Greece P.S.Z.N.I. *Mar. Ecol.* 16(1): 41–56.
- Malea, P., & Haritonidis, S. (1999a). Metal content in *Enteromorpha linza* Linnaeus in Thermaikos Gulf (Greece). *Hydrobiologia* 394: 103–112.
- Malea, P., & Haritonidis, S. (1999b). Seasonal accumulation of metals by red alga *Gracilaria verrucosa* (Hud.) Papenf. From Thermaikos Gulf, Greece. *J. Appl. Phycol.* 11: 503–509.
- Malea, P., & Haritonidis, S. (2000). Use of the green alga *Ulva rigida* C. Agardh as an indicator species to reassess metal pollution in the Thermaikos Gulf, Greece, *J. Appl. Phycol.* 12: 169–176.
- Marcomini, A., Sfriso, A., & Zanette, M. (1993). Macroalgal blooms, nutrient and trace metal cycles in a coastal lagoon. Proc. COST-48 Syrup. Sub Group III, Thessaloniki, 24-26 September.
- Margulis, L., & Schwartz, K. (1998). Five kingdoms, an illustrated guide to the phyla of life on earth. 3<sup>rd</sup> ed. W. H. Freeman and Company. New York.
- Mathieson, A. C., Norton, T. A., & Neushul, M. (1981). The taxonomic implications of genetic and environmentally induced variations in seaweed morphology. *Bot. Rev.* 47: 313-347.
- Mattox, K. R., & Stewart, K. D. (1984). Classification of the green algae: a concept based on comparative cytology. In Irvin, D. E. G., John, D., (Eds.): Systematics of the Green Algae. 29- 72. - London, New York: Academic Press.
- Maxam, A., & Gilbert, W. (1977). *Proc. Nat. Acad. Sci. USA* 74: 560-564.
- Mayer, A. M. S., & Lehmann, V. K. B. (2000). Marine pharmacologyin 1998: marine compounds with antibacterial, anticoagulant, antifungal, anti-inflammatory, anthelmintic, antiplatelet, antiprotozoal, and antiviral activities; with actions on the cardiovascular, endocrine, immune, and nervous systems; and other miscellaneous mechanisms of action. *Pharmacologist* 42: 62–69.

- McConnel, O. P., Hughes, P. A., Targett, N. M., & Daley, J. (1982). Effects of secondary metabolites from marine algae on feeding by the sea urchin, *Lytechinus variegatus*. *J. Chem. Ecol.* 8:1437–1453.
- McHugh, D. J. (2003). A guide to the seaweed industry. FAO Fisheries Technical Paper No. 441, Rome, p. 105.
- McLachlan, A. (1985). The biomass of macro and interstitial fauna on clean and wrack-covered beaches in Western Australia. *Estuarine, Coastal and Shelf Science*. 21: 587-599.
- Meinesz, A., & Hesse, B. (1991). Introduction et invasion de l'algue tropicale *Caulerpa taxifolia* en Méditerranée nord-occidentale. *Oceanol. Acta* 14 (4): 415–426.
- Meinesz, A., Belsher, T., Thibaut, T., Antolic, B., Ben Mustapha, K., Boudouresque, C. F., & et al. (2001). The introduced green alga *Caulerpa taxifolia* continues to spread in the Mediterranean. *Bio. Inv.* 3: 201–210.
- Meinesz, A., Benichou, L., Blachier, J., Komatsu, T., Lemeé, R., Molenaar, H., & et al. (1995). Variations in the structure, morphology and biomass of *Caulerpa taxifolia* in the Mediterranean. *Bot. Mar.* 38: 499– 508.
- Meñez, E., & Calumpong, H. (1982). The genus *Caulerpa* from Central Visayas, Philippines. Smiths. *Contrib. Mar. Sci.* 17: 1-21.
- Menzel, D. (1987). The cytoskeleton of the giant coenocytic green alga *Caulerpa* visualized by immunocytochemistry. *Protoplasma* 139: 71-76.
- Meusnier, I., Olsen, J. L., Stam, W. T., Destombe, C., & Valero, M. (2001). Phylogenetic analyses of *Caulerpa taxifolia* (Chlorophyta) and of its associated bacterial microflora provide clues to the origin of the Mediterranean introduction. *Mol. Ecol.* 10: 931–946.
- Miyamura, S., & Hori, T. (1989). Presence of DNA in the pyrenoid matrix of the siphonous green alga, *Caulerpa okamurae* Web. v. Bos. *Plant Morphol* 1: 19-22.
- Miyamura, S., & Hori, T. (1991). DNA is present in the pyrenoid core of the siphonous green algae of the genus *Caulerpa* and yellow-green algae of the genus *Pseudodichotomosiphon*. *Protoplasma* 161: 192-196.
- Miyamura, S., Hori, T., Ohya, T., Tohma, T., & Ikawa, T. (1996). Co-localization of chloroplast DNA and ribulose-1, 5-bisphosphate carboxylase/oxygenase in the so-called pyrenoid of the siphonous green alga *Caulerpa lentillifera* (Caulerpales, Chlorophyta). *Phycologia* 35(2): 156-160.
- Mizukami, Y., Kito, H., Kunimoto, M., & Kobayashi, M. (1998). Effect of DNA preparation from laver (*Porphyra yezoensis*) thalli on reproducibility of RAPD (random amplified polymorphic DNA) patterns. *J. Appl. Phycol.* 10: 23–29.

- Modena, M., Matricardi, G., Vacchi, M., & Guidetti, P. (2000). Spreading of *Caulerpa racemosa* (Forsska°l) J. Agardh (Bryopsidaceae, Chlorophyta) along the coasts of the Ligurian Sea. *Cryptogamie Algol.* 21: 301–304.
- Moore, L. J., & Coleman, A. W. (1987). Do the chloroplast DNAs of *Pandorina morum* (Volvocales, Chlorophyta) evolve at the same rate as cp DNAs of flowering plant species. 14th Int. Bot. Congr. Berhn, Abstr. 178.
- Munda, I. M. (1982). The effect of different pollutants on benthic marine algae. Rapp. Com. Int. Mer. M\_edit. VI J. Poll. Cannes. pp. 721–726.
- Munda, I. M., & Hundnik, V. (1991). Trace metal content in some seaweeds from the Northern Adriatic. *Bot. Mar.* 34: 241–249.
- Neori, A., Krom, M. D., Ellner, S. P., Boyd, C. E., Popper, D., Rabinovitch, R., & et al. (1996). Seaweed biofilters as regulators of water quality in integrated fish-seaweed culture units. *Aquaculture* 141: 183–199.
- Neori, A., Cohen, I., & Gordin, H. (1991). *Ulva lactuca* biofilters for marine fishpond effluent. II. Growth rate, yield and C: N ratio. *Bot. Mar.* 34: 483–489.
- Neushul, P. (1987). Energy from marine biomass: the historical record. In Bird KT, Benson PH (eds) Seaweed cultivation for renewable resources. Elsevier, Amsterdam, pp. 1-37.
- Nizamuddin, M. (1964). Studies on the genus *Caulerpa* from Karachi. *Bot. Mar.* 6: 204–223.
- Norris, J. N., & Fenical, W. (1982). Chemical defense in tropical marine algae. In: Rutzler, K., Macintyre, I. G. (ed.) The Atlantic Barrier Reef ecosystem at Carrie Bow Cay, Belize 1: structure and communities. *Smithsonian Contr. mar. Sci.* 12: 417-431.
- North, W. J. (1980). Trace metals in giant kelp. *Macrocystis*. *Am .J. Bot.* 67:1097-101
- North, W. J. (1987). Oceanic farming of *Macrocystis*, the problems and non-problems. In Bird KT, Benson PH (eds), Seaweed Cultivation for Renewable Resources. Elsevier, Amsterdam, pp. 39-67.
- Nuber, N., Gornic, O., Lauc, G., Bauer, N., Zuljevi, A., Papes, D., & et al. (2007). Genetic evidence for the identity of *Caulerpa racemosa* (Forsska°l) J. Agardh (Caulerpales, Chlorophyta) in the Adriatic Sea. *Eur. J. Phycol.* 42 (1): 113–120.
- Ogden, J. C. (1976). Some aspects of herbivore-plant relationships on Caribbean reefs and seagrass beds. *J. Aquat. Bot.* 2: 103-116C.
- Ogden, J. C., & Lobel, P. S. (1981). The role of herbivorous fishes andurchins in coral reef communities. *J. Envir. Biol. Fish.* 3: 49-63.

- Ohba, H., & Enomoto, S. (1987). Culture studies on *Caulerpa* (Caulerpales, Chlorophyceae). II. Morphological variation of *C. racemosa* var. *laetevirens* under various culture conditions. *Jpn. J. Phycol.* 25: 178–88.
- Ohba, H., Nashima, H., & Enomoto, S. (1992). Culture studies on *Caulerpa* (Caulerpales, Chlorophyceae). Reproduction, development, and morphological variation of laboratory-cultured *C. racemosa* var. *peltata*. *Bot. Mag. Tokyo* 105: 589–600.
- Okabe, Y., & Okada, M. (1988). Isolation of native pyrenoid core matrix having ribulose 1,5-bisphosphate carboxylase activity from the green alga *Bryopsis maxima*. *Plant Cell Physiol.* 29: 88–96.
- Okada, M., Okabe, M., Kono, M., & Nakayama, K. (1991). Peptide composition and enzyme activities of isolated pyrenoids from the green alga *Bryopsis maxima*. *Can J. Bot.* 69: 1053–1061.
- Panayotidis, P., & Zuljevic, A. (2001). Sexual reproduction of the invasive green alga *Caulerpa racemosa* var. *occidentalis* in the Mediterranean sea. *Oceanologica acta* 24: 199–203.
- Papenfuss, G. F. (1968). A history, catalogue, and bibliography of the Red Sea benthic algae. *Israel Journal of Botany* 17: 1–118.
- Parent-Massin, D., Fournier, V., Amade, P., Lemeé, R., Durand-Clement, M., Delescluze, C., & et al. (1996). Evaluation of caulerpenyne toxicological risk on human cells in culture: hematopoietic progenitors, melanocytes and keratinocyte cell lines. *J. Toxicol. Env. Health* 47: 47–59.
- Park, Y. T., Park, J. B., Chung, S. Y., Song, B. C., Lim, W. A., Kim, C. H., & et al. (1998). Isolation of marine bacteria killing red tide microalgae. I. Isolation and algicidal properties of *Micrococcus* sp. LG-1 possessing killing activity for harmful Dinoflagellate *Cochlodinium polykrikoides*. *J. Korean Fish. Soc.* 31: 767–773.
- Patwary, M. U., MacKay, R. M., & van der Meer, J. P. (1993). Revealing genetic markers in *Gelidium vagum* (Rhodophyta) through the random amplified polymorphic DNA (RAPD) technique. *J. Phycol.* 29: 216–222.
- Paul, V. J. (1985). The natural products chemistry and chemical ecology of tropical green algae of the order Caulerpales. Ph.D. dissertation, Scripps Institution of Oceanography, Univ. of California, San Diego.
- Paul, V. J., & Fenical, W. (1983). Isolation of halimedatriah Chemical defense adaptation in the calcareous reef building alga *Halimeda*. *J. Science Wash. D.C.* 221: 747–749.
- Paul, V. J., & Fenical, W. (1986 a,b). Chemical defense in tropical green algae, order Caulerpales. *J. Mar. Ecol. Prog. Ser.* 34: 157–169.

- Paul, V. J., Littler, M. M., Littler, D. S., & Fenical, W. (1987). Evidence for chemical defense in tropical green alga *Caulerpa ashmeadii* (Caulerpaceae: Chlorophyta): isolation of new bioactive sesquiterpenoids. *J. chem. Ecol.* 13 (5): 1171-1185.
- Payri, C., De N'Yeurt, A., & Orempuller, J. (2000). Algae of French Polynesia. Au Vent des Iles, Papeete, Tahiti.
- Pereira, L., Sousa, A., Amado, A. M., & Ribeiro, P. J. A. (2003). Use of FTIR, FT-Raman and <sup>13</sup>C-NMR spectroscopy for identification of some seaweed phycocolloids. *Biomol. Eng.* 20: 223-228.
- Pesando, D., Leme e, R., Ferrua, C., Amade, P., & Girard, J. P. (1996). Effects of caulerpenyne, the major toxin from *Caulerpa taxifolia* on mechanisms related to sea urchin egg cleavage. *J. Aquat Toxicol.* 35: 139-155.
- Peterson, R. D. (1972). Effect of light intensity on the morphology and productivity of *Caulerpa racemosa*. *Micronesica* 8: 63-86.
- Pham-Ho ng, H. (1969). Rong bi n Vi tnam. Marine algae of South Vietnam. pp. 558.
- Phang, S. M. (1994). Some species of *Gracilaria* from Peninsular Malaysia and Singapore. In Abbott IA (ed), Taxonomy of Economic Seaweeds Vol. IV. California Sea Grant Programme, University of California, California, pp.125-134.
- Phang, S. M. (1998). The seaweed resources of Malaysia. In Critchley, A. T. & M. Ohno (eds), Seaweed Resources of the World. JICA, Japan: 79-91.
- Phang, S. M., Lim, P. E., Yeong, H. Y., Ng, W. S., & Song, S. L. (2010). Marine algae collected during the scientific expedition to Bachok, Kelantan and the Islands of Terengganu with one New Record, *Pterocladiella* from Malaysia. *Malaysian J. Sci.* 29: 31-46.
- Phillips, J. A. (1997). Algae. In: *Queensland Plants: Names and Distribution*. (Henderson, R.J.F. Eds), pp. 223-240. Indooroopilly, Queensland: Queensland Herbarium, Department of Environment.
- Phillips, J. A. (2002). Algae. In: Names and distribution of Queensland plants, algae and lichens. (Henderson, R.J.F. Eds), pp. 228-244. Brisbane: Queensland Government Environmental Protection Agency.
- Piazzi, L., Balestri, E., & Cinelli, F. (1994). Presence of *Caulerpa racemosa* in the north-western Mediterranean. *Cryptogamie., Algol.* 15: 183-189.
- Piazzi, L., Balestri, E., Magri, M., & Cinelli, F. (1997). Expansion de l'algue tropicale *Caulerpa racemosa* (Forsska l) J. Agardh (Bryopsidophyceae, Chlorophyta) le long de la co te toscane (Italie). *Cryptogamie., Algol.* 18, 343-350.

Pickett-Heaps, J. D. (1975). Green algae. Structure, Reproduction and Evolution in Selected Genera. - Sunderland, Mass.: Sinauer Associates.

Pillman, A., Woolcott, G. W., Olsen, J. L., Stam, W. T., & King, R. J. (1997). Inter- and intraspecific genetic variation in *Caulerpa* (Chlorophyta) based on nuclear rDNA ITS sequences. *Eur. J. Phycol.* 32: 379–86.

Potin, P., Bouarab, K., Kupper, F., & Kloareg, B. (1999). Oligosaccharide recognition signals and defence reactions in marine plant–microbe interactions. *Curr. Opin. Microbiol.* 2: 276–283.

Potoky, P., & Maze, J. (1988). Effect of seaweed inclusion in compost preparation on the quality of the composts' obtained. In P. Morand & E. H. Schulte (eds), Aquatic Primary Biomass (Marine Macroalgae): Biomass Conversion, Removal and Use of Nutrients. I. Proceedings of the 1st Workshop of the COST 48 Sub-Group 3. L'Houmeau, France. February 12-14, 1987. E.E.C., Brussels: 67-86.

Primavera, J. H. (1993). A critical review of shrimp pond culture in the Philippines. *Rev. fish. Sci.* 1: 151–201.

Provasoli, L., & Pintner, I. J. (1980). Bacteria induced polymorphism in an axenic laboratory strain of *Ulva lactuca* (Chlorophyceae). *J. Phycol.* 16: 196–201.

Prud'homme van Reine, W. F., & Lokhorst, G. M. (1992). *Caulerpella* gen. nov. a non holocarpic member of the Caulerpales (Chlorophyta). *Nova. Hedwigia.* 54:113-126.

Prud'homme van Reine, W. F., Verheij, E., & Coppejans, E. (1996). Species and ecads of *Caulerpa* (Ulvophyceae, Chlorophyta) in Malesia (South-East Asia): Taxonomy, biogeography and biodiversity. *Netherl. J. aquatic Ecol.* 30: 83-98.

Rangaswami, G., & Bagyaraj, D.J. (2004). Agricultural microbiology. 2<sup>nd</sup> ed. pp. 60-135.

Rice, E. L., & Bird, C. J. (1990). Relationships among geographically distant populations of *Gracilaria verrucosa* (Gracilariales, Rhodophyta) and related species. *J. Phycol.* 29: 501–510.

Roels, O. A., Haines, K. C., & Sunderlin, J. B. (1976). The potential yield of artificial upwelling mariculture. In Persson G, Jaspers E (eds), Proc. 10th *Eur. Symp. Mar. Biol.*, Ostend, Belgium, Sept. 17–23, 1975. Universa Press, Wetteren. 1: 358–394.

Ryther, J. H., Goldman, J. C., Gifford, J. E., Huguenin, J. E., Wing, A. S., Clarner, J. P., & et al. (1975) Physical models of integrated waste recycling-marine polyculture systems. *Aquaculture.* 5: 163–177.

Sahoo, D., Nivedita & Debasish (2001). *Seaweeds of Indian coast.* pp. xxi + 283. New Delhi: A.P.H. Publishing.

- Salisbury, J. L., & Floyd, G. L. (1978). Molecular, enzymatic and ultra structure characterization of the pyrenoids of scaly green monad *Micromonas squamata*. *J. Phycol.* 14: 362-368.
- Sanger, F., & Coulson, A. R. (1977). The use of thin acrylamide gels for DNA sequencing. *FEBS Lett.* 87: 107-110.
- Satoh, H., Okada, M., Nakayama, K., & Miyaji, K. (1984). Purification and further characterization of pyrenoid proteins and ribulose-1,5-bisphosphate carboxylate-oxygenase from the green alga *Bryopsis maxima*. *Plant. cell. Physiol.* 25: 1205-1214.
- Sfriso, A., Marcomini, A., & Zanette, M. (1995). Heavy metals in sediments, SPM and phytozoobenthos of the lagoon of Venice. *Mar. Pollut. Bull.* 30 (2): 116–124.
- Shiba, T. (1992). The genus *Roseobacter*. In The Prokaryotes: a Handbook on the Biology of Bacteria: Ecophysiology, Isolation, Identification, Applications (eds Starr, M. P., Stolp, H., Trüpper, H. G., Balows, A., Schlegel, H. G.). pp. 202–206.
- Silva, P. C., Basson, P. W., & Moe, R. L. (1996). Catalogue of the benthic marine algae of the Indian Ocean. *U. Calif. Public. Bot.* 79: 1-1259.
- Silva, P. C., Meñez, E. G. & Moe, R. L. (1987). Catalog of the benthic marine algae of the Philippines. *Smithsonian Contributions to Marine Sciences* 27: 1-179.
- Skelton, P. A., & South, G. R. (2004). New records and notes on marine benthic algae of American Samoa - Chlorophyta & Phaeophyta. *Cryptogamie Algologie* 25: 291-312.
- Stam, W. T., Olsen, J. L., Frisch Zaleski, S., Murray, S. N., Brown, K. R., & Walters, L. J. (2006). A forensic and phylogenetic survey of *Caulerpa* species (Caulerpales, Chlorophyta) from the Florida coast, local aquarium shops, and ecommerce: Establishing a proactive baseline for early detection. *J. Phycol.* 42: 1113-1124.
- Staves, M. P., & LaClaire, J. W. (1985). Nuclear synchrony in *Valoniamacrophypha* (Chlorophyta): light microscopy and flow cytometry. *J. Phycol.* 21:68–81.
- Svedelius, N. (1906). Reports on the marine algae of Ceylon. No. I. Ecological and systematic studies of the Ceylon species of *Caulerpa*. *Ceylon Mar. Biol. Rep.* 4: 81–144.
- Swofford, D. L. (2002) PAUP\*. Phylogenetic Analysis Using Parsimony (\*and Other Methods), 4.0 edn, Sinauer, Sunderland, MA.
- Sze, P. (1986). A Biology of the Algae. Wm. C. Brown Publishers. Dubuque, Iowa.
- Sze, P. (1998). A biology of the algae, Third ed. McGraw-Hill, Boston, 278 p.

- Targett, N. M., Targett, T. E., Vrolijk, N. H., & Ogden, J. C. (1986). The effect of macrophyte secondary metabolites or feeding preferences of the herbivorous parrotfish *Sparisoma radians*. *Mar. Biol.* 92: 141-148.
- Taylor, W. R. (1960). Marine Algae of the Eastern Tropical and Subtropical Coasts of the Americas. Ann Arbor, University Michigan Press, pp. 870.
- Teo, L. W., & Wee, Y. C. (1983). Seaweeds of Singapore. pp. 123. Singapore: Singapore University Press.
- Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22: 4673–4680.
- Trono, G. C., & Ganzon-Fortes, E. T. (1988). Philippine Seaweeds. National Book Store, Inc., Manila, Philippines. pp. 330.
- Tseng, C. K. (1981). Commercial Cultivation. In Lobban, C. S. and Wynne, M. J. (eds.), *The Biology of Seaweeds*, 680-725. Blackwell Scientific Publication, Oxford.
- Vadas, R. L. (1977). Preferential feeding: an optimization strategy in sea urchins. *J. Ecol. Monogr.* 47: 337-371.
- Valatka, S., Makinen, A., & Yli-Mattila, T. (2000). Analysis of genetic diversity of *Furcellaria lumbricalis* (Gigartinales, Rhodophyta) in the Baltic Sea by RAPD-PCR technique. *J. Phycol.* 39(2): 109–117.
- Vallini, G., A. Pera, F. Cecchi, M. M., & Sicurani, M. A. (1993). Compost Stabilization of Algal Biomass Drawn In Eutrophic Lagoon Ecosystems. *Compost Sci. Util.* 1 (2): 49-53.
- Van den Heede, C. (1994). De Codiales (Chlorophyta) van Kenya, Tanzania en de Seychellen. MSc thesis Universiteit Gent. Deel 1: 160pp., deel 2: 73 pl.
- Van Den Heede, C., & Coppejans, E. (1995). The genus *Codium* (Chlorophyta, Codiales) from Kenya, Tanzania (Zanzibar) and the Seychelles *Nova Hedwigia* 62: 389-417.
- Van den Hoek, C., Mann, D. G., & Jahns, H. M. (1995). Algae. An Introduction to Phycology. Cambridge University Press, Cambridge, pp. 623.
- Van Oppen, M. J. H., Klerk, H., de Graaf, M., Stam, W. T., & Olsen, J. L. (1996). Assessing the limits of random amplified polymorphic DNAs (RAPDs) in seaweed biogeography. *J. Phycol.* 32: 433–444.

- Vandermeulen, H., & Gordin, H. (1990). Ammonium uptake using *Ulva* (Clorophyceae) in intensive fishpond systems: mass culture and treatment of effluent. *J. Appl. Phycol.* 2: 263–374.
- Verellen, A. (1990). *Avrainvillea, Chlorodesmis, Halimeda, Udotea*: Caulerpales (Chlorophyta) van de Keniaanse kust. MSc thesis Universiteit Gent. Deel 1: 99pp., deel 2: 70pp.
- Verheij, E., & Prud'homme van Reine, W. F. (1993). Seaweeds of the Spermonde Archipelago, SW Sulawesi, Indonesia. *Blumea* 37: 385-510.
- Verlaque, M., Afonso-Carrillo, J., Gil-Rodriguez, M. C., Durand, C., Boudouresque, C. F., & Le Parco, Y. (2004). Blitzkrieg in a marine invasion: *Caulerpa racemosa* var. *cylindracea* (Bryopsidales, Chlorophyta) reaches the Canary Islands (north-east Atlantic). *Biol. Invas.* 6: 269–281.
- Verlaque, M., Afonso-Carrillo, J., Gil-Rodriguez, M. C., Durand, C., Boudouresque, C. F., & et al.(2004). Blitzkrieg in a marine invasion: *Caulerpa racemosa* var. *cylindracea* (Bryopsidales, Chlorophyta) reaches the Canary Islands (north-east Atlantic). *Biol. Invas.* 6: 269–281.
- Verlaque, M., Boudouresque, C. F., Meinesz, A., & Gravez, V. (2000). The *Caulerpa racemosa* complex (Caulerpales, Ulvophyceae) in the Mediterranean Sea. *Bot. Mar.* 43: 49–68.
- Verlaque, M., Durand, C., Huisman, J. M., Boudouresque, C. F., & Le Parco, Y. (2003). On the identity and origin of the Mediterranean invasive *Caulerpa racemosa* (Caulerpales, Chlorophyta). *Eur. J. Phycol.* 38: 325–339.
- Volckaert F. A. M., Batargas C., Cana'rio A., Chatziplis D., Chistiakov D. A., Haley C., & et al. (2008). The European sea bass (*Dicentrarchus labrax* L.) and its genomic resources. In: Genome Mapping and Genomics in Fishes and Aquatic Animals (Ed. by C. Kole), pp. 117–34. Springer Verlag, Berlin.
- Vos, P., Hogers, R., Bleeker, R., Reijans, M., Van de Lee, T., Horne, M., & et al. (1995). AFLP: a new technique for DNA Fingerprinting. *Nucleic Acids Res.* 23: 4407-4414.
- Weber-van Bosse, A. (1898). Monographie des Caulerpes. *Ann. Jardin Bot. Buitenzorg.* 15: 243–401.
- Welsh, J., & McClelland, M. (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res.* 18: 7213.
- Wiedenmann, J., Baumstark, A., Pillen, T. L., Meinesz, A., & Vogel, W. (2001). DNA fingerprints of *Caulerpa taxifolia* provide evidence for the introduction of an aquarium strain into the Mediterranean Sea and its close relationship to an Australian population. *Mar. Biol.* 138: 229–234.

- Williams, J. G. K., Kubelik, A. R., Livak, K., Rafalski, J., & Tingey, S. V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res.* 18: 6531-6535.
- Witvrouw, M., & De Clercq, E. (1997). Sulfated polysaccharides extracted from sea algae as potential antiviral drugs. *Gen. Pharmacol.* 29: 497–511.
- Wylie, C. R., & Paul, V. J. (1988). Feeding preferences of the surgeonfish Zebrasoma flavescens in relation to chemical defenses of tropical algae. *J. Mar. Ecol. Prog. Ser.* 45: 23-32.
- Yamada, Y. (1950). A list of marine algae from Ryukyusho, Formosa. *Sci. Pap. Inst. Algol. Res., Fac. Sci., Hokkaido Imperial Univ.* 3: 173-194.
- Yeh, W. J., & Chen, G. Y. (2004). Nuclear rDNA and internal transcribed spacer sequences clarify *Caulerpa racemosa* vars. from other *Caulerpa* species. *Aquat. Bot.* 80 (3): 193-207.