

REFERENCES

- Abdullah, R., Alizah, R., Wee, Y. H., Leaw, C. L., Yeap, C. B., Salwa, A. S., Yap, W. S. P., Joseph, J. L., Muhammad Rashdan, M.. and Yuen, L. H. (2005). Immature embryo: a useful tool for oil palm (*Elaeis guineensis. Jacq*) genetic transformation studies. *Electronic journal of Biotechnology* 8: 24-34.
- Altschul, S. F., Madden, T. L., Schaffer, A..A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389-3402.
- Advisory Committee on Novel Foods and Processes (ACNFP). (1994).
- Aragão, F. J. L. and Brasileiro, A. C. M.. (2002). Positive, negative and marker-free strategies for transgenic plant selection. *Braz. J. Plant Physiol.* 14: 1-10.
- Aswath, C. R., Mo, S. Y., Kim, D. H. and Park, S. W. (2006). *Agrobacterium* and biolistic transformation of onion using non-antibiotic selection marker phosphomannose isomerase. *Plant Cell Rep.* 25: 92-99.
- Barb, A, W. (2002). Phosphomannose isomerase in *Nicotiana tabacum* L. NT1 and *Apium graveolens* var. *dulce* L. cell suspension cultures. MSc Thesis Submitted to North Carolina State University.
- Basiron, Y. and Chan, K. W. (2005). The role of research and development strategies in food safety and good agricultural, manufacturing and distribution practices in the Malaysian palm oil industry. *Oil Palm Industry Economic Journal* 5: 1-16.
- Bevan, M. W., Flavell, R. B. and Chilton, M. D. (1983). A chimaeric antibiotic resistance gene as a selectable marker for plant cell transformation. *Nature* 304: 184-187.
- Bhatti, K. H. and He, C. (2009). *Agrobacterium* mediated tobacco transformation with rice Fae gene and segregation analysis of T1 generation. *Pak. J. Bot.*, 41(1): 403-412.
- Birch, R. G. (1997). Plant Transformation: Problems And Strategies For Practical Application. *Annu. Rev. Plant Physiol. Mol. Biol.* 48: 297-326.
- Bojsen, K., Donaldson, I., Haldrup, A., Joersbo, M., Kreiberg, J., Nielsen, J., Okkels, F. T. and Petersen, S. G. (1998). Mannose Or Xylose Based Positive Selection. US Patent 5767378.
- Boscariol, R. L., Almeidawab., Derbyshire, M. T. V. C., Mourao Filho, F. A. A. and Mendes, B. M. J. (2003). The use of the PMI/Mannose selection system to recover transgenic sweet orange plants (*Citrus Sinensis* L. Osbeck). *Plant Cell Rep.* 22:122–128.
- Brasileiro, A. C. M.. and Dusi, D. M. A. (1999). *Transformação Genética De Plantas*. In: Torres AC, Caldas LS, Buso JA (Eds), *Cultura De Tecidos E Transformação Genética De Plantas*, Pp.679-735. Embrapa-SPI/Embrapa-CNPH, Brasília, Brazil.

- Briza, J., Pavingerova, D., Prikrylova, P., Gazdova, J., Vlasak, J. and Niedermeierova, H. (2008). Use of phosphomannose isomerase-based selection system for *Agrobacterium*-mediated transformation of tomato and potato. *Biologia Plantarum* 52 (3): 453-461.
- Carvalho, C. H. S., Zehr, U. B., Gunaratna, N., Anderson, J., Kononowicz, H. H., Hodges, T. K. and Axtell, J. D. (2004). *Agrobacterium*-mediated transformation of sorghum: factors that affect transformation efficiency. *Genet Mol Biol* 27:259–269.
- Chalfie, M., Tu, Y., Euskirchen, G., Ward, W. W. and Prasher, D. C. (1994). Green fluorescent protein as a marker for gene expression. *Science* 263: 802-805.
- Cheng, X., Sardana, R., Kaplan, H. and Altosaar, I. (1998). *Agrobacterium*-mediated rice plants expressing synthetic *cryIA (b)* and *cryIA (c)* genes are highly toxic to striped stem borer and yellow stem borer. *Proc. Natl. Acad. Sci. USA*, 95: 2767–2772.
- Chia, T. F., Chan, Y. S. and Chua, N. H. (1994). The firefly luciferase gene as a non-invasive reporter for *Dendrobium* transformation. *Plant J.* 6: 441-446.
- Christensen, A. H. and Quail, P. H. (1996). Ubiquitin promoter-based vectors for high-level expression of selectable and/or screenable marker genes in monocotyledonous plants. *Trans Res* 5:213–218.
- Chowdhury, M.K.U., Parveez, G.K.A. and Saleh, N.M. (1997). Evaluation of five promoters for use in transformation of oil palm (*Elaeis Guineensis* Jacq.). *Plant Cell Reports*. 16: 277-281.
- Collins, L. V. and Hackett, J. (1991). Molecular cloning, characterization, and nucleotide sequence of the *rfc* gene, which encodes an O-antigen polymerase of *Salmonella typhimurium*. *J Bacteriol* 173: 2521±2529.
- Comai, L., Facciotti, D. Hiatt, W. R., Thompson, G., Rose, R. E. and Stalker, D. M. (1985). Expression in plants of a mutant *Aroa* gene from *Salmonella Typhimurium* confers resistance to glyphosate. *Nature* 317: 741-744.
- Corley, R. H. V. (1983). Potential productivity of tropical perennial crops. *Experimental Agriculture* 19: 217-237.
- Corley, R. H. V., Lee, C. H., Law, L. H. and Wong, C. Y. (1986). Abnormal development in oil palm clones. *The Planter*, Kuala Lumpur 62:233–240.
- Corley, R.H.V. and Tinker P.B. (2003). *The Oil Palm* (4th ed.). Blackwell Science Ltd, Oxford, pp. 201-215.
- Daniell, H., Wiebe, P. O. and Millan, A. F. (2001). Antibiotic-free chloroplast genetic engineering - an environmentally friendly approach. *Trends Plant Sci.* 6:237-239.
- Darzins, A., Frantz, B., Vanags, R. I. and Chakrabarty, A. M. (1986). Nucleotide sequence analysis of the phosphomannose isomerase gene (*pmi*) of *pseudomonas*

aeruginosa and comparison with the corresponding *Escherichia coli* gene *manA*. *Gene* 42:293-302.

- Datta, K., Baisakh, N., Oliva, N., Torrizo, L., Abrigo, E., Tan, J., Rai, M., Rehana, S., Al-Babili, S., Beyer, P., Potrykus, I. and Datta, S. K. (2003). Bioengineered ‘golden’ indica rice cultivars with •-carotene metabolism in the endosperm with hygromycin and mannose selection systems. *Plant Biotechnol J* 1:81–90.
- Dayang Izawati, A. M., Parveez, G. K. A. and Masani, M. Y. A. (2008). Method for producing transgenic plants. Transfer of Technology 2008. Malaysian Palm Oil Board.
- Dayang Izawati, A. M. (2009). “Development of technique for production of transgenic plant using positive selectable marker (2-Deoxyglucose-6-Phosphate Phosphatase)”. Personal communication.
- De Block, M., Botterman, J., Vandewiele, M., Dockx, J., Thoen, C., Gossele, V., Movva, N. R., Thompson, C., Van Montagu, M. and Leemans, J. (1987). Engineering herbicide resistance in plants by expression of a detoxifying enzyme. *EMBO J.* 6: 2513-2518.
- Degenhardt, J., Poppe, A., Montag, J. and Szankowski, I. (2006). The use of the phosphomannose isomerase/mannose selection system to recover transgenic apple plants. *Plant Cell Rep* 25: 1149-1156.
- Dekeyser, R. B., Claes, M., Marichal, M., Van Montagu, M. and Caplan, A. (1989). Evaluation of selectable markers for rice transformation. *Plant Physiol.* 90: 217-223.
- Doyle, J. J. and Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissues. *Phytochemical Bull.* 19: 11-15.
- Eichholtz, D. A., Rogers, S. G., Horsch, R. B., Klee, H. J., Hayford, M. and Hoffman, N. L. (1987). Expression of mouse dihydro folate reductase gene confers methotrexate resistance in transgenic petunia plants. *Somatic Cell Mol Genet.* 13:67–76.
- Enríquez-Obregón, G.A., Vázquez-Padrón, R.I., Prieto-Samsónov, D.L., Pérez, M. and Selman-Housein, G. (1997). Genetic transformation of sugarcane by *Agrobacterium tumefaciens* using antioxidants compounds. *Biotecnología Aplicada* 14: 169-174
- Enríquez-Obregón, G. A., Vázquez-padrón, R. I., Prieto-sansonov, D. L., de la Riva, G. A. and Selman-Housein, G. (1998). Herbicide resistant sugarcane (*Saccharum officinarum* L.) plants by *Agrobacterium*-mediated transformation. *Planta* 206: 20-27.
- Evans, R., Wang, A., Hanten, J., Altendorf, P. and Mettler, I. (1996). A positive selection system for maize transformation. *In Vitro* 32 (3. Part II): 72A (abstract).

- Feeney, M. and Punja, Z. K. (2003). Tissue culture and *Agrobacterium* mediated transformation of hemp (*Cannabis Sativa L.*). *In Vitro Cell. Dev. Biol. Plant.* 39: 578-585.
- Ferber, D. (1999) GM crops in the cross hairs. *Science* 286: 1662–1666.
- Ferguson, J. D., Street, H. E. and Avid, S. B. (1958). The carbohydrate nutrition of tomato roots. IV. The inhibition of excised root growth by galactose and mannose and its reversal by dextrose and xylose. *Ann. Bot.* 22: 525-538.
- Finer, J. J. and McMullen, M. D. (1991). Transformation of soybean via particle bombardment of embryogenic suspension culture tissue. *In Vitro Cell. Dev. Biol.* 27: 115-182.
- Finer, J. J., Vain, P., Jones, M. W. and McMullen, M. (1992). Development of the particle inflow gun for DNA delivery to plant cells. *Plant Cell Rep.* 11: 323-328.
- Fraley, R. T., Rogers, S. G., Horsch, R. B., Sanders, P. R., Flick, J. S., Adams, S. P., Bittner, M.L., Brand, L.A., Fink, C.L., Fry, J.S., Galluppi, G.R., Goldberg, S.B., Hoffmann. N.L. and Woo, S.C. (1983). Expression of bacterial genes in plant cells. *Proc Nat. Acad. Sci. USA* 80: 4803-4807.
- Fromm, M., Taylor, L. P. and Walbot, V. (1986). Stable transformation of maize after gene transfer by electroporation. *Nature* 319: 791-793.
- Fu, D. L., Xiao, Y. M., Muthukrishnan, S. and Liang, G. H. (2005). *In Vivo* performance of a dual genetic marker, *ManA-Gfp*, in transgenic bentgrass. *Genome*. 48: 722-730.
- Gadaleta, A., Giancaspro, A. and Blanco, A. (2004). Phosphomannose Isomerase, *PMI*, as selectable marker gene in durum wheat transformation. Proceedings of the XLVIII Italian Society of Agricultural Genetics.
- Gao, Z., Xie, X., Ling, Y., Muthukrisnan, S. and Liang, G. H. (2005). *Agrobacterium Tumefaciens*- mediated sorghum transformation using a mannose selection system. *Plant Biotechnology Journal*. 3(6): 591.
- Gelvin, S. B. (2003). *Agrobacterium*-mediated plant transformation: The biology behind the “gene-jockeying” tool. *Microbiology and Molecular Biology Reviews* 67 (1): 16-37.
- George, E. F. (1993). *Plant propagation by tissue culture*. Edington Wilts, Extergetics LTD.
- Goddijn, O. J. M., Van der Duyn Schouten, P. M., Schilperoort, R. A., and Hoge, J. H. C. (1993). A chimeric tryptophan decarboxylase gene as a novel selectable marker in plant cells. *Plant Mol Biol.* 22:907–12.
- Goldsworthy, A. and Street, H. E. (1965). The carbohydrate nutrition of tomato roots VIII. The mechanism of the inhibition by D-mannose of the respiration of excised roots. *Ann Bot* 29: 45-58.

- Gordon-Kamm, W. J., Spencer, T. M., Mangano, M. L., Adams, T. R., Daines, R. J., Start, W. G. O., Brian, J. V., Chambers, S. A., Adams, J. W. R., Willetts, N. G., Rice, T. B., Mackey, C. J., Krueger, W., Kausch, A. P. and Lemaux, P. G. (1990). Transformation of maize cells and regeneration of fertile transgenic plants. *Plant Cell* 2: 603-618.
- Gracy, R.W. and Noltmann, E.A. (1968). Studies on phosphomannose isomerase. II. Characterization as a zinc metalloenzyme. *J. Biol. Chem.* 243: 4109–4116.
- Graham, L. A., Liou, Y. C., Walker, V. K. and Davies, P. L. (1997). Hyperactive antifreeze protein from beetles. *Nature* 388: 728-729.
- Haldrup, A., Petersen, S. G. and Okkels, F. T. (1998a). Positive selection: A plant selection principle based on xylose isomerase, an enzyme used in the food industry. *Plant Cell Rep.* 18: 76-81.
- Haldrup, A., Petersen, S. G. and Okkels, F. T. (1998b). The xylose isomerase gene from *Thermoanaerobacterium thermosulfurogenes* allows effective selection of transgenic plant cells using D-xylose as the selection agent. *Plant Mol. Biol.* 37: 287-296.
- Hanahan, D. (1982). Studies on transformation of *Escherichia coli* with plasmids. *Journal of Molecular and Biology* 166: 557-580.
- Hansen, G., Shillito, R. D. and Chilton, M. D. (1997). T-strand integration in maize protoplasts after codelivery of a T-DNA substrate and virulence genes. *Proceedings of the National Academy of Sciences USA* 94: 11726-11730.
- Hansen, G. and Wright, M. S. (1999). Recent advances in the transformation of plants. *Trends Plant Sci* 4: 226-231.
- Hartley, C. W. S. (1988). "The oil palm". 3rd ed. Essex: Longman Scientific and Technical.
- Haughn, G. W., Smith, J., Mazur, B. and Somerville, C. (1988) Transformation with a mutant *Arabidopsis* acetolactate synthase gene renders tobacco resistant to sulfonylurea herbicides. *Mol. Gen. Genet.* 211: 266-271.
- Hayford, M. B., Medford, J. I., Hoffman, N. L., Rogers, S. G. and Klee, H. J. (1988). Development of a plant transformation selection system based on expression of genes encoding gentamicin acetyltransferase. *Plant Physiol.* 86: 1216-1222.
- He, Z., Fu, Y., Si, H., Hu, G., Zhang, S., Yu, Y. and Sun, Z. (2004). Phosphomannose isomerase (*pmi*) gene as a selectable marker for rice transformation via *Agrobacterium*. *Plant Sci* 166: 17–22.
- He, Z. Q., Duan, Z. Z., Liang, W., Chen, F. J., Yao, W., Liang, H. W., Yue, C. Y., Sun, Z. X., Chen, F. and Dai, J. W. (2006). Mannose selection system used for cucumber transformation. *Plant Cell Rep.* 25: 953-958.

- Herrera-Estrella, L., De Block, M., Messens, E., Hernalsteens, J. P., Van Montagu, M. and Schell, J. (1983). Chimeric genes as dominant selectable markers in plant cells. *EMBO J.* 2: 987-995.
- Herold, A. and Lewis. (1977). Mannose and green plants: occurrence, physiology and metabolism, and use as a tool to study the role of orthophosphate. *New Phytol.* 79: 1-40.
- Hille, J., Verheggen, F., Roelvink, P., Franssen, H., Van Kammen, A. and Zabel, P. (1986). Bleomycin resistance: a new dominant selectable marker for plant cell transformation. *Plant Mol. Biol.* 7: 171-176.
- Hu, T., Metz, S., Chay, C., Zhou, H. P., Biest, N., Chen, G., Cheng, M., Feng, X., Radionenko, M., Lu, F. and Fry, J. (2003) Agrobacterium-mediated large-scale transformation of wheat (*Triticum aestivum* L.) using glyphosate selection. *Plant Cell Rep.* 21: 1010-1019.
- Irdani, T., Bogani, P., Mengoni, A., Mastromei, G. and Buiatti, M. (1998). Construction of a new vector conferring methotrexate resistance in *Nicotiana tabacum* plants. *Plant Mol Biol.* 37:1079-84.
- Jain, M., Chengalrayan, K., Abouzid, A. and Gallo, M. (2007). Prospecting the utility of a PMI/mannose selection system for the recovery of transgenic sugar cane (*Saccharum* spp. hybrid) plants. *Plant Cell Rep* 26:581-590.
- Jang, J. C. and Sheen, J. (1994). Sugar sensing in higher plants. *Plant Cell* 6: 1665-1679.
- Jang, J. C. and Sheen, J. (1997). Sugar sensing in higher plants. *Trends Plant Sci.* 2: 208-214.
- Jefferson, R. A. (1987). Assaying chimeric genes in plants: The GUS gene fusion system. *Plant Mol. Biol. Rep.* 5: 387-405.
- Jefferson, R. A. (1989). The GUS Reporter Gene System. *Nature.* 342: 837-838.
- Joersbo, M. and Okkels, T. (1996). A novel principle for selection of transgenic plant cells: positive selection. *Plant Cell Rep* 16: 219-221.
- Joersbo, M., Donaldson, I., Kreiberg, J., Petersen, S. G., Brunstedt, J. and Okkels, F. T. (1998). Analysis of mannose selection used for transformation of sugar beet. *Mol Breeding* 4: 111-117.
- Joersbo, M., Pederson, S. G. and Okkels, F. T. (1999). Parameters interacting with mannose selection employed for the production of transgenic sugar beet. *Physiol Plant* 105: 109-115.
- Joersbo, M., Mikkelsen, D. J. and Brunstedt, J. (2000). Relationship between promoter strength and transformation frequencies using mannose selection for the production of transgenic sugar beet. *Mol Breeding* 6: 207-213.

- Joersbo, M. (2001). Advances in the selection of transgenic plants using non-antibiotic marker genes. *Physiol Plant* 111: 269-272.
- Jones, L. H. (1974). Propagation of clonal oil palms by tissue culture. *Oil Palm News*. 17: 1-8.
- Jones, J. D. G., Svab, Z., Harper, E. C., Hurwitz, C. D. and Maliga, P. (1987). A dominant nuclear streptomycin resistance marker for plant cell transformation. *Mol. Gen. Genet.* 210: 86-91.
- Kanchanapoom, K., Nakkaew, A., Kanchanapoom, K. and Phongdara, A. (2008). Determination of the most efficient target tissue and helium pressure for biolistic transformation of oil palm (*Elaeis guineensis* Jacq.). *Songklanakarin J. Sci. Technol.* 30 (2): 153-157.
- Ke, J., Khan, R., Johnson, T., Somers, D. A. and Das, A. (2001). High-efficiency gene transfer to recalcitrant plants by *Agrobacterium tumefaciens*. *Plant Cell Rep.* 20, 150–156.
- Kim, J. Y., Jung, M., Kim, H. S., Lee, Y. H., Choi, S. H., Lim, Y. P., Min, B. W., Yang, S. G. and Harn, C. H. (2002). A new selection system for pepper regeneration by mannose. *J. Plant Biotechnol* 4: 129-134.
- Klein, T. M. and Jones, T. (1999). Methods of genetic transformation: the gene gun. P21-42. In: Vasil IK (Ed.) *Molecular improvement of cereal crops*. Kluwer, Dordrecht, The Netherlands.
- Klein, T. M., Wolf, E. D., Wu, R. and Sanford, J. C. (1987). High-velocity microprojectiles for delivering nucleic acids into living cells. *Nature* 327: 70-73.
- Koncz, C., Németh, K., Redei, G. P. and Scell, J. (1994). Homologous recombination and gene silencing in plants. Kluwer, Dordrecht, The Netherlands. *J Paszkowski* (ed.). 167-189
- Koziel, M. G., Adams, T. L., Hazlet, M. A., Damm, D., Miller, J. and Dahlbeck, D. (1984). A cauliflower mosaic virus promoter directs expression of kanamycin resistance in morphogenic transformed plant cells. *J Mol Appl Genet.* 2:549–62.
- Ku, J. J., Park, Y. W. and Park, Y. D. (2006). A non-antibiotic selection system uses the Phosphomannose-Isomerase (PMI) gene for *Agrobacterium*-mediated transformation of chinese cabbage. *J. Plant Biol.* 49: 115-122.
- Kunze, I., Ebneth, M., Heim, U., Geiger, M., Sonnewald, U. and Herbers, K. (2001). 2-Deoxyglucose resistance: A novel selection marker for plant transformation. *Mol Breed* 7: 221-227.
- Lamblin, F., Aime, A., Hano, C., Roussy, I., Domon, J. M., Droogenbroeck, B. V. and Laine, E. (2007). The use of the phosphomannose isomerase gene as alternative selectable marker for *Agrobacterium*-mediated transformation of flax (*Linum usitatissimum*). *Plant Cell Rep.* 26:765–772.

- Lee, B. T. and Matheson, N. K. (1984). Phosphomannoisomerase and Phosphoglucoisomerase in seeds of *Cassia Coluteoides* and some other legumes that synthesize galactomannan. *Phytochemistry* 23: 983-987.
- Leduc, N., Matthys-Rochon, E., Rougier, M., Mogensen, L., Holm, P., Magnard, J. and Dumas, C. (1996). Isolated maize zygotes mimic *in vivo* embryogenic development and express microinjected genes when cultured *in vitro*. *Dev. Biol.* 177: 190-203.
- Li, H. Q., Kang, P. J., Li, M. L. and Li, M. R. (2007). Genetic transformation of *Torenia Fournieri* using the PMI/Mannose selection system. *Plant Cell Tissue Organ Cult.* 90:103–109.
- Li, Z., Upadhyaya, N. M., Meena, S., Gibbs, A. J. and Waterhouse, P. M. (1997). Comparison of promoters and selectable marker genes for use in Indian rice transformation. *Mol Breed* 3: 1-14.
- Lord, S and Vovola, P. (2004). Clonal micropropagation of oil palm (*Elaeis Guineensis Jacq.*). KULIM Conference. <http://www.nbpol.com.pg/report/index.html>.
- Loughman, B. C. (1966). The mechanism of absorption and utilization of phosphate by barley plants in relation to subsequent transport to the shoot. *New Phytol* 64: 388-397.
- Lucca, P., Ye, X. and Potrykus, I. (2001). Effective selection and regeneration of transgenic rice plants with mannose as selective agent. *Mol Breeding* 7: 43-49.
- Lurquin, P. F. (1997). Gene transfer by electroporation. *Mol. Biotechnol.* 7: 5-35.
- Luthra, R., Varsha Dubey, R. K., Srivastava, A. K. and Kumar, S. (1997). Microprojectile mediated plant transformation: A bibliographic search. *Euphytica* 95: 260-294.
- Malaysian Palm Oil Council. (2009). Malaysian Palm Oil Industry. <http://www.mpoc.org.my/Malaysian%20Palm%20Oil%20Industry.aspx>. Accessed on 15 May 2009.
- Malca, I., Endo, R. M. and Long, M. R. (1967). Mechanism of glucose counteraction of inhibition of root elongation by galactose, mannose, and glucosamine. *Phytopathology* 57: 272-278.
- Masani, A. M. Y., Ho, C. L. and Parveez, G. K. A.. (2001). Construction of PHB gene expression vectors for the production of biodegradable plastics in transgenic oil palm. Proceeding of the 2001 PIPOC International Palm Oil Congress, Kuala Lumpur. Pp 694-711.
- Masani, A. M. Y. and Parveez, G. K. A. (2008). Development of transformation vectors of the production of potentially high oleate transgenic oil palm. *Electronic Journal of Biotechnology* 11: 1-9.
- Masura, S. S. (2009). Isolation and characterization of a constitutive promoter from oil palm. Msc Thesis Submitted to Universiti Kebangsaan Malaysia.

- Masura S. S; Parveez G, K. A. and Ismanizan, I. (2010). Isolation and characterization of oil palm constitutive promoter derived from ubiquitin extension protein (*Uep1*) gene. *New Biotechnol J* 27: 289-299
- Mazur, B. J. and Falco, S. C. (1989). The development of herbicide resistant crops. *Annu. Rev. Plant Physiol.* 40:441-470.
- McCabe, D. E., Swain, W. F., Martinez, B. J. and Christou, P. (1988). Stable transformation of soybean (*Glycine max*) by particle acceleration. *Bio/Technology* 6: 923-926.
- Melanson, D., Roussy, I. and Geneviève, H. (1999). The use of phosphomannose isomerase as a selectable marker to recover transgenic *Arabidopsis* plants. Abstract from 20th Annual Crown Gall Conference Nov. p. 59. Houston, Texas.
- Mengiste, T., Amedeo, P. and Paskowski, J. (1997). High-efficiency transformation of *Arabidopsis thaliana* with a selectable marker gene regulated by the T-DNA 1'promoter. *Plant J* 12: 945-948.
- Miles, J. S. and Guest, J. R. (1984). Nucleotide sequence and transcriptional start point of the phosphomannose isomerase gene (*ManA*)Of *Escherichia Coli*. *Gene* 32: 41-48.
- Miki, B. and McHugh, S. (2004). Selectable marker genes in transgenic plants: Applications, alternatives and biosafety. *J. Biotechnol.* 107: 193-232
- Min, B. W., Cho, Y. N., Song, M . J., Noh, T. K., Kim, B. K., Chae, W. K., Park, Y. S., Choi, Y. D. and Ham, C. H. (2007). Successful genetic transformation of chinese cabbage using phosphomannose isomerase as a selection marker. *Plant Cell Rep.* 26:337–344.
- Mohd Basri, W. (2009). Overview of the Malaysian oil palm industry 2008. http://econ.mpopb.gov.my/economy/EID_web.htm.
- Murakami, T., Anzai, H., Imai, S., Satoh, A., Nagaoka, K. and Thompson, C. J. (1986). The bialaphos biosynthetic genes of *Streptomyces Hygroscopicus*: molecular cloning and characterization of the gene cluster. *Mol. Gen. Genet.* 205:42-50.
- Murashige, T. and Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant.* 15:473–497.
- Murata, T. (1975). Studies on the Phosphomannose Isomerase of *Amorphophallus konjac* C. Koch I. Its isolation and some enzymatic properties. *Plant Cell Physiol* 16:953-961.
- Na'imatalapidah, A. M. and Parvez, G. K. A. (2007). Evaluation of green fluorescence protein (GFP) as a selectable marker for oil palm transformation via transient expression. *Asia Pacific Journal Of Molecular Biology And Biotechnology*. 15, 1-8.

- Negrotto, D., Jolley, M., Beer, S., Wenck, A.R. and Hansen, G. (2000). The use of phosphomannose-isomerase as a selectable marker to recover transgenic maize plants (*Zea Mays L.*) via transformation. *Plant Cell Rep* 19: 798-803.
- Norelli, J. L. and Aldwinckle, H. S. (1993). The role of aminoglycoside antibiotics in the regeneration and selection of neomycin phosphotransferase-transgenic apple tissue. *J. Am. Soc. Hortic. Sci.* 118:311-316.
- Olczak, M. and Watorek, W. (2000). Structural analysis of N-Glycans from yellow lupine (*Lupinus Luteus*) seed diphosphonucleotide phosphate phosphodiesterase. *Biochem Et Biophys Acta* 1523: 236-245.
- O'Kennedy, M. M., Burger, J. T. and Botha, F. C. (2004). Pearl Millet transformation system using the positive selectable marker gene phosphomannose isomerase. *Plant Cell Rep* 22(9): 684-690.
- Ow, D., Wood, K., Deluca, M., Dewet, J., Helinski, D. and Howell, S. (1986). Transient and stable expression of the firefly luciferase gene in plant cells and transgenic plants. *Science* 234:856-859.
- Parveez, G. K. A., Chowdhury, M. K. U. and Saleh, N. M. (1996). Determination of minimal inhibitory concentration of selection agents for oil palm (*Elaeis Guineensis Jacq.*) transformation. *Asia Pacific Journal Of Molecular Biology and Biotechnology*. 4: 219-228.
- Parveez, G. K. A., Chowdhury, M. K. U. and Saleh, N. M. (1997). Physical parameters affecting transient GUS gene expression in oil palm (*Elaeis Guineensis Jacq.*) using the biolistic device. *Industrial Crops And Products*. 6, 41-50.
- Parveez, G. K. A. (1998). Optimization of parameters involved in the transformed of oil palm using biolistic method. Ph.D. Thesis Submitted To Universiti Putra Malaysia. 243 Pages.
- Parveez, G. K. A., Chowdhury, M. K. U. and Saleh, N. M. (1998). Biological parameters affecting transient GUS gene expression in oil palm (*Elaeis Guineensis Jacq.*) embryogenic calli via microprojectile bombardment. *Industrial Crops and Products*. 8: 17-27.
- Parveez, G. K. A., Sambanthamurthi, R., Siti Nor Akmar, A., Rasid, O., Masri, M. M. and Cheah, S. C. (1999). Production of transgenic oil palm-Current success and future considerations. Proceedings of the 1999 PORIM International Palm Oil Congress (Agriculture), ed. Ariffin D, Chan, K W and Sharifah, S.R. S. A. pp 3-13.
- Parveez, G. K. A. (2000). In '*Molecular Biology Of Woody Plants*'. (Eds. S.M. Jain and S.C. Minocha). Vol. 2, Pp. 27-350. (Kluwer Academic Publishers).
- Parveez, G. K. A., Masnita, M. M., Alizah, Z. A., Na'imatalapidah, A. M., Masani A. M.Y., Haliza, H. F., Omar, R., and Suan-Choo, C. (2000). Transgenic oil palm: Production and Projection. *Biochemical Society Transaction*. 28(6): 969-972.

- Parveez, G. K. A. (2003). Novel products from transgenic oil palm. *Agriculture and Biotechnology Network* 5: 113.
- Parveez, G. K. A., Abrizah, O., Masani, A. M. Y., Umi Salamah, R., Sambanthamurthi, R., Bahariah, B., Nur Fariza, M. S., Yuen, L. H., Nor Hanin, A., Ahmad Tarmizi, H., Zamzuri, I., Cheah, S. C., Ahmad Kushairi, D., Mohd Basri, W., York, G. and Yeong Bae, J. (2005). Value addition of oil palm through genetic engineering. Proceeding Conference Biotechnology Plant Comodities. 198-209.
- Parveez, G. K. A., Na'imatuapidah, A. M., Alizah, Z. and Rasid, A. O. (2007). Determination of minimal inhibitory concentration of selection agents for selecting transformed immature embryos of oil palm. *Asia Pacific Journal of Molecular Biology and Biotechnology*. 15(3): 133-146.
- Parveez, G. K. A. and Na'imatulapida, A. M. (2008). Factors affecting green fluorescence protein (GFP) gene expression in oil palm after microprojectile bombardment. *Journal of Oil Palm Research* 20: 494-506.
- Parveez, G. K. A., Bahariah, B., Nor Hanin, A., Masani, A. M. Y., Rasid, O. A., Ahmad Tarmizi, H., Zamzuri, I., Arif, A. M. A., Kushairi, A. D., York, G., Jo, B. Y. and Sinskey, J. A. (2008). Transformation of PHB and PHBV genes driven by maize ubiquitin promoter into oil palm for the production of biodegradable plastics. *Journal of Oil Palm Research Special Issue* 2: 77-86.
- Penna, S., Sagi, L. and Swennen, R. (2002). Positive selectable marker genes for routine plant transformation. *In Vitro Cell. Dev. Biol. Plant* 38:125-128.
- Perl, A., Galili, S., Shaul, O., Ben-Tzvi, I. and Galili, G. (1993). Bacterial dihydrolipicollate synthase and desensitized aspartic kinase: two novel selectable markers for plant transformation. *Biotechnology* 11: 715–718.
- Permyakova, N., Shumnyi, V. and Deineko, E. (2009). *Agrobacterium*-mediated transformation of plants: Transfer of vector DNA fragments in the plant genome. *Russian Journal of Genetics* 45: 266-275.
- Privalle, L. S., Wright, M., Reed, J., Hansen, G., Dawson, J., Dunder, E. M., Chang, Y. F., Powell, M. L. and Meghji, M. (2000). Phosphomannose Isomerase, a novel selectable plant selection system: mode of action and safety assessment. In Fairbairn C, Scoles G, Mchughen A, Eds, *Proceedings Of The 6th International Symposium On The Biosafety Of Genetically Modified Organisms*, Saskatoon, Canada, University Extension Press, Univ. Saskatchewan, Pp 171-178.
- Privalle, L. S. (2002). Phosphomannose Isomerase, a novel plant selection system: potential allergenicity assessment. *Ann N Y Acad Sci*. 964:129–138.
- Proudfoot, A. E. I., Turcatti, G., Wells, T.N.C., Payton, M. A. and Smith, D.J. (1994). Purification, cDNA cloning and heterologous expression of human phosphomannose isomerase. *Eur J Biochem* 219:415-423.
- Paszkowski, J., Shillito, R. D., Saul, M. W., Mandak, V., Hohn, T., Hohn, B. and Potrykus, I. (1984). Direct gene transfer to plants. *EMBO J*. 3:2717-2722.

- Puonti-Kaerlas, J., Klöti, A. and Potrykus, I. (1999). Biotechnological Contributions To Food Security With Cassava And Rice. *Plant Biotechnol.* 16: 39 – 48.
- Raberchault, H., Ahee, J. and Guenin, G. (1970). Colonies cellulaires et formes embryos *in vitro* à partir de cultures d'embryons de palmier à huile (*Elaeis guineensis* Jacq.) à l'aide de cultures de tissus foliaires. *Comptes Rendus de l'Académie des Sciences Paris Série. D*283:1735–1737.
- Raemakers, C. J. J. M., Sofiari, E., Taylor, N., Henshaw, G., Jacobsen, E. and Visser, R. G. F. (1996). Production of transgenic cassava (*Manihot Esculenta* Crantz) plants by particle bombardment using luciferase activity as selection marker. *Mol. Breed.* 2: 339–349.
- Rajanaidu, N., Jailani, B. S., Cheah, S. C. and Ahmad Kushairi, D. (1993). Oil palm breeding: Current issues and future developments. Proc. 1993 PORIM National Oil palm conference: 1-29.
- Ramesh, S. A., Kaiser, B. N., Franks, T., Collins, G. and Sedgley, M. (2006). Improved methods in *Agrobacterium*-mediated transformation of almond using positive (Mannose/*pmi*) or negative (Kanamycin resistance) selection-based protocols. *Plant Cell Rep.* 25:821–828.
- Ramessar, K., Peremarti, A., Gomez-Galera, S., Naqvi, S., Moralejo, M., Munoz, P., Capell, T. and Christou, P. (2007). Biosafety and risk assessment framework for selectable marker genes in transgenic crop plants: a case of the science not supporting the politics. *Transgenic Res.* 16: 261-280.
- Reed, J., Privalle, L., Powell, M. L., Meghji, M., Dawson, J., Dunder, E., Suttie, J., Wenck, A., Launis, K., Kramer, C., Chang, Y. F., Hansen, G. and Wright, M. (2001). Phosphomannose Isomerase: An efficient selectable marker for plant transformation. *In Vitro Cell. Dev. Biol. Plant* 37: 127-132.
- Sambanthamurthi, R., Parveez, G. K. A. and Cheah, S. C. (2000). Genetic engineering of oil palm. In: Yusof, B., Jailani, B S and Chan, K W. (eds.). Advances in Oil Palm Research, pp. 284-331. Kuala Lumpur: MPOB.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). Molecular Cloning: A laboratory manual, 2nd edn. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Sanford, J. C., Klein, T. M., Wolf, E. D. and Allen, N. (1987). Delivery of substances into cells and tissues using a particle bombardment process. *Journal of Particulate Science and Technology*. 5: 27-37.
- Sawahel, W. A. (1994). Transgenic plants: performance, release and containment. *World J. Microbiol. Biotech.* 10:139-144.
- Schmidt, M., Arnold, W., Niemann, A., Kleickmann, A. and Pühler, A. (1992) The *Rhizobium meliloti pmi* gene encodes a new type of phosphomannose isomerase. *Gene* 122:35-43.

- Schnorf, M., Neuhaus-Url, G., Galli, A., Potrykus, I. and Neuhaus, G. (1991). An improved approach for transformation of plant cells by microinjection: molecular and genetic analysis. *Transgenic Res.* 1: 12-30.
- Schöpke, C., Taylor, N., Carcamo, R., Konan, N. K., Marmey, P., Henshaw, G. G., Beachy, R. N. and Fauquet, C. (1996). Regeneration of transgenic cassava plants (*Manihot Esculenta* Crantz) from microbombarded embryogenic suspension cultures. *Nature Biotechol.* 14: 731-735.
- Shah, D. M., Horsch, R. B., Klee, H. J., Kishore, G. M., Winter, J. A., Turner, N.E., Hironaka, C. M., Sanders, P. R., Gasser, C. S., Aykent, S., Siegel, N. R., Rogers, S. G. and Fraley, R. T. (1986). Engineering herbicide tolerance in transgenic plants. *Science* 233: 478-481 .
- Shillito, R. (1999). Methods of genetic transformation: electroporation and polyethylene glycol treatment. p9-22. In: Vasil I (ed.) Molecular improvement of cereal crop. Kluwer, Dordrecht, The Netherlands.
- Shinabarger, D., Berry, A., May, T. B., Rothmel, R., Fialho, A. and Chakrabarty, A. M. (1991). Purification and characterization of phosphomannose isomerase-guanosine diphospho-D-mannose pyrophosphorylase. *J Biol Chem* 266:2080-2088.
- Shrawat, A. K., Becker, D. and Lorz, H. (2007). *Agrobacterium tumefaciens*-mediated genetic transformation of barley (*Hordeum vulgare* L.) *Plant Science* 172: 281-290.
- Slein, M. W. (1950). Phosphomannose isomerase. *J Biol Chem* 186:753-761.
- Smith, D. J., Proudfoot, A., Friedli, L., Klig, L. S., Paravicini, G. and Payton, M. A. (1992). PMI40, an intron-containing gene required for early steps in yeast mannosylation. *Mol. Cell. Biol.* 12: 2924-2930.
- Sogek, A. K. (1998). Stages in the vegetative propagation of oil palm, (*Elaeis guineensis*) Jacq. through tissue culture. *Journal of Oil Palm Research* 10 (2): 1-9.
- Songul, G., Ekrem, G., Rajvinder, K., Joshua, W., Ling, M., Han-Qi, T. and Peggy, G. L. (2009). Efficient, reproducible *Agrobacterium*-mediated transformation of sorghum using heat treatment of immature embryos. *Plant Cell Reports.* 28: 429-444.
- Sonnewald, U., and Ebneth, M. (1998). 2-Deoxyglucose-6-phosphate (2-DOG-6-P) phosphatase sequences as selection markers in plants. WO 98/45456.
- Staritsky, G. (1970). Tissue culture of oil palm (*Elaeis guineensis*) as a tool for its vegetative propagation. *Ephytica.*19:288–292. doi: 10.1007/BF01904205.
- Stein, J. C. and Hansen, G. (1999). Mannose induces an endonuclease responsible for DNA laddering in plant cell. *Plant Physiol.* 121:1-9.
- Stoop, J. M. H., Williamson, J. D. and Pharr, D. M. (1996) Mannitol metabolism in plants: A method for coping with stress. *Trends Plant Sci* 1:139-144.

- Strahl-Bolsinger, S., Gentzsch, M. and Tanner, W. (1999). Protein O-Mannosylation. *Biochem Et Biophys Acta* 1426: 297-307.
- Svab, Z., Hajdukiewicz, P. and Maliga, P. (1990a). Stable transformation of plastics in higher plants. *Proc. Natl. Acad. Sci. USA* 87: 8562-8530.
- Svab, Z., Harper, E. C., Jones J. D. G. and Maliga, P. (1990b). Aminoglycoside-3.-adenyltransferase confers resistance to spectinomycin and streptomycin in *Nicotiana tabacum*. *Plant Mol. Biol.* 14: 197-206.
- Thompson, J. A., Drayton, P. R., Frame, B. R., Wang, K, and Dunwell, J. M. (1995). Maize transformation utilizing silicon carbide whiskers; a review, *Euphytica* 85: 75-80.
- Todd, R. and Tague, B. W. (2001). Phosphomannose Isomerase: A versatile selectable marker for *Arabidopsis Thaliana* germ-line transformation. *Plant Molecular Biology Reporter*. 19: 307-319.
- Tzfira, T. and Citovsky, V. (2000). From host recognition to T-DNA integration: the function of bacterial and plant genes in the *Agrobacterium* - plant cell interaction. *Molecular Plant Pathology*, 1 (4): 201-212.
- Ursin, V. M. (1996). Aldehyde dehydrogenase selectable markers for plant transformation. WO 96/12029.
- Vain, P., De Buyser, J., Bui Trang, V., Haicour, R. and Henry, Y. (1995). Foreign gene delivery into monocotyledonous species. *Biotechnol. Adv.* 13: 943-947.
- Van Den Elzen, P. J. M., Townsend, J., Lee, K. Y. and Bedbrook, J. R. (1985). A chimaeric hygromycin resistance gene as a selectable marker in plant cells. *Plant Mol. Biol.* 5: 299-302.
- Vasil I. K. (1994) Molecular improvement of cereals. *Plant Mol. Biol.* 25:925-937.
- Vasil, V., Srivastava, V., Castillo, A. M., Fromm, M. E. and Vasil, I. K. (1993). Rapid production of transgenic wheat plants obtained by direct bombardment of cultured immature embryos. *Bio/Technology* 11: 1553-1558.
- Vasil, V., Castillo, A. M., Fromm, M. E. and Vasil, I. K. (1992). Herbicide resistant transgenic wheat plants obtained by microprojectile bombardment of regenerable embryogenic cells. *Bio/Technology* 10: 667-674.
- Waldron, C., Murphy, E. B., Roberts, J. L., Gustafson, G. D., Armour, S. L. and Malcolm, S. K. (1985). Resistance to hygromycin B: A new marker for plant transformation studies. *Plant Mol. Biol.* 5:103-108.
- Wang, A. S., Evans, R. A., Altendorf, P. R., Hanten, J. A., Doyle, M. C. and Rosichan, J. L. (2000). A mannose selection system for production of fertile transgenic maize plants from protoplasts. *Plant Cell Rep.* 19: 654-660.
- Wheeler, G. L., Jones, M. A. and Smirnoff, N. (1998) The biosynthetic pathway of vitamin C in higher plants. *Nature* 393:365-369.

- Wilmink, A. and Dons, J. J. M. (1993). Selective agents and marker genes for use in transformation of monocotyledonous plants. *Plant Mol. Biol. Reporter*. 11: 165-185.
- Wright, M., Dawson, J., Dunder, E., Suttie, J., Reed, J., Kramer, C., Chang, Y., Novitzky, R., Wang, H. and Artim-Moore, L. (2001). Efficient biolistic transformation of maize (*Zea mays* L.) and wheat (*Triticum aestivum* L.) using the phosphomannose isomerase gene, PMI, as the selectable marker. *Plant Cell Rep* 20:429-436.
- Zai-Song, D., Ming, Z., Yu-Xiang, J., Liang-Bi, L. and Ting-Yun, K. (2006). Efficient *Agrobacterium*-mediated transformation of rice by Phosphomannose Isomerase/Mannose selection. *Plant Molecular Biology Reporter* 24: 295–303.
- Zamzuri, I., Mohd Arif, S., Rajanaidu, N. and Rohani, O. (1999). Commercial feasibility of clonal oil palm planting material production. *PORIM Occasional Paper* No. 40, Pp 52.
- Zhang, P. and Puonti-Kaerlas, J. (2000). PIG-mediated cassava transformation using positive and negative selection. *Plant Cell Rep* 19:1041–1048.
- Zhang, P., Potrykus, I. and Puonti-Kaerlas, J. (2000). Efficient production of transgenic cassava using negative and positive selection. *Transgenic Res.* 9: 405-415.
- Zhao, F. Y., Li, Y. F. and Xu, P. (2006a). *Agrobacterium*-mediated transformation of cotton (*Gossypium hirsutum* L. cv. Zhongmian 35) using glyphosate as a selectable marker. *Biotechnology Letters* 28:1199-1207.
- Zhao, J., Jin, Z. Q. and Xu, B. (2006b). Progress on *Agrobacterium*-mediated transformation of banana. Publication of Medicine, National Center for Biotechnology Information (12): 1619-1626.
- Zhu, Y. J., Agbayani, R., Mccaffery, H., Albert, H. H. and Moore, P. H. (2005). Effective selection of transgenic papaya plants with the PMI/Man selection system. *Plant Cell Rep.* 24(7): 426-432.