

## References

- Allan, J. & Robinson, R. (1924), An accessible derivative of chromonol *J. Chem. Soc., Trans.*, 125, 2192.
- Arkoudis, E.; Lykakis, I. N.; Gryparis, C.; Stratakis, M. (2009) Biomimetic synthesis of dimeric metabolite Acremine G via a highly regioselective and stereoselective Diels-Alder reaction. *Org Lett.* 11, 2988-91.
- Ares, J. J., Outt, P. E., Randall, J. L., Murray, P. D., Weisshaar, P. S., O'Brien, L. M., *et al.* (1995). Synthesis and biological evaluation of substituted flavones as gastroprotective agents. *J. Med. Chem.*, 38, 4937-4943.
- v. Auwers, K. & Muller, K. (1908) *Ber. Dtsch. Chem. Ges.*, 41, 4233.
- Awuah, E., & Capretta, A. (2009). Access to flavones via a microwave-assisted, one-pot Sonogashira-carbonylation-annulation reaction. *Org. Lett.*, 11, 3210-3213.
- Baker, W. (1933). Molecular rearrangement of some *o*-acyloxyacetophenones and the mechanism of the production of 3-acylchromones. *J. Chem. Soc.*, 1381-1389.
- Beecher, G. R. (2003). Overview of dietary flavonoids: nomenclature, occurrence and intake. *J Nutr*; 133, 3248S-3254S.
- Bianco, A., Cavarischia, C., Farina, A., Guiso, M. & Marra., C. (2003). A new synthesis of flavonoids via Heck reaction. *Tetrahedron Lett.*, 44, 9107-9109
- Bois, F.; Beney, C.; Mariotte, A. M.; Boumendjel, A. (1999). A one-step synthesis of 5-hydroxyflavones . *Synlett*. 1480.
- Boots, A. W., Wilms, L. C., Swennen, E. L., Kleinjans, J. C., Bast, A., & Haenen, G. R. (2008). *In vitro* and *ex vivo* anti-inflammatory activity of quercetin in healthy volunteers. *Nutrition*, 24, 703-710.
- Broadley, K. J. & Kelly, D. R. (2001) Muscarinic receptor agonists and antagonists. *Molecules* 6, 142-193.
- Caltagirone S., Rossi C., Poggi A., *et al.* (2000). Flavonoids apigenin and quercetin inhibit melanoma growth and metastatic potential. *Int. J. Cancer*, 87, 595–600.

Chee, C. F.; Abdullah, I.; Buckle, M. J. C.; Rahman, N. A. (2010) An efficient synthesis of ( $\pm$ )-panduratin A and ( $\pm$ )-isopanduratin A, inhibitors of dengue-2 viral activity. *Tetrahedron Lett.* 51, 495-498.

Cheenpracha, S.; Karalai, C.; Ponglimanont, C.; Subhadhirasakul, S.; Tewtrakul, S. (2006) Anti-HIV-1 protease activity of compounds from *Boesenbergia pandurata*. *Bioorg. Med. Chem.* 14, 1710-4.

Chung, L.Y., Yap, K.F., Mustafa, M.R., Goh, S.H. & Imiyabir, Z. (2005) Muscarinic receptor activity of some Malaysian plant species, *Pharm. Biol.* 43, 672-682.

Chung, L.Y., Yap, K.F., Mustafa, M.R., Goh, S.H. and Imiyabir, Z. (2008) Muscarinic receptor binding activity of polyoxygenated flavones from *Melicope subunifoliolata*. *Phytochem.* 69, 1548-1554.

Cong, H.; Ledbetter, D.; Rowe, G. T.; Caradonna, J. P.; Porco, J. A., Jr. (2008) Electron transfer-initiated Diels-Alder cycloadditions of 2'-hydroxychalcones. *J. Am. Chem. Soc.* 130, 9424–9425.

Corbett, J. L.; Weavers, R. T. (2008) Aluminosilicate catalysis of chalcone Diels–Alder reactions. *Synth. Commun.* 38, 489–498.

Costantino, L.; Rastelli, G.; Albasini, A.. (1996) A rational approach to the design of flavones as xanthine oxidase inhibitors. *Eur. J. Med. Chem.* 31, 693.

Cotelle, N. (2001). Role of flavonoids in oxidative stress. *Curr. Top. Med. Chem.*, 1, 569-590.

Damas J., Bourdon V., Remacle-Volon G., Lecomte J. (1985). Pro-inflammatory flavonoids which are inhibitors of prostaglandin biosynthesis. *Prostaglandins Leukot Med.* 19, 11–24.

Dat, N. T.; Jin, X.; Lee, K.; Hong, Y. S.; Kim, Y. H.; Lee, J. J. (2009) Hypoxia-inducible factor-1 inhibitory benzofurans and chalcone-derived Diels-Alder adducts from *Morus* species. *J. Nat. Prod.* 72, 39-43.

de Groot H. (1994). Reactive oxygen species in tissue injury. *Hepatogastroenterology*, 41, 328–32.

Eddarir, S., Cotelle, N., Bakkour, Y & Rolando, C. (2003). An efficient synthesis of chalcones based on the Suzuki reaction. *Tetrahedron Lett.*, 44, 5359-5363

Forkmann, G. (1991). Flavonoids as flower pigments: the formation of the natural spectrum and its extension by genetic engineering; Blackwell: Berlin, ALLEMAGNE, Vol. 106.

Fotsis T., Pepper M.S., Aktas E., et al. (1997). Flavonoids, dietary-derived inhibitors of cell proliferation and in vitro angiogenesis. *Cancer Res.* 57, 2916–21.

Ganguly, A. K.; Kaur, S.; Mahata, P. K.; Biswas, D.; Pramanik, B. N.; Chan, T. M. (2005). Synthesis and properties of 3-acyl- $\gamma$ -pyrones, a novel class of flavones and chromones. *Tetrahedron Lett.* 46, 4119.

Ganguly, A. K.; Mahata, P. K.; Biswas, D. (2006) Synthesis of oxygen heterocycles. *Tetrahedron Lett.* 47, 1347.

Gryglewski R.J., Korbut R., Robak J., Swies J. (1987). On the mechanism of antithrombotic action of flavonoids. *Biochem Pharmacol.*, 36, 317–22.

Gu, J. Q.; Park, E. J.; Vigo, J. S.; Graham, J. G.; Fong, H. H. S.; Pezzuto, J. M.; Kinghorn, A. D. (2002) Activity-guided isolation of constituents of *Renealmia nicolaoides* with the potential to induce the phase II enzyme quinone reductase. *J. Nat. Prod.* 65, 1616–1620.

Haddach, M. & McCarthy, J. R. (1999). A new method for the synthesis of ketones: The palladium-catalyzed cross-coupling of acid chlorides with arylboronic acids. *Tetrahedron Lett.* 40, 3109-3112

Halliwell, B. & Gutteridge, J. M. C. (1998). In *Free Radicals in Biology and Medicine*; Oxford University Press: Oxford.

Hanasaki Y, Ogawa S, Fukui S. (1994). The correlation between active oxygens scavenging and antioxidative effects of flavonoids. *Free Radic. Biol. Med.*, 16, 845–50.

Harborne, J. B. (1994). In The flavonoids: Advances in Research Since 1986: Chapman and Hall: London.

Harrington, P. J.; Hegedus, L. S.; McDaniel, K. F. (1987) Palladium-catalyzed reactions in the synthesis of 3- and 4-substituted indoles. 2. Total synthesis of the *N*-acetyl methyl ester of ( $\pm$ )-clavicipitic acids. *J. Am. Chem. Soc.* 109, 4335–38.

Havsteen B. (1983). Flavonoids, a class of natural products of high pharmacological potency. *Biochem Pharmacol.* 32, 1141-8.

Hertog M.G., Feskens E.J., Hollman P.C., Katan M.B., Kromhout D. (1993). Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study. *Lancet* 342, 1007–11.

Hertog M.G., Kromhout D., Aravanis C., et al. (1995). Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study. *Arch. Intern. Med.*, 155, 381–6.

Hoang, D. M.; Ngoc, T. M.; Dat, N. T.; Ha do, T.; Kim, Y. H.; Luong, H. V.; Ahn, J.S.; Bae, K (2009) Protein tyrosine phosphatase 1B inhibitors isolated from *Morus bombycis*. *Bioorg Med Chem Lett*. 19, 6759-61.

Hoesch K., (1915). Eine Neue Synthese aromatischer Ketone. I. Darstellung einiger Phenol-ketone. *Chem. Ber.*, 48, 1122-1133.

Houben J., (1926). *ibid.*, 59, 2878.

Huk I., Brovkovich V., Nanobash V.J., et al. (1998). Bioflavonoid quercetin scavenges superoxide and increases nitric oxide concentration in ischaemia-reperfusion injury: an experimental study. *Br. J. Surg.*, 85, 1080–5.

Iwase, Y., Takemura, Y., Ju-ichi, M., Mukainaka, T., Ichiishi, E., Ito, C., et al. (2001). Inhibitory effect of flavonoid derivatives on Epstein-Barr virus activation and two-stage carcinogenesis of skin tumors. *Cancer Lett.*, 173, 105-109.

Jain, P. K.; Makrandi, J. K.; Grover, S. K. (1982) A Facile Baker-Venkataraman Synthesis of Flavones using Phase Transfer Catalysis. *Synthesis*. 221.

Jin, C. H., Shin, E. J., Park, J. B., Jang, C. G., Li, Z., Kim, M. S., et al. (2009). Fustin flavonoid attenuates beta-amyloid (1-42)-induced learning impairment. *J Neurosci Res*, 87, 3658-3670.

Jung, E. M.; Lee, Y. R. Bull. (2008) First concise total syntheses of biologically interesting nicolaoidesin C, crinatusin C1, and crinatusin C2. *Korean Chem. Soc.* 29, 1199–1204.

Kaul T.N., Middleton E. Jr, Ogra P.L. (1985). Antiviral effect of flavonoids on human viruses. *J Med. Virol.*, 15, 71–9.

Kantam, M. L., Ranganath, K. V. S., Sateesh, M., Kumar, K. B. S. & Choudary, B. M. (2005). Friedel-Crafts acylation of aromatics and heteroaromatics by beta zeolite. *J. Mol. Catal. A: Chem.*, 225, 15-20.

Kitamura, K., Honda, M., Yoshizaki, H., Yamamoto, S., Nakane, H., Fukushima, M., et al. (1998). Baicalin, an inhibitor of HIV-1 production in vitro. *Antiviral Res*, 37, 131-140.

Knek P., Jarvinen R., Seppanen R., et al. (1997). Dietary flavonoids and the risk of lung cancer and other malignant neoplasms. *Am. J. Epidemiol.*, 146, 223–30.

Korkina L.G., Afanas'ev I.B. (1997). Antioxidant and chelating properties of flavonoids. *Adv Pharmacol*, 38, 151–63.

Lagiou, P., Samoli, E., Lagiou, A., Tzonou, A., Kalandidi, A., Peterson, J., et al. (2004). Intake of specific flavonoid classes and coronary heart disease-a case-control study in Greece. *Eur J Clin Nutr*, 58, 1643-1648.

Larson, R. A. (1988). The antioxidants of higher plants. *Phytochemistry*, 27, 969-978.

Lee Y. K.; Tan S. K.; Wahab, H. A.; Yusof, R.; Rahman, N. A. (2007) Nonsubstrate based inhibitors of dengue virus serine protease: A molecular docking approach to study binding interactions between protease and inhibitors. *AsPac J. Mol. Biol. Biotechnol.* 15, 53-59.

Liang, Y. C., Tsai, S. H., Tsai, D. C., Lin-Shiau, S. Y., & Lin, J. K. (2001). Suppression of inducible cyclooxygenase and nitric oxide synthase through activation of peroxisome proliferator-activated receptor-gamma by flavonoids in mouse macrophages. *FEBS Lett*, 496, 12-18.

Loke, W. M., Proudfoot, J. M., Stewart, S., McKinley, A. J., Needs, P. W., Kroon, P. A., et al. (2008). Metabolic transformation has a profound effect on anti-inflammatory activity of flavonoids such as quercetin: lack of association between antioxidant and lipoxygenase inhibitory activity. *Biochem Pharmacol*, 75, 1045-1053.

Looker, J. H.; Hanneman, W. W. (1962) Physical and chemical properties of hydroxyflavones. I. Infrared absorption spectra of monohydroxyflavones and their *O*-methyl and *O*-acetyl derivatives. *J. Org. Chem.* 27, 381.

Looker, J. H.; Hanneman, W. W. (1962) Physical and chemical properties of hydroxyflavones. II. 3-aroyl-5-hydroxyflavones. Synthetic and infrared spectral studies. *J. Org. Chem.* 27, 3261.

Liu, B., Yang, J., Wen, Q., & Li, Y. (2008). Isoliquiritigenin, a flavonoid from licorice, relaxes guinea-pig tracheal smooth muscle *in vitro* and *in vivo*: role of cGMP/PKG pathway. *Eur J Pharmacol*, 587, 257-266.

Mahal, H.S.; Venkataraman, K. (1933). A synthesis of flavones at room temperature. *Curr. Sci.* 2, 214.

Mahal, H. S.; Venkataraman, K. (1934). Synthetical experiments in the chromone group. Part XIV. The action of sodamide on 1-acyloxy-2-acetonaphthones. *J. Chem. Soc.*, 1767-1769.

Marais, J. P., Ferreira, D., & Slade, D. (2005). Stereoselective synthesis of monomeric flavonoids. *Phytochem.*, 66, 2145-2176.

Middleton, E., Jr., Kandaswami, C., & Theoharides, T. C. (2000). The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease, and cancer. *Pharmacol Rev*, 52, 673-751.

Minassi, A., Giana, A., Ech-Chahad, A., & Appendino, G. (2008). A regiodivergent synthesis of ring A prenylflavones. *Org. Lett.*, 10, 2267-2270.

Morikawa, T.; Funakoshi, K.; Ninomiya, K.; Yasuda, D.; Miyagawa, K.; Matsuda, H.; Yoshikawa, M. (2008) Medicinal foodstuffs. XXXIV.<sup>1</sup> Structures of new prenylchalcones and prenylflavanones with TNF- $\alpha$  and aminopeptidase N inhibitory activities from *Boesenbergia rotunda*. *Chem. Pharm. Bull.* 56, 956-962.

Mruk, J. S., Webster, M. W., Heras, M., Reid, J. M., Grill, D. E., & Chesebro, J. H. (2000). Flavone-8-acetic acid (Flavonoid) profoundly reduces platelet-dependent thrombosis and vasoconstriction after deep arterial injury *In vivo*. *Circulation*, 101, 324-328.

Mursu, J., Voutilainen, S., Nurmi, T., Tuomainen, T. P., Kurl, S., & Salonen, J. T. (2008). Flavonoid intake and the risk of ischaemic stroke and CVD mortality in middle-aged Finnish men: the Kuopio Ischaemic Heart Disease Risk Factor Study. *Br J Nutr*, 100, 890-895.

Narender, T. & Papi Reddy, K. (2007). A simple and highly efficient method for the synthesis of chalcones by using borontrifluoride-etherate. *Tetrahedron Lett.* 48, 3177-3180.

Nijveldt, R. J., van Nood, E., van Hoorn, D. E., Boelens, P. G., van Norren, K., & van Leeuwen, P. A. (2001). Flavonoids: a review of probable mechanisms of action and potential applications. *Am J Clin Nutr*, 74, 418-425.

Osman H.E., Maalej N., Shanmuganayagam D., Folts J.D. (1998). Grape juice but not orange or grapefruit juice inhibits platelet activity in dogs and monkeys. *J. Nutr.*, 128, 2307-12.

Pandji, C.; Grimm, C.; Wray, V.; Witte, L.; Proksch, P. (1993) Insecticidal constituents from four species of the Zingiberaceae. *Phytochemistry*. 34, 415-419.

Rao, V. S., Paiva, L. A., Souza, M. F., Campos, A. R., Ribeiro, R. A., Brito, G. A., et al. (2003). Ternatin, an anti-inflammatory flavonoid, inhibits thioglycolate-elicited rat peritoneal neutrophil accumulation and LPS-activated nitric oxide production in murine macrophages. *Planta Med*, 69, 851-853.

Piatta, P. G. (2000). Flavonoids as antioxidants *J. Nat. Prod.* 63, 1035-42.

Rajkumar, S., & Jebanesan, A. (2008). Bioactivity of flavonoid compounds from *Poncirus trifoliata* L. (Family: Rutaceae) against the dengue vector, *Aedes aegypti* L. (Diptera: Culicidae). *Parasitol Res*, 104, 19-25.

Riva, C.; DeToma, C.; Donadd, L.; Boi, C.; Pennini, R.; Motta, G. & Leonardi, A. (1997). New DBU (1,8-diazabicyclo[5.4.0]undec-7-ene) assisted one-pot synthesis of 2,8-disubstituted 4H-1-benzopyran-4-ones. *Synthesis*. 195.

Ruhemann, S.; Stapleton, H. E. (1900) Condensation of Phenols with Esters of the Acetylene Series. Part III. Synthesis of Benzo- $\gamma$ -pyrone. *J. Chem. Soc.*, 77, 1179-1185.

Ruhemann, S. (1913) Über ein einfaches Verfahren zur Darstellung der Flavone und über die Synthese des Thioflavones. (A Simple Procedure for the Preparation of Flavones and the Synthesis of Thioflavone.) *Ber.*, 46, 2188-2197.

Sanchez, I., Gomez-Garibay, F., Taboada, J., & Ruiz, B. H. (2000). Antiviral effect of flavonoids on the dengue virus. *Phytother Res*, 14, 89-92.

Sanhueza J., Valdes J., Campos R., Garrido A., Valenzuela A. (1992). Changes in the xanthine dehydrogenase/xanthine oxidase ratio in the rat kidney subjected to ischemia-reperfusion stress: preventive effect of some flavonoids. *Res. Commun. Chem. Pathol. Pharmacol.*, 78, 211-8.

Shibata, K.; Tatsukawa, A.; Umeoka, K.; Lee, H. S.; Ochi, M. (2000) Crinatusins, bioactive Diels-Alder adducts from *Cyathocalyx crinatus*. *Tetrahedron*. 56, 8821-8824;

Sugamoto, K.; Kurogi, C.; Matsushita, Y.; Matsui, T. (2008) Synthesis of 4-hydroxyderricin and related derivatives. *Tetrahedron Lett*. 49, 6639-41.

Tan, G.H. (2004). The Synthesis of Chrysin under Houben-Hoesch Reaction Conditions., B.Sc Thesis, University of Malaya.

Tan, S. K.; Pippen, R.; Yusof, R.; Ibrahim, H.; Khalid, N.; Rahman, N. A. (2006) Inhibitory activity of cyclohexenyl chalcone derivatives and flavonoids of

fingerroot, *Boesenbergia rotunda* (L.), towards dengue-2 virus NS3 protease. *Bioorg. Med. Chem. Lett.* 16, 3337-40.

Tsopmo, A; Tene, M; Kamnaing, P; Ayafor, J.F.; Sterner, O. (1999) A new Diels–Alder-type adduct flavonoid from *Dorstenia barteri*. *J. Nat. Prod.* 62, 1432.

Tuntiwachwuttikul, P.; Pancharoen, O.; Reutrakul, V.; Byrne, L. T. (1984) (1'RS,2'SR,6'RS)-(2,6-dihydroxy-4-methoxyphenyl)-[3'-methyl-2'-(3"-methylbut-2"-enyl)-6'-phenyl-cyclohex-3'-enyl]methanone (panduratin A)-a constituent of the red rhizomers of a variety of *Boesenbergia pandurata*. *Aust. J. Chem.* 37, 449–453.

Ueng, Y. F., Shyu, C. C., Liu, T. Y., Oda, Y., Lin, Y. L., Liao, J. F., et al. (2001). Protective effects of baicalein and wogonin against benzo[a]pyrene- and aflatoxin B (1)-induced genotoxicities. *Biochem Pharmacol*, 62, 1653-1660.

Valerio, D. A., Georgetti, S. R., Magro, D. A., Casagrande, R., Cunha, T. M., Vicentini, F. T., et al. (2009). Quercetin reduces inflammatory pain: inhibition of oxidative stress and cytokine production. *J Nat Prod*, 72, 1975-1979.

van Acker S.A., Tromp M.N., Haenen G.R., van der Vijgh W.J., Bast A. (1995). Flavonoids as scavengers of nitric oxide radical. *Biochem. Biophys. Res. Commun.*, 214, 755–9.

Verma, A. K., & Pratap, R. (2010). The biological potential of flavones. *Nat Prod Rep*, 27, 1571-1593.

Wei, K.; Li, W.; Koike, K.; Nikaido, T. (2005). Cobalt(II)-Catalyzed Intermolecular Diels–Alder Reaction of Piperine. *Org. Lett.* 7, 2833-2835.

Wessely, F.; Moser, G. H. *Monatsch. Chem.* (1930). Synthese und Konstitution des Skutellareins. 56, 97–105.

Wheeler, T. S. (1963) Flavone. *Organic Syntheses*, New York: Wiley, Collect. Vol. IV, 478-481.

Win, N. N.; Awale, S.; Esumi, H.; Tezuka, Y.; Kadota, S. (2007) Bioactive secondary metabolites from *Boesenbergia pandurata* of Myanmar and their preferential cytotoxicity against human pancreatic cancer PANC-1 cell line in nutrient-deprived medium. *J. Nat. Prod.* 70,1582–1587.

Yoshio Hano, Miwa Aida, Taro Nomura and Shinichi Ueda. (1992) A novel way of determining the structure of artonin I, an optically active Diels–Alder type adduct, with the aid of an enzyme system of *Morus alba* cell cultures. *J. Chem. Soc. Chem. Commun.* 1177.