

MALOCCLUSION STATUS AND ORTHODONTIC TREATMENT
NEEDS OF 14-YEAR-OLD YEMENI ADOLESCENTS

RASHEED ABDULSALAM AL-MAQTARI

THESIS SUBMITTED IN FULFILLMENT OF THE
REQUIREMENTS FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

DEPARTMENT OF CHILDREN'S DENTISTRY AND ORTHODONTIC
FACULTY OF DENTISTRY
UNIVERSITY OF MALAYA
KUALA LUMPUR

2011

ABSTRACT

The aim of this study was to develop a national database on the prevalence, severity of malocclusion and orthodontic treatment needs among Yemeni adolescents. Other variables included were geographical zones, gender and urban-rural locations.

A multi-stage stratified random sample of 2400 subjects from 60 schools with equal proportion of male and female were selected from nine governorates covering the whole topography of Yemen (coastal, plateau, mountains, desert and islands). Measuring instruments were the conventional FDI/WHO method of occlusal traits (Baume et al., 1973; Berzoukov et al., 1979) and the Index of Orthodontic Treatment Needs (IOTN), (Brooke and Shaw, 1989; Richmond et al., 1995). Other occlusal traits included based on clinical oral examination were canine relationship and bimaxillary protrusion. Data collection was carried out by one examiner assisted by a recorder, using mouth mirror and the orthodontic ruler, with patient seating on a portable dental chair or classroom chair and natural light. Prior to the oral examination, the examiner was calibrated against a gold standard on children of the same age in Malaysia. A pilot study was again conducted on Yemeni children in Thamar a week before data collection began. All information gathered was checked for completeness and data was transferred into a laptop using the SPSS software data entry program. Analysis was carried out using the SPSS version 15 program.

Measurement of occlusal traits based on FDI/WHO objective method showed that dental discrepancies was observed in 14.6% (impacted 5.7%, congenitally absent 2.2%, retained deciduous 3.2%, missing due to extraction or trauma 2.8% and supernumerary 0.7%) of the sample examined. Crowding was observed in 53.3% of the sample, of whom 27.8% had crowding of ≥ 2 mm. Spacing was observed in 2.9% of the sample, mostly in the maxilla. Midline diastema was also found to occur in 5.4% of the sample

examined. Asymmetrical molar relationship was only observed in 11.3% of the sample, mostly of Class I/Class II relationship (9.4%). Overjet was observed in 9.1% of the sample. Only a small proportion (2%) had anterior crossbite, deepbite (10.3%), anterior openbite (4.5%), posterior openbite (2.9%), posterior crossbite (5.2%) and 2.7% scissor bite. Other occlusal traits measured clinically were partially erupted teeth (6.5%), Class II canine relationship (right 17.4%, left 18.4%), Class III canine relationship (right 1.8%, left 1.7%) and bimaxillary protrusion (9.3%).

Normative orthodontic treatment needs as assessed using the Dental Health Component criteria of (IOTN) showed that slightly less than half (44.3%) of the adolescents needed some form of orthodontic treatment (Grades 3,4 and 5). Of this a quarter (26.8%) 'definitely' needed treatment. Assessment of whether examiner or subject perceived orthodontic treatment need was made using the Aesthetic Component of IOTN. Findings showed that examiner perceived at least 28.9% of the adolescents in this study needed orthodontic treatment (18.8% 'borderline' and 10.1% 'definite' need); on the contrary subjects themselves perceived a much lower proportion (13.4%). Further analysis was carried out to compare the findings according to gender, urban-rural areas and country's topography.

In conclusion, findings showed the prevalence of malocclusion was 74.4% when measured using the conventional WHO/FDI criteria. But when measured with Index of IOTN, the prevalence of malocclusion observed was 68.2%. Although both examiner measurements of normative and aesthetic perceived need indicated a relatively high, subjects self-perceived needs was however low. These findings suggest a need for a more conservative treatment approach in dealing with malocclusion problems among Yemeni adolescents.

DECLARATION

I certify that this thesis is based on my own independent work, except where acknowledged in the text or by reference. No part of this work has been submitted for a degree or diploma to this or any other university.

Dr. Rasheed Abdulsalam Al-Maqtari

Date:

Supervisor: **Associate Professor Dr. Halimah Awang**
Department of Children's Dentistry and Orthodontic,
Faculty of Dentistry,
University of Malaya,
Kuala Lumpur,
Malaysia.

Co-Supervisor: **Professor Dr. Rahimah Abdul Kadir**
Deputy Dean,
Postgraduates, Research, Internationalisation & Industry,
Faculty of Dentistry,
Universiti Kebangsaan Malaysia,
Kuala Lumpur,
Malaysia.

Co-Supervisor: **Dr. Norzakiah Mohamed Zam Zam**
Part-time lecturer,
Department of Children's Dentistry and Orthodontic,
Faculty of Dentistry,
University of Malaya,
Kuala Lumpur,
Malaysia.

ACKNOWLEDGEMENTS

First of all, I thank “God” for inspiring me with the ability, strength and patience to perform this modest study.

My deepest gratitude and appreciation to my supervisor *Professor Dr. Rahimah Abdul Kadir* for her wisdom, encouragement, guidance, patience and continuous valuable scientific suggestions throughout the preparation of my thesis, as well as for the time she has spent reading the various drafts. Her critical commentary on my work has played a major role in both the content and presentation of my arguments.

I would like to express my sincere gratitude to my supervisor *Associate Professor Dr. Halimah Awang* for her guidance, valuable advice, continuous support, patience and valuable time throughout my work and very helpful comments.

I am most grateful to my supervisor *Dr. Norzakiah Mohamed Zam Zam* for her bright ideas, guidance and for being generous with her knowledge, experience and time in supervising and encouraging me greatly in every step along the way of this work.

My grateful thanks and praise goes to the Dean, *Professor Dr. Rosnah Zain*, and *Professor Dato’ Dr. Ishak Abdul Razak* former Dean, Faculty of Dentistry, University of Malaya for their continuous support to all the postgraduate students including myself.

I would like to express my deepest gratitude to *Associate Professor Dr. Noor Hayaty Abu Kasim*, Deputy Dean (Research and Postgraduate), Faculty of Dentistry, University of Malaya for managing all the necessities for postgraduate program.

I wish to express my most sincere appreciation to all staff and members of the Department of Children’s Dentistry and Orthodontic for their invaluable supports. Also

grateful thanks to *Dr. Marhazlinda Jamaludin*, Head of Informatic Unit, Faculty of Dentistry, University of Malaya for her statistical help and advice.

I would like to offer my appreciation to *Professor Dr. Ahmad Abdo Saleh* Vice-Rector for Postgraduate Studies and Scientific Research, Thamar University, Yemen for his support, cooperation and communication with Ministry of Health and Ministry of Education throughout my work.

My special thanks to *Dr. Mohamed Al-Tabib*, *Dr. Nasr Al-Qudimi* and *Lutf Al-Yemeni* for their help in organising the survey and data collection. Also my sincere thanks to *Professor Dr. S. Richmond*, for having kindly provided some references related to this study.

I would like to thank University of Malaya, Malaysia and University of Thamar, Yemen for giving me this opportunity to continue my study and also for providing an effective environment and financial supports.

My grateful thanks and appreciation to headmasters, all the school staffs and students involved in this study for their kind help and cooperation. The author would also like to express special thanks to so many others unmentioned but certainly not forgotten.

Finally, my acknowledgement would not be completed without recognising the role of my wife *Lubna* and my beloved children *Emad*, *Bara'a* and *Khaled* for their love and patience. I should also like to express my deepest gratitude to the memory of my father, my mother and every member in my family for all that they have done for me.

CONTENTS

Title page	i
Abstract	ii
Declaration	iv
Acknowledgements	v
Contents	vii
List of figures	xiv
List of tables	xvi
List of abbreviations	xx
List of appendices	xxi
CHAPTER ONE: INTRODUCTION	1
CHAPTER TWO: REVIEW OF LITERATURE	6
2.1 Country Background	7
2.1.1 Country profile	7
2.1.2 Background history	8
2.1.3 Population	9
2.1.4 Geographical outlook	9
2.1.5 Economic development	10
2.1.6 Dental education and services in Yemen	11
2.2 Dental Occlusion	12
2.2.1 Ideal dental occlusion	12
2.2.2 Normal dental occlusion	13
2.2.3 Malocclusion	15
2.2.3.1 Aetiology of malocclusion	15
2.2.3.2 Prevalence of malocclusion	16
2.2.3.3 Psychological effect of malocclusion	17
2.3 Occlusal Traits	18
2.3.1 Dental discrepancies	18
2.3.2 Space discrepancies	21
2.3.2.1 Crowding and spacing	21
2.3.2.2 Maxillary midline diastema	24
2.3.3 Occlusal discrepancies in antero-posterior relationship	25

2.3.3.1	Overjet	25
2.3.3.2	Anterior crossbite	27
2.3.3.3	Bimaxillary protrusion	28
2.3.4	Occlusal discrepancies in vertical relationship	29
2.3.4.1	Overbite	29
2.3.4.2	Openbite	31
2.3.5	Occlusal discrepancies in transversal relationship	33
2.3.5.1	Posterior crossbite and scissor bite	33
2.4	Methods of Malocclusion Measurements	34
2.4.1	Requirements for an ideal method for malocclusion measurement	35
2.4.2	Classification of malocclusion assessment methods	36
2.4.3	Evolution of malocclusion classifications and indices	39
2.4.3.1	Angle Classification (1899)	39
2.4.3.1.1	Criticisms of Angle Classification	40
2.4.3.1.2	Modifications of Angle Classification	42
2.4.3.2	Massler and Frankel (1951)	45
2.4.3.3	Malalignment Index (1959)	45
2.4.3.4	Handicapping Labio-lingual Deviations Index, HLDI (1960)	46
2.4.3.5	Malocclusion Severity Estimate, MSE (1961)	48
2.4.3.6	Occlusal Feature Index, OFI (1961)	48
2.4.3.7	Method for Epidemiological Registration of Malocclusion, MERM (1964)	49
2.4.3.8	Treatment Priority Index, TPI (1967)	50
2.4.3.9	Handicapping Malocclusion Assessment Record, HMAR (1968)	51
2.4.3.10	Occlusal Index (1971)	52
2.4.3.11	Method for Measuring Occlusal Traits (FDI, 1973)	53
2.4.3.12	Swedish Public Health Index (1974)	55
2.4.3.13	Eismann Index (1974)	56
2.4.3.14	Indication Index (1977)	56
2.4.3.15	British Standards Institute Classification (1983)	57
2.4.3.16	Severity Index (1986)	58
2.4.3.17	Dental Aesthetic Index, DAI (1986)	58

2.4.3.18	Index of Orthodontic Treatment Need, IOTN (1989)	60
2.4.3.18.1	Dental Health Component (DHC) of IOTN	61
2.4.3.18.2	Aesthetic Component (AC) of IOTN	63
2.4.3.18.3	Validity and reliability of IOTN	66
2.4.3.19	Need for Orthodontic Treatment Index, NOTI (1992)	68
2.4.3.20	Index of Complexity Outcome and Need, ICON (2000)	69
2.4.3.21	Treatment Difficulty Index (2006)	71
2.5	Selection of Measurement Methods for this Study	73
2.6	Purpose of the Study	75
2.6.1	Aim	75
2.6.2	Specific objectives	75
	CHAPTER THREE: METHODOLOGY	77
3.1	Study Area	78
3.2	Study Design	80
3.3	Sampling	80
3.3.1	Sample size calculation	80
3.3.2	Determination of age sample	82
3.3.3	Selection of schools	83
3.3.4	Selection of sample	84
3.4	Administration of Study Conduct	85
3.4.1	Ethical approval and permissions	85
3.4.2	Instruments and equipment	85
3.4.3	Infection control	86
3.5	Measurement Tool and Criteria	87
3.5.1	Occlusal traits based on FDI/WHO method	88
3.5.1.1	Measurement of dental discrepancies	88
3.5.1.1.1	Missing permanent teeth	88
3.5.1.1.2	Supernumerary teeth	90
3.5.1.1.3	Incisors malformation	90
3.5.1.1.4	Retained deciduous teeth	90
3.5.1.2	Measurement of space discrepancies	91

3.5.1.2.1	Crowding	91
3.5.1.2.2	Spacing	92
3.5.1.2.3	Maxillary midline diastema	92
3.5.1.3	Measurement of antero-posterior occlusal relationship	92
3.5.1.3.1	Molar relationship	93
3.5.1.3.2	Overjet	96
3.5.1.3.3	Anterior crossbite	96
3.5.1.4	Measurement of vertical occlusal relationship	97
3.5.1.4.1	Overbite	97
3.5.1.4.2	Anterior openbite	97
3.5.1.4.3	Posterior openbite	98
3.5.1.5	Measurement of transversal occlusion relationship	98
3.5.1.5.1	Posterior crossbite	98
3.5.1.5.1	Scissor bite	99
3.5.2	Measurement of additional occlusal traits observed	99
3.5.2.1	Partially erupted teeth	99
3.5.2.2	Canine relationship	100
3.5.2.3	Bimaxillary protrusion	103
3.5.2.4	Incisor Classification	103
3.5.3	Measurement of orthodontic treatment need	105
3.5.3.1	Dental Health Component (DHC)	105
3.5.3.2	Aesthetic Component (AC)	106
3.5.3.2.1	Aesthetic Component evaluation by examiner	106
3.5.3.2.2	Aesthetic Component evaluation by subjects	106
3.6	Calibration	108
3.6.1	Laboratory calibration	108
3.6.2	Field clinical calibration	108
3.6.3	Calibration results	110
3.7	Pilot Study	114
3.7.1	Sampling and examination area	114
3.7.2	Duplicate examinations	115
3.7.3	Results of pilot study	115
3.8	Main Study Implementation	116
3.8.1	Examination area	116
3.8.2	General and personal information	117

3.8.3 Clinical examination	117
3.9 Statistical Analysis	118
CHAPTER FOUR: RESULTS	119
4.1 Occlusal Traits Based on FDI/WHO Method	120
4.1.1 Dental discrepancies assessment	120
4.1.1.1 Missing permanent teeth	123
4.1.1.2 Supernumerary and retained deciduous teeth	124
4.1.2 Space discrepancies assessment	125
4.1.2.1 Crowding	125
4.1.2.1.1 Maxillary crowding	126
4.1.2.1.2 Mandibular crowding	128
4.1.2.2 Spacing	130
4.1.2.2.1 Maxillary and mandibular spacing	131
4.1.2.3 Maxillary midline diastema	134
4.1.3 Occlusal Discrepancies assessment	135
4.1.3.1 Occlusal discrepancies in antero-posterior relationship	135
4.1.3.1.1 Molar relationship	135
4.1.3.1.2 Overjet	140
4.1.3.1.3 Anterior crossbite	143
4.1.3.2 Occlusal discrepancies in vertical relationship	145
4.1.3.2.1 Overbite	145
4.1.3.2.2 Anterior openbite	147
4.1.3.2.3 Posterior openbite	148
4.1.3.3 Occlusal discrepancies in transversal relationship	149
4.1.3.3.1 Posterior crossbite	149
4.1.3.3.2 Scissor bite	151
4.2 Additional Occlusal Traits Observed	152
4.2.1 Partially erupted teeth	152
4.2.2 Canine relationship	153
4.2.3 Bimaxillary protrusion	157

4.3	Orthodontic Treatment Needs Based on IOTN	158
4.3.1	Normative orthodontic treatment need assessment	158
4.3.1.1	Distribution of normative orthodontic treatment need of the sample by demographic variables	159
4.3.2	Orthodontic treatment need as measured by Aesthetic Component of IOTN	161
4.3.2.1	Orthodontic treatment need as perceived by examiner	162
4.3.2.2	Orthodontic treatment need as perceived by subjects	163
4.4	Relationship of Orthodontic Treatment Need between Examiner and Subjects	165
4.5	Malocclusion as Measured by Different Methods	166
4.5.1	Prevalence of malocclusion based on FDI/WHO method	166
4.5.2	Prevalence of malocclusion based on IOTN	168
4.5.3	Comparison between malocclusion as measured by FDI/WHO method and IOTN	171
	CHAPTER FIVE: DISCUSSION	172
5.1	General Overview of the Findings	173
5.2	Occlusal Traits Based on FDI/WHO Method	175
5.2.1	Dental discrepancies	175
5.2.1.1	Missing permanent teeth	175
5.2.1.2	Supernumerary teeth and retained deciduous teeth	177
5.2.2	Space discrepancies	178
5.2.2.1	Crowding and spacing	178
5.2.2.2	Maxillary midline diastema	180
5.2.3	Occlusal discrepancies	182
5.2.3.1	Occlusal discrepancies in antero-posterior relationship	182
5.2.3.1.1	Molar relationship	182
5.2.3.1.2	Overjet	184
5.2.3.1.3	Anterior crossbite	186
5.2.3.2	Occlusal discrepancies in vertical relationship	187
5.2.3.2.1	Overbite	187
5.2.3.2.2	Anterior openbite	188

5.2.3.2.3	Posterior openbite	189
5.2.3.3	Occlusal discrepancies in transversal relationship	190
5.2.3.3.1	Posterior crossbite	190
5.2.3.3.2	Scissor bite	191
5.3	Additional Occlusal Traits Observed	191
5.3.1	Partially erupted teeth	191
5.3.2	Canine relationship	192
5.3.3	Bimaxillary protrusion	193
5.4	Orthodontic Treatment Needs as Measured by IOTN	195
5.4.1	Normative orthodontic treatment need assessment	195
5.4.2	Orthodontic treatment need as measured by the Aesthetic Component (AC) of IOTN	198
5.4.2.1	Aesthetic Component (AC) assessment by examiner	198
5.4.2.2	Aesthetic Component (AC) assessment by subjects	199
5.5	Relationship of Orthodontic Treatment Need Assessment by Examiner and Subjects	200
5.6	Prevalence of Malocclusion as Measured by Different Methods	202
5.7	Modification of Orthodontic Measurement Tool for Yemeni Population	203
5.8	The Need for Orthodontists to Improve Orthodontic Services in Yemen	206
5.9	Limitations of This Study	207
	CHAPTER SIX: CONCLUSIONS AND RECOMMENDATIONS	209
6.1	Conclusions	210
6.2	Recommendations	212
	REFERENCES	214
	APPENDICES	232

LIST OF FIGURES

Figure	Description	Page
CHAPTER THREE: METHODOLOGY		
3.1	Distribution of nine governorates (green colour) in map of Yemen	79
3.2	Flow chart of the sample selection (giving middle zone as an example)	84
3.3	Instruments and equipment used in the study	86
3.4	Impacted lower canine clinically observed as bulging ridge	89
3.5	Clinically absent lateral incisors	89
3.6	Missing central incisor due to extraction which was confirmed from history and reduction in alveolar bone	89
3.7	Supernumerary tooth with displacement of teeth	90
3.8	Crowding of lower arch (distance between anatomical contact points)	91
3.9	Generalised spacing	92
3.10	Class I molar relationship	93
3.11	Class II molar relationship	94
3.12	Class III molar relationship	95
3.13	Anterior crossbite of upper right lateral incisor	96
3.14	Anterior openbite	97
3.15	Posterior crossbite	98
3.16	Scissor bite	99
3.17	Partially erupted lower second premolar tipped and impacted against first premolar and first molar	99
3.18	Class I canine relationship	100
3.19	Class II canine relationship	101
3.20	Class III canine relationship	102
3.21	Incisor classification relationship	104
3.22	Disposable DHC ruler used in measurements of orthodontic treatment need	106

3.23	Aesthetic component evaluation on a female subject in urban area	107
3.24	Aesthetic Component male evaluation on a male subject in rural area	107
3.25	The gold standard and recorder	109
3.26	Intra-examiners calibration	110
3.27	Position of the examiner, subject and recorder	116

CHAPTER FOUR: RESULTS

4.1	Prevalence of dental discrepancies of the sample	120
4.2	Occlusion status based on FDI/WHO method	167

LIST OF TABLES

Table	Description	Page
CHAPTER TWO: REVIEW OF LITERATURE		
2.1	Summary of studies on prevalence of malocclusion	17
2.2	Summary of studies on prevalence of impacted, congenital and supernumerary teeth	20
2.3	Summary of studies on prevalence of crowding and spacing	23
2.4	Summary of studies on prevalence of maxillary diastema	25
2.5	Summary of studies on prevalence of overjet	26
2.6	Summary of studies on prevalence of anterior crossbite	28
2.7	Summary of studies on prevalence of deep overbite	31
2.8	Summary of studies on prevalence of anterior and posterior openbite	32
2.9	Summary of studies on prevalence of posterior crossbite and scissor bite	34
2.10	Summary of epidemiological studies on malocclusion according to Angle classes assessment	44
2.11	Severity and treatment priority of malocclusion according to Treatment Priority Index (TPI)	50
2.12	Summary of studies based on normative orthodontic treatment need	62
2.13	Summary of studies of orthodontic treatment need according to subject's perception	64
2.14	Summary of studies of treatment need according to examiner assessment	65
2.15	Need, acceptability and complexity score range of Index of Complexity Outcome and Need (ICON)	70
2.16	Improvement grades of treatment according to Index of Complexity Outcome and Need (ICON)	71
CHAPTER THREE: METHODOLOGY		
3.1	Number of 14 year old intermediate school students of nine included governorates and total population of Yemen	79
3.2	Distribution of sample by zones, location (urban & rural) and gender	82

3.3	Results of inter-examiner calibration of occlusal traits	111
3.4	Crosstabulation of inter-examiner calibration of Dental Health Component of IOTN measurement	112
3.5	Crosstabulation of intra-examiner calibration of Dental Health Component of IOTN measurement	112
3.6	Crosstabulation of inter-examiner calibration of Aesthetic Component of IOTN measurement	113
3.7	Crosstabulation of intra-examiner calibration of Aesthetic Component of IOTN measurement	113
3.8	Intra-examiner calibration of malocclusion classified by Incisor, DHC and AC of IOTN (Pilot study sample)	115

CHAPTER FOUR: RESULTS

4.1	Distribution of dental discrepancies by zones	121
4.2	Distribution of dental discrepancies by location (urban & rural) and gender	122
4.3	Distribution of missing permanent teeth	123
4.4	Distribution of supernumerary, partially erupted and retained deciduous teeth	124
4.5	Crosstabulation between maxillary and mandibular crowding	125
4.6	Distribution of maxillary crowding by zones	127
4.7	Distribution of maxillary crowding by location (urban & rural) and gender	127
4.8	Distribution of mandibular crowding by zones	129
4.9	Distribution of mandibular crowding by location (urban & rural) and gender	129
4.10	Crosstabulation between maxillary and mandibular spacing	130
4.11	Distribution of maxillary spacing between zones	132
4.12	Distribution of maxillary spacing between location (urban & rural) and gender	132
4.13	Distribution of mandibular spacing between zones	133
4.14	Distribution of mandibular spacing between location (urban & rural) and gender	133
4.15	Association of maxillary diastema between demographic variables	134
4.16	Crosstabulation of right and left molar relationship classes	135

4.17	Distribution of molar relationship discrepancy in right and left sides of occlusion by zones	137
4.18	Distribution of molar relationship discrepancy in right and left sides of occlusion by location (urban & rural) and gender	138
4.19	Distribution of units Class II molar relationship by zones	139
4.20	Distribution of maxillary overjet of the sample by zones	141
4.21	Distribution of maxillary overjet of the sample by location (urban & rural) and gender	141
4.22	Distribution of reverse overjet by zones	142
4.23	Distribution of reverse overjet by location (urban & rural) and gender	142
4.24	Distribution of anterior crossbite of the sample according to zones	143
4.25	Distribution of anterior crossbite according to location (urban & rural) and gender	143
4.26	Association of anterior crossbite between demographic variables	144
4.27	Association of overbite of the sample between zones	146
4.28	Association of overbite of the sample between location (urban & rural) and gender	146
4.29	Distribution of anterior openbite by zones	147
4.30	Distribution of anterior openbite by location (urban & rural) and gender	147
4.31	Distribution of posterior openbite by zones	148
4.32	Distribution of posterior openbite by location (urban & rural) and gender	148
4.33	Distribution of posterior crossbite by zones	149
4.34	Distribution of posterior crossbite among location (urban & rural) and gender	149
4.35	Association of posterior openbite between demographic variables	150
4.36	Distribution of scissor bite by zones	151
4.37	Distribution of scissor bite by location (urban & rural) and gender	151
4.38	Distribution of partially erupted teeth for the jaw	152
4.39	Distribution of partially erupted teeth between demographic variables	153
4.40	Crosstabulation of right and left canine relationship classes	154

4.41	Distribution of canine relationship discrepancy in right and left sides of occlusion by zones	155
4.42	Distribution of canine relationship discrepancy in right and left sides of occlusion by location and gender	156
4.43	Distribution of bimaxillary protrusion by zones	157
4.44	Distribution of bimaxillary protrusion by location and gender	157
4.45	Distribution of normative orthodontic treatment need as measured by the Dental Health Component (DHC) of IOTN	158
4.46	Distribution of normative orthodontic treatment need by zones	159
4.47	Distribution of normative orthodontic treatment need by location (urban & rural) and gender	160
4.48	Distribution of Aesthetic Component grades as evaluated by examiner and subjects	161
4.49	Distributions of treatment need categories as evaluated by examiner using Aesthetic Component by zones	162
4.50	Distributions of treatment need categories evaluated by examiner using Aesthetic Component by location (urban & rural) and gender	163
4.51	Subject evaluation of treatment need by zones as measured by Aesthetic Component of IOTN	164
4.52	Subject evaluation of treatment need by location and gender as measured by Aesthetic Component of IOTN	164
4.53	Relationship of normative treatment and subjects perceived needs	165
4.54	Malocclusion prevalence as measured using FDI/WHO method	167
4.55	Prevalence of malocclusion prevalence as measured based on IOTN	169
4.56	Distribution of different severity of malocclusion responsible in determining grades of treatment need of IOTN	170
4.57	Occlusion status comparison as measured by FDI/WHO method of occlusal traits and IOTN	171

LIST OF ABBREVIATIONS USED IN THE THESIS

Abbreviation	Meaning
ABO	American Board of Orthodontics
AC	Aesthetic Component
AD	Anno Domini
BC	Before Christ
CalMod	California modification
COCSTOC	Commission on Classification and Statistics for Oral Conditions
DAI	Dental Aesthetic Index
DHC	Dental Health Component
FDI	Fédération Dentaire Internationale
HLDI	Handicapping Labio-lingual Deviations Index
HMAR	Handicapping Malocclusion Assessment Record
ICON	Index of Complexity Outcome and Need
IOTN	Index of Orthodontic Treatment Need
MERM	Method for Epidemiological Registration of Malocclusion
MOCDO	Missing, Overjet, Crossbites, Displacement, Overbite
MSE	Malocclusion Severity Estimate
NOTI	Need for Orthodontic Treatment Index
°C	Degree Celsius
OFI	Occlusal Feature Index
PAR	Peer Assessment Rating
Per annum	Per year
SCAN	Index of Standardised Continuum of Aesthetic Need
TDI	Treatment Difficulty Index
TPI	Treatment Priority Index
WHO	World Health Organisation

List of Appendices

Appendix Number	Description	Page
1	Details of Dental Health Component grades of IOTN	233
2	Some interpretation for Index of Orthodontic Treatment Need	235
3	Dental Health Component (DHC) ruler description	236
4	Aesthetic Component grades of IOTN	237
5	Ethics approval	238
6	Letter from supervisor and certified of Yemeni Embassy in Kuala Lumpur to Thamar University in Yemen	239
7	A letter from Vice-Rector, Postgraduate Studies & Scientific Research in Thamar University to Yemeni Director of Education [in English]	240
8	A letter from Vice-Rector, Postgraduate Studies & Scientific Research in Thamar University to Yemeni Director of Education [in Arabic]	241
9	Protocol and permission latter to participating schools' headmasters [in English]	242
10	Protocol and permission latter to participating schools' headmasters [in Arabic]	243
11	Participant consent letter for photograph [in English]	244
12	Participant consent letter for photograph [in Arabic]	245
13	Clinical examination form	246
14	Distribution of dental discrepancies of pilot study	250
15	Distribution of space discrepancies of pilot study	250
16	Distribution of antero-posterior occlusal relationship of pilot study	251
17	Distribution of vertical occlusal relationship of pilot study	251
18	Distribution of molar and canine relationships of pilot study	252
19	Treatment need according to Dental Health Component of IOTN - results of pilot study	252

20	Treatment need of pilot study according to Aesthetic Component of IOTN	253
21	Distribution of occlusal traits of IOTN grades among the sample	254
22	Distribution of occlusal traits of IOTN grades by zones	255
23	Distribution of occlusal traits of IOTN grades by location and gender	256
24	Correlation of normative and subjects perceived orthodontic treatment need categories	257
25	Correlation of examiner and subjects according to Aesthetic Component treatment need categories	257
26	Letter from Yemen Dental Association regarding to the number of orthodontists in Yemen	258
27	Crosstabulation of orthodontic treatment need categories with Incisor Classification Classes	259