

REFERENCES

- Abdullah ,N, A, B (2008) : Development of recombinant *Mycobacterium bovis* Bacille calmette guerin(rBCG) expressing the 19kDa-c terminus of merozoite surface protein-1(MSP-1C) and the 22kDa of serine repeat antigen (SE22) of *plasmodium falciparum* as a potential blood-stage malaria vaccine , PHD thesis .
- Aikawa, M., Iseki, M., Barnwell, J. W., Taylor, D., Oo, M. M., & Howard, R. J. (1990). The pathology of human cerebral malaria. *Am J Trop Med Hyg*, 43(2 Pt 2), 30-37.
- Al Yaman, F. M., Mokela, D., Genton, B., Rockett, K. A., Alpers, M. P., & Clark, I. A. (1996). Association between serum levels of reactive nitrogen intermediates and coma in children with cerebral malaria in Papua New Guinea. *Trans R Soc Trop Med Hyg*, 90(3), 270-273.
- Al-Yaman, F., Genton, B., Reeder, J. C., Mokela, D., Anders, R. F., & Alpers, M. P. (1997). Humoral response to defined *Plasmodium falciparum* antigens in cerebral and uncomplicated malaria and their relationship to parasite genotype. *Am J Trop Med Hyg*, 56(4), 430-435.
- Anderson, T. J., Haubold, B., Williams, J. T., Estrada-Franco, J. G., Richardson, L., Mollinedo, R., et al. (2000). Microsatellite markers reveal a spectrum of population structures in the malaria parasite *Plasmodium falciparum*. *Mol Biol Evol*, 17(10), 1467-1482.
- Arez, A. P., Snounou, G., Pinto, J., Sousa, C. A., Modiano, D., Ribeiro, H., et al. (1999). A clonal *Plasmodium falciparum* population in an isolated outbreak of malaria in the Republic of Cabo Verde. *Parasitology*, 118 (Pt 4), 347-355.

- Ariey, F., Chalvet, W., Hommel, D., Peneau, C., Hulin, A., Mercereau-Puijalon, O., et al. (1999). *Plasmodium falciparum* parasites in French Guiana: limited genetic diversity and high selfing rate. *Am J Trop Med Hyg*, 61(6), 978-985.
- Ariey, F., Hommel, D., Le Scanf, C., Duchemin, J. B., Peneau, C., Hulin, A., et al. (2001). Association of severe malaria with a specific *Plasmodium falciparum* genotype in French Guiana. *J Infect Dis*, 184(2), 237-241.
- Arnot, D. (1998). Unstable malaria in Sudan: the influence of the dry season. Clone multiplicity of *Plasmodium falciparum* infections in individuals exposed to variable levels of disease transmission. *Trans R Soc Trop Med Hyg*, 92(6), 580-585.
- Aubouy, A., Migot-Nabias, F., & Deloron, P. (2003). Polymorphism in two merozoite surface proteins of *Plasmodium falciparum* isolates from Gabon. *Malar J*, 2, 12.
- Babiker, H. A., Charlwood, J. D., Smith, T., & Walliker, D. (1995). Gene flow and cross-mating in *Plasmodium falciparum* in households in a Tanzanian village. *Parasitology*, 433-442.
- Babiker, H. A., Lines, J., Hill, W. G., & Walliker, D. (1997). Population structure of *Plasmodium falciparum* in villages with different malaria endemicity in east Africa. *Am J Trop Med Hyg*, 56(2), 141-147.
- Babiker, H. A., Ranford-Cartwright, L. C., & Walliker, D. (1999). Genetic structure and dynamics of *Plasmodium falciparum* infections in the Kilombero region of Tanzania. *Trans R Soc Trop Med Hyg*, 93 Suppl 1, 11-14.

- Babiker, H. A., Satti, G., & Walliker, D. (1995). Genetic changes in the population of *Plasmodium falciparum* in a Sudanese village over a three-year period. *Am J Trop Med Hyg*, 53(1), 7-15.
- Basco, L. K., Tahar, R., & Escalante, A. (2004). Molecular epidemiology of malaria in Cameroon. XVIII. Polymorphisms of the *Plasmodium falciparum* merozoite surface antigen-2 gene in isolates from symptomatic patients. *Am J Trop Med Hyg*, 70(3), 238-244.
- Beck, H. P., Felger, I., Huber, W., Steiger, S., Smith, T., Weiss, N., et al. (1997). Analysis of multiple *Plasmodium falciparum* infections in Tanzanian children during the phase III trial of the malaria vaccine SPf66. *J Infect Dis*, 175(4), 921-926.
- Bendixen, M., Msangeni, H. A., Pedersen, B. V., Shayo, D., & Bodker, R. (2001). Diversity of *Plasmodium falciparum* populations and complexity of infections in relation to transmission intensity and host age: a study from the Usambara Mountains, Tanzania. *Trans R Soc Trop Med Hyg*, 95(2), 143-148.
- Black, C. G., Wu, T., Wang, L., Hibbs, A. R., & Coppel, R. L. (2001). Merozoite surface protein 8 of *Plasmodium falciparum* contains two epidermal growth factor-like domains. *Mol Biochem Parasitol*, 114(2), 217-226.
- Blackman, M. J., & Holder, A. A. (1992). Secondary processing of the *Plasmodium falciparum* merozoite surface protein-1 (*MSP-1*) by a calcium-dependent membrane-bound serine protease: shedding of *MSP-133* as a noncovalently associated complex with other fragments of the *MSP-1*. *Mol Biochem Parasitol*, 50(2), 307-315.

- Bogreau, H., Renaud, F., Bouchiba, H., Durand, P., Assi, S. B., Henry, M. C., et al. (2006). Genetic diversity and structure of African *Plasmodium falciparum* populations in urban and rural areas. *Am J Trop Med Hyg*, 74(6), 953-959.
- Bruce, M. C., Galinski, M. R., Barnwell, J. W., Donnelly, C. A., Walmsley, M., Alpers, M. P., et al. (2000). Genetic diversity and dynamics of *plasmodium falciparum* and *P. vivax* populations in multiply infected children with asymptomatic malaria infections in Papua New Guinea. *Parasitology*, 121 (Pt 3), 257-272.
- Cattamanchi, A., Kyabayinze, D., Hubbard, A., Rosenthal, P. J., & Dorsey, G. (2003). Distinguishing recrudescence from reinfection in a longitudinal antimalarial drug efficacy study: comparison of results based on genotyping of *MSP-1*, *MSP-2*, and *glurp*. *Am J Trop Med Hyg*, 68(2), 133-139.
- Chappel, J. A., & Holder, A. A. (1993). Monoclonal antibodies that inhibit *Plasmodium falciparum* invasion in vitro recognise the first growth factor-like domain of merozoite surface protein-1. *Mol Biochem Parasitol*, 60(2), 303-311.
- Chenet, S. M., Branch, O. H., Escalante, A. A., Lucas, C. M., & Bacon, D. J. (2008). Genetic diversity of vaccine candidate antigens in *Plasmodium falciparum* isolates from the Amazon basin of Peru. *Malar J*, 7, 93.
- Clark, J. T., Donachie, S., Anand, R., Wilson, C. F., Heidrich, H. G., & McBride, J. S. (1989). 46-53 kilodalton glycoprotein from the surface of *Plasmodium falciparum* merozoites. *Mol Biochem Parasitol*, 32(1), 15-24.
- Conway, D. J., Greenwood, B. M., & McBride, J. S. (1991). The epidemiology of multiple-clone *Plasmodium falciparum* infections in Gambian patients. *Parasitology*, 103 Pt 1, 1-6.

- Coppel, R. L., Cowman, A. F., Lingelbach, K. R., Brown, G. V., Saint, R. B., Kemp, D. J., et al. (1983). Isolate-specific S-antigen of *Plasmodium falciparum* contains a repeated sequence of eleven amino acids. *Nature*, 306(5945), 751-756.
- Daubersies, P., Sallenave-Sales, S., Magne, S., Trape, J. F., Contamin, H., Fandeur, T., et al. (1996). Rapid turnover of *Plasmodium falciparum* populations in asymptomatic individuals living in a high transmission area. *Am J Trop Med Hyg*, 54(1), 18-26.
- Davis, T. M., Supanaranond, W., Pukrittayakamee, S., Silamut, K., & White, N. J. (2003). Evidence for continued two-brood replication of *Plasmodium falciparum* in vivo during quinine treatment. *Acta Trop*, 89(1), 41-45.
- David, P. H., Hadley, T. J., Aikawa, M., & Miller, L. H. (1984). Processing of a major parasite surface glycoprotein during the ultimate stages of differentiation in *Plasmodium knowlesi*. *Mol Biochem Parasitol*, 11, 267-282.
- Decuypere, S., Elinck, E., Van Overmeir, C., Talisuna, A. O., D'Alessandro, U., & Dujardin, J. C. (2003). Pathogen genotyping in polyclonal infections: application of a fluorogenic polymerase-chain-reaction assay in malaria. *J Infect Dis*, 188(8), 1245-1249.
- Deleersnijder, W., Hendrix, D., Bendahman, N., Hanegreefs, J., Brijs, L., Hamers-Casterman, C., et al. (1990). Molecular cloning and sequence analysis of the gene encoding the major merozoite surface antigen of *Plasmodium chabaudi* IP-PC1. *Mol Biochem Parasitol*, 43(2), 231-244.
- Dent, A. E., Yohn, C. T., Zimmerman, P. A., Vulule, J., Kazura, J. W., & Moormann, A. M. (2007). A polymerase chain reaction/ligase detection reaction fluorescent

- microsphere assay to determine *Plasmodium falciparum* MSP-119 haplotypes. *Am J Trop Med Hyg*, 77(2), 250-255.
- Eliades, M. J., Wolkon, A., Morgah, K., Crawford, S. B., Dorkenoo, A., Sodahlon, Y., et al. (2006). Burden of malaria at community level in children less than 5 years of age in Togo. *Am J Trop Med Hyg*, 75(4), 622-629.
- Engelbrecht, F., Togel, E., Beck, H. P., Enwezor, F., Oettli, A., & Felger, I. (2000). Analysis of *Plasmodium falciparum* infections in a village community in Northern Nigeria: determination of MSP-2 genotypes and parasite-specific IgG responses. *Acta Trop*, 74(1), 63-71.
- Epstein, N., Miller, L. H., Kaushel, D. C., Udeinya, I. J., Rener, J., Howard, R. J., et al. (1981). Monoclonal antibodies against a specific surface determinant on malarial (*Plasmodium knowlesi*) merozoites block erythrocyte invasion. *J Immunol*, 127(1), 212-217.
- Farnert, A., Arez, A. P., Babiker, H. A., Beck, H. P., Benito, A., Bjorkman, A., et al. (2001). Genotyping of *Plasmodium falciparum* infections by PCR: a comparative multicentre study. *Trans R Soc Trop Med Hyg*, 95(2), 225-232.
- Farnert, A., Rooth, I., Svensson, Snounou, G., & Bjorkman, A. (1999). Complexity of *Plasmodium falciparum* infections is consistent over time and protects against clinical disease in Tanzanian children. *J Infect Dis*, 179(4), 989-995.
- Farnert, A., Snounou, G., Rooth, I., & Bjorkman, A. (1997). Daily dynamics of *Plasmodium falciparum* subpopulations in asymptomatic children in a holoendemic area. *Am J Trop Med Hyg*, 56(5), 538-547.

- Felger, I., Marshal, V. M., Reeder, J. C., Hunt, J. A., Mgone, C. S., & Beck, H. P. (1997). Sequence diversity and molecular evolution of the merozoite surface antigen 2 of *Plasmodium falciparum*. *J Mol Evol*, 45(2), 154-160.
- Felger, I., Smith, T., Edoh, D., Kitua, A., Alonso, P., Tanner, M., et al. (1999). Multiple *Plasmodium falciparum* infections in Tanzanian infants. *Trans R Soc Trop Med Hyg*, 93 Suppl 1, 29-34.
- Felger, I., Tavul, L., Kabintik, S., Marshall, V., Genton, B., Alpers, M., et al. (1994). *Plasmodium falciparum*: extensive polymorphism in merozoite surface antigen 2 alleles in an area with endemic malaria in Papua New Guinea. *Exp Parasitol*, 79(2), 106-116.
- Fenton, B., Clark, J. T., Wilson, C. F., McBride, J. S., & Walliker, D. (1989). Polymorphism of a 35-48 kDa *Plasmodium falciparum* merozoite surface antigen. *Mol Biochem Parasitol*, 34(1), 79-86.
- Garcia, J. E., Puentes, A., & Patarroyo, M. E. (2006). Developmental biology of sporozoite-host interactions in *Plasmodium falciparum* malaria: implications for vaccine design. *Clin Microbiol Rev*, 19(4), 686-707.
- Gardner, M. J., Hall, N., Fung, E., White, O., Berriman, M., Hyman, R. W., et al. (2002). Genome sequence of the human malaria parasite *Plasmodium falciparum*. *Nature*, 419(6906), 498-511.
- Genton, B., Betuela, I., Felger, I., Al-Yaman, F., Anders, R. F., Saul, A., et al. (2002). A recombinant blood-stage malaria vaccine reduces *Plasmodium falciparum* density and exerts selective pressure on parasite populations in a phase 1-2b trial in Papua New Guinea. *J Infect Dis*, 185(6), 820-827.

- Gerold, P., Schofield, L., Blackman, M. J., Holder, A. A., & Schwarz, R. T. (1996). Structural analysis of the glycosyl-phosphatidylinositol membrane anchor of the merozoite surface proteins-1 and -2 of *Plasmodium falciparum*. *Mol Biochem Parasitol*, 75(2), 131-143.
- Ghanchi, N. K., Martensson, A., Ursing, J., Jafri, S., Bereczky, S., Hussain, R. (2010). Genetic diversity among *Plasmodium falciparum* field isolates in Pakistan measured with PCR genotyping of the merozoite surface protein 1 and 2. *Malar J*, 9, 1.
- Gibson, H. L., Tucker, J. E., Kaslow, D. C., Krettli, A. U., Collins, W. E., Kiefer, M. C., et al. (1992). Structure and expression of the gene for Pv200, a major blood-stage surface antigen of *Plasmodium vivax*. *Mol Biochem Parasitol*, 50(2), 325-333.
- Gilles hm, Warrell, da (1993) Bruce-Chwatt's essential malariology. 3rd ed. Edward Arnold,Hodder and Stoughton, London.
- Gomez, D., Chaparro, J., Rubiano, C., Rojas, M. O., & Wasserman, M. (2002). Genetic diversity of *Plasmodium falciparum* field samples from an isolated Colombian village. *Am J Trop Med Hyg*, 67(6), 611-616.
- Guinovart, C., Navia, M. M., Tanner, M., & Alonso, P. L. (2006). Malaria: burden of disease. *Curr Mol Med*, 6(2), 137-140.
- Haddad, D., Snounou, G., Mattei, D., Enamorado, I. G., Figueroa, J., Stahl, S., et al. (1999). Limited genetic diversity of *Plasmodium falciparum* in field isolates from Honduras. *Am J Trop Med Hyg*, 60(1), 30-34.
- Happi, C. T., Gbotosho, G. O., Sowunmi, A., Falade, C. O., Akinboye, D. O., Gerena, L., et al. (2004). Molecular analysis of *Plasmodium falciparum* recrudescence malaria

- infections in children treated with chloroquine in Nigeria. *Am J Trop Med Hyg*, 70(1), 20-26.
- Healer, J., Murphy, V., Hodder, A. N., Masciantonio, R., Gemmill, A. W., Anders, R. F., et al. (2004). Allelic polymorphisms in apical membrane antigen-1 are responsible for evasion of antibody-mediated inhibition in *Plasmodium falciparum*. *Mol Microbiol*, 52(1), 159-168.
- Heidari, A., Keshavarz, H., Rokni, M. B., & Jelinek, T. (2007). Genetic diversity in merozoite surface protein (MSP)-1 and *MSP-2* genes of *Plasmodium falciparum* in a major endemic region of Iran. *Korean J Parasitol*, 45(1), 59-63.
- Herrera, S., Rudin, W., Herrera, M., Clavijo, P., Mancilla, L., de Plata, C., et al. (1993). A conserved region of the *MSP-1* surface protein of *Plasmodium falciparum* contains a recognition sequence for erythrocyte spectrin. *EMBO J*, 12(4), 1607-1614.
- Holder, A. A., & Freeman, R. R. (1984). The three major antigens on the surface of *Plasmodium falciparum* merozoites are derived from a single high molecular weight precursor. *J Exp Med*, 160(2), 624-629.
- Holder, A. A., & Blackman, M. J. (1994). What is the function of MSP-I on the malaria merozoite? *Parasitol Today*, 10(5), 182-184.
- Irion, A., Beck, H. P., & Felger, I. (1997). New repeat unit and hot spot of recombination in FC27-type alleles of the gene coding for *Plasmodium falciparum* merozoite surface protein 2. *Mol Biochem Parasitol*, 90(1), 367-370.

- Irion, A. (2000). Molecular diversity and immunological properties of the *Plasmodium falciparum* Merozoite Surface Protein 2 (MSP-2). PhD Thesis ,University of Basel, Switzerland.
- Issifou, S., Djikou, S., Sanni, A., Lekoulou, F., & Ntoumi, F. (2001). No influence of season of transmission nor age of patients on the complexity and genetic diversity of *Plasmodium falciparum* infection in Cotonou, Benin. *Bull Soc Pathol Exot*, 94(2 Pt 2), 195-198.
- Joshi, H., Valecha, N., Verma, A., Kaul, A., Mallick, P. k. Shalini, S. et al. (2007). Genetic structure of *Plasmodium falciparum* field isolates in eastern and north-eastern India. *Malar J*, 6, 60.
- Kang, J. M., Moon, S. U., Kim, J. Y., Cho, S. H., Lin, K., Sohn, W. M., (2010). Genetic polymorphism of merozoite surface protein-1 and merozoite surface protein-2 in *Plasmodium falciparum* field isolates from Myanmar. *Malar J*, 9, 131.
- Khatoon, L., Baliraine, F. N., Bonizzoni, M., Malik, S. A., & Yan, G (2010). Genetic structure of *Plasmodium vivax* and *Plasmodium falciparum* in the Bannu district of Pakistan. *Malar J*, 9, 112.
- Konate, L., Zwetyenga, J., Rogier, C., Bischoff, E., Fontenille, D. Tall, A. (1999). Variation of *Plasmodium falciparum* MSP-1 block 2 and MSP-2 allele prevalence and of infection complexity in two neighbouring Senegalese villages with different transmission conditions. *Trans R Soc Trop Med Hyg*, 93 Suppl 1, 21-28.
- Krajden, S., Panisko, D. M., Tobe, B., Yang, J., & Keystone, J. S. (1991). Prolonged infection with *Plasmodium falciparum* in a semi-immune patient. *Trans R Soc Trop Med Hyg*, 85(6), 731-732.

- Kumar, S., Epstein, J. E., & Richie, T. L. (2002). Vaccines against asexual stage malaria parasites. *Chem Immunol*, 80, 262-286.
- Kushwaha, A., Rao, P. P., Suresh, R. P., & Chauhan, V. S. (2001). Immunogenicity of recombinant fragments of *Plasmodium falciparum* acidic basic repeat antigen produced in *Escherichia coli*. *Parasite Immunol*, 23(8), 435-444.
- Kyes, S., Harding, R., Black, G., Craig, A., Peshu, N., Newbold, C., et al. (1997). Limited spatial clustering of individual *Plasmodium falciparum* alleles in field isolates from coastal Kenya. *Am J Trop Med Hyg*, 57(2), 205-215.
- Lawrence, N., Stowers, A., Mann, V., Taylor, D., & Saul, A. (2000). Recombinant chimeric proteins generated from conserved regions of *Plasmodium falciparum* merozoite surface protein 2 generate antiparasite humoral responses in mice. *Parasite Immunol*, 22(5), 211-221.
- Lengeler, C. (2004). Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Syst Rev*(2), CD000363.
- Lewis, A. P. (1989). Cloning and analysis of the gene encoding the 230-kilodalton merozoite surface antigen of *Plasmodium yoelii*. *Mol Biochem Parasitol*, 36(3), 271-282.
- Liljander, A. (2010). *Malaria : multiclonal infections and protective immunity*. Karolinska Institutet, Stockholm, Sweden.
- MacLeod, A. (2004). Minisatellites and MVR-PCR for the individual identification of parasite isolates. *Methods Mol Biol*, 270, 187-202.
- Marks, F., Meyer, C. G., Sievertsen, J., Timmann, C., Evans, J., Horstmann, R. D., et al. (2004). Genotyping of *Plasmodium falciparum* pyrimethamine resistance by

- matrix-assisted laser desorption-ionization time-of-flight mass spectrometry. *Antimicrob Agents Chemother*, 48(2), 466-472.
- Marshall, V. M., Anthony, R. L., Bangs, M. J., Purnomo, Anders, R. F., & Coppel, R. L. (1994). Allelic variants of the *Plasmodium falciparum* merozoite surface antigen 2 (MSA-2) in a geographically restricted area of Irian Jaya. *Mol Biochem Parasitol*, 63(1), 13-21.
- Marshall, V. M., Coppel, R. L., Anders, R. F., & Kemp, D. J. (1992). Two novel alleles within subfamilies of the merozoite surface antigen 2 (MSA-2) of *Plasmodium falciparum* *Mol Biochem Parasitol*, 50(1), 181-184.
- Marshall, V. M., Silva, A., Foley, M., Cranmer, S., Wang, L., McColl, D. J., et al. (1997). A second merozoite surface protein (MSP-4) of *Plasmodium falciparum* that contains an epidermal growth factor-like domain. *Infect Immun*, 65(11), 4460-4467.
- Marshall, V. M., Tieqiao, W., & Coppel, R. L. (1998). Close linkage of three merozoite surface protein genes on chromosome 2 of *Plasmodium falciparum*. *Mol Biochem Parasitol*, 94(1), 13-25.
- Mayor, A., Saute, F., Aponte, J. J., Almeda, J., Gomez-Olive, F. X., Dgedge, M., et al. (2003). *Plasmodium falciparum* multiple infections in Mozambique, its relation to other malariological indices and to prospective risk of malaria morbidity. *Trop Med Int Health*, 8(1), 3-11.
- McColl, D. J., Silva, A., Foley, M., Kun, J. F., Favaloro, J. M., Thompson, J. K., et al. (1994). Molecular variation in a novel polymorphic antigen associated with *Plasmodium falciparum* merozoites. *Mol Biochem Parasitol*, 68(1), 53-67.

- McKean, P. G., O'Dea, K., & Brown, K. N. (1993). Nucleotide sequence analysis and epitope mapping of the merozoite surface protein 1 from *Plasmodium chabaudi chabaudi* AS. *Mol Biochem Parasitol*, 62(2), 199-209.
- Menegon, M., Severini, C., Sannella, A., Paglia, M. G., Sangare, D., Abdel-Wahab, A., et al. (2000). Genotyping of *Plasmodium falciparum* gametocytes by reverse transcriptase polymerase chain reaction. *Mol Biochem Parasitol*, 111(1), 153-161.
- Miller, L. H., Roberts, T., Shahabuddin, M., & McCutchan, T. F. (1993). Analysis of sequence diversity in the *Plasmodium falciparum* merozoite surface protein-1 (MSP-1). *Mol Biochem Parasitol*, 59(1), 1-14.
- Montoya, L., Maestre, A., Carmona, J., Lopes, D., Do Rosario, V., & Blair, S. (2003). *Plasmodium falciparum*: diversity studies of isolates from two Colombian regions with different endemicity. *Exp Parasitol*, 104(1-2), 14-19.
- Mugittu, K., Adjuik, M., Snounou, G., Ntoumi, F., Taylor, W., Mshinda, H., et al. (2006). Molecular genotyping to distinguish between recrudescents and new infections in treatment trials of *Plasmodium falciparum* malaria conducted in Sub-Saharan Africa: adjustment of parasitological outcomes and assessment of genotyping effectiveness. *Trop Med Int Health*, 11(9), 1350-1359.
- Muller, O., Traore, C., Becher, H., & Kouyate, B. (2003). Malaria morbidity, treatment-seeking behaviour, and mortality in a cohort of young children in rural Burkina Faso. *Trop Med Int Health*, 8(4), 290-296.
- Nwagwu, M., Haynes, J. D., Orlandi, P. A., & Chulay, J. D. (1992). *Plasmodium falciparum*: chymotryptic-like proteolysis associated with a 101-kDa acidic-basic repeat antigen. *Exp Parasitol*, 75(4), 399-414.

- O'Dempsey, T. J., McArdle, T. F., Laurence, B. E., Lamont, A. C., Todd, J. E., & Greenwood, B. M. (1993). Overlap in the clinical features of pneumonia and malaria in African children. *Trans R Soc Trop Med Hyg*, 87(6), 662-665.
- Oeuvray, C., Bouharoun-Tayoun, H., Gras-Masse, H., Bottius, E., Kaidoh, T., Aikawa, M., et al. (1994). Merozoite surface protein-3: a malaria protein inducing antibodies that promote *Plasmodium falciparum* killing by cooperation with blood monocytes. *Blood*, 84(5), 1594-1602.
- Pachebat, J. A., Ling, I. T., Grainger, M., Trucco, C., Howell, S., Fernandez-Reyes, D., et al. (2001). The 22 kDa component of the protein complex on the surface of *Plasmodium falciparum* merozoites is derived from a larger precursor, merozoite surface protein 7. *Mol Biochem Parasitol*, 117(1), 83-89.
- Paul, R. E., & Day, K. P. (1998). Mating patterns of *Plasmodium falciparum*. *Parasitol Today*, 14(5), 197-202.
- Paul, R. E., Hackford, I., Brockman, A., Muller-Graf, C., Price, R., Luxemburger, C., et al. (1998). Transmission intensity and *Plasmodium falciparum* diversity on the northwestern border of Thailand. *Am J Trop Med Hyg*, 58(2), 195-203.
- Paul, R. E., Packer, M. J., Walmsley, M., Lagog, M., Ranford-Cartwright, L. C., Paru, R., et al. (1995). Mating patterns in malaria parasite populations of Papua New Guinea. *Science*, 269(5231), 1709-1711.
- Pearce, J. A., Mills, K., Triglia, T., Cowman, A. F., & Anders, R. F. (2005). Characterisation of two novel proteins from the asexual stage of *Plasmodium falciparum*, H101 and H103. *Mol Biochem Parasitol*, 139(2), 141-151.

- Peyerl-Hoffmann, G., Jelinek, T., Kilian, A., Kabagambe, G., Metzger, W. G., & von Sonnenburg, F. (2001). Genetic diversity of *Plasmodium falciparum* and its relationship to parasite density in an area with different malaria endemicities in West Uganda. *Trop Med Int Health*, 6(8), 607-613.
- Ranjit, M. R., Das, A., Das, B. P., Das, B. N., Dash, B. P., & Chhotray, G. P. (2005). Distribution of *Plasmodium falciparum* genotypes in clinically mild and severe malaria cases in Orissa, India. *Trans R Soc Trop Med Hyg*, 99(5), 389-395.
- Robert, F., Ntoumi, F., Angel, G., Candito, D., Rogier, C., Fandeur, T., et al. (1996). Extensive genetic diversity of *Plasmodium falciparum* isolates collected from patients with severe malaria in Dakar, Senegal. *Trans R Soc Trop Med Hyg*, 90(6), 704-711.
- Rubio, J. M., Berzosa, P. J., & Benito, A. (2001). Amplified fragment length polymorphism (AFLP) protocol for genotyping the malarial parasite *Plasmodium falciparum*. *Parasitology*, 123(Pt 4), 331-336.
- Sachs, J., & Malaney, P. (2002). The economic and social burden of malaria. *Nature*, 415(6872), 680-685.
- Sakihama, N., Ohmae, H., Bakote'e, B., Kawabata, M., Hirayama, K., & Tanabe, K. (2006). Limited allelic diversity of *Plasmodium falciparum* merozoite surface protein 1 gene from populations in the Solomon Islands. *Am J Trop Med Hyg*, 74(1), 31-40.
- Sallenave-Sales, S., Ferreira-da-Cruz, M. F., Faria, C. P., Cerruti, C., Jr., Daniel-Ribeiro, C. T., & Zalis, M. G. (2003). *Plasmodium falciparum*: limited genetic diversity of

- MSP-2* in isolates circulating in Brazilian endemic areas. *Exp Parasitol*, 103(3-4), 127-135.
- Schoepflin, S., Valsangiacomo, F., Lin, E., Kiniboro, B., Mueller, I., & Felger, I. (2009). Comparison of *Plasmodium falciparum* allelic frequency distribution in different endemic settings by high-resolution genotyping. *Malar J*, 8, 250.
- Silva, N. S., Silveira, L. A., Machado, R. L., Pova, M. M., & Ferreira, M. U. (2000). Temporal and spatial distribution of the variants of merozoite surface protein-1 (*MSP-1*) in *Plasmodium falciparum* populations in Brazil. *Ann Trop Med Parasitol*, 94(7), 675-688.
- Smith, T., Beck, H. P., Kitua, A., Mwankusye, S., Felger, I., Fraser-Hurt, N., et al. (1999). Age dependence of the multiplicity of *Plasmodium falciparum* infections and of other malariological indices in an area of high endemicity. *Trans R Soc Trop Med Hyg*, 93 Suppl 1, 15-20.
- Smythe, J. A., Coppel, R. L., Brown, G. V., Ramasamy, R., Kemp, D. J., & Anders, R. F. (1988). Identification of two integral membrane proteins of *Plasmodium falciparum*. *Proc Natl Acad Sci U S A*, 85(14), 5195-5199.
- Smythe, J. A., Peterson, M. G., Coppel, R. L., Saul, A. J., Kemp, D. J., & Anders, R. F. (1990). Structural diversity in the 45-kilodalton merozoite surface antigen of *Plasmodium falciparum*. *Mol Biochem Parasitol*, 39(2), 227-234.
- Snewin, V. A., Herrera, M., Sanchez, G., Scherf, A., Langsley, G., & Herrera, S. (1991). Polymorphism of the alleles of the merozoite surface antigens MSA1 and MSA2 in *Plasmodium falciparum* wild isolates from Colombia. *Mol Biochem Parasitol*, 49(2), 265-275.

- Snounou, G., Viriyakosol, S., Jarra, W., Thaithong, S., & Brown, K. N. (1993). Identification of the four human malaria parasite species in field samples by the polymerase chain reaction and detection of a high prevalence of mixed infections. *Mol Biochem Parasitol*, 58(2), 283-292.
- Snounou, G., Zhu, X., Siripoon, N., Jarra, W., Thaithong, S., Brown, K. N., et al. (1999). Biased distribution of *MSP-1* and *MSP-2* allelic variants in *Plasmodium falciparum* populations in Thailand. *Trans R Soc Trop Med Hyg*, 93(4), 369-374.
- Snow, R. W., Guerra, C. A., Noor, A. M., Myint, H. Y., & Hay, S. I. (2005). The global distribution of clinical episodes of *Plasmodium falciparum* malaria. *Nature*, 434(7030), 214-217.
- Soulama, I., Nebie, I., Ouedraogo, A., Gansane, A., Diarra, A., Tiono, A. B., et al. (2009). *Plasmodium falciparum* genotypes diversity in symptomatic malaria of children living in an urban and a rural setting in Burkina Faso. *Malar J*, 8, 135.
- Soulama, I., Sawadogo, M., Nebie, I., Diarra, A., Tiono, A. B., Konate, A., et al. (2006). Genetic diversity of *Plasmodium falciparum* and pathogenesis of the severe malarial anaemia in children under 5 years old in the province of Boulgou, Burkina Faso. *Bull Soc Pathol Exot*, 99(3), 166-170.
- Subbarao, S. K., & Sharma, V. P. (1997). Anopheline species complexes & malaria control. *Indian J Med Res*, 106, 164-173.
- Takala, S., Branch, O., Escalante, A. A., Kariuki, S., Wootton, J., & Lal, A. A. (2002). Evidence for intragenic recombination in *Plasmodium falciparum*: identification of a novel allele family in block 2 of merozoite surface protein-1: Asembo Bay Area Cohort Project XIV. *Mol Biochem Parasitol*, 125(1-2), 163-171

- Tibayrenc, M. (1998). Beyond strain typing and molecular epidemiology: integrated genetic epidemiology of infectious diseases. *Parasitol Today*, 14(8), 323-329.
- Toebe, C. S., Clements, J. D., Cardenas, L., Jennings, G. J., & Wiser, M. F. (1997). Evaluation of immunogenicity of an oral *Salmonella* vaccine expressing recombinant *Plasmodium berghei* merozoite surface protein-1. *Am J Trop Med Hyg*, 56(2), 192-199.
- Uthapibull, C., Aufiero, B., Syed, S. E., Hansen, B., Guevara Patino, J. A., Angov, E., et al. (2001). Inhibitory and blocking monoclonal antibody epitopes on merozoite surface protein 1 of the malaria parasite *Plasmodium falciparum*. *J Mol Biol*, 307(5), 1381-1394.
- Vafa, M., Troye-Blomberg, M., Anchang, J., Garcia, A., & Migot-Nabias, F. (2008). Multiplicity of *Plasmodium falciparum* infection in asymptomatic children in Senegal: relation to transmission, age and erythrocyte variants. *Malar J*, 7, 17.
- Vargas-Serrato, E., Barnwell, J. W., Ingravallo, P., Perler, F. B., & Galinski, M. R. (2002). Merozoite surface protein-9 of *Plasmodium vivax* and related simian malaria parasites is orthologous to p101/ABRA of *P. falciparum*. *Mol Biochem Parasitol*, 120(1), 41-52.
- Walliker, D. (1983). The genetic basis of diversity in malaria parasites. *Adv Parasitol*, 22, 217-259.
- White, N. J. (2008). *Plasmodium knowlesi*: the fifth human malaria parasite. *Clin Infect Dis*, 46(2), 172-173.
- World Health Organization (2008). "World Malaria Report 2008.
- World Health Organization (2004). "World Malaria Report 2004.

- World Health Organization (2009). "World Malaria Report 2009.
- World Health Organization (2010). "World Malaria Report 2010.
- Wongsrichanalai, C., Barcus, M. J., Muth, S., Sutamihardja, A., & Wernsdorfer, W. H. (2007). A review of malaria diagnostic tools: microscopy and rapid diagnostic test (RDT). *Am J Trop Med Hyg*, 77(6 Suppl), 119-127.
- Zakeri, S., Bereczky, S., Naimi, P., Pedro Gil, J., Djadid, N. D., Farnert, A., et al. (2005). Multiple genotypes of the merozoite surface proteins 1 and 2 in *Plasmodium falciparum* infections in a hypoendemic area in Iran. *Trop Med Int Health*, 10(10), 1060-1064.
- Zwetyenga, J., Rogier, C., Tall, A., Fontenille, D., Snounou, G., Trape, J. F., et al. (1998). No influence of age on infection complexity and allelic distribution in *Plasmodium falciparum* infections in Ndiop, a Senegalese village with seasonal, mesoendemic malaria. *Am J Trop Med Hyg*, 59(5), 726-735.