

REFERENCES

- Aguilera-Luiz, M. M., Plaza-Bolaños, P., Romero-González, R., Vidal, J. L. M., & Frenich, A. G. (2011). Comparison of the efficiency of different extraction methods for the simultaneous determination of mycotoxins and pesticides in milk samples by ultra high-performance liquid chromatography-tandem mass spectrometry. *Analytical and Bioanalytical Chemistry*, 399, 2863-2875.
- Ambrosino, P., Galvano, F., Fogliano, V., Logrieco, A., Fresa, R., & Ritieni, A. (2004). Supercritical fluid extraction of Beauvericin from maize. *Talanta*, 62(3), 523-530.
- Anastassiades, M., Lehotay, S. J., Stajnbaher, D., & Schenck, F. J. (2003). Fast and easy multiresidue method employing acetonitrile extraction/partitioning and “dispersive solid-phase extraction” for the determination of pesticide residues in produce. *Journal of AOAC INTERNATIONAL*, 86(2), 412-431.
- Ardic, M., Karakaya, Y., Atasever, M., & Durmaz, H. (2008). Determination of aflatoxin B1 levels in deep-red ground pepper (isot) using immunoaffinity column combined with ELISA. *Food and Chemical Toxicology*, 46(5), 1596-1599.
- Ardrey, R. E. (2003). *Liquid Chromatography –Mass Spectrometry: An Introduction*. West Sussex, England: John Wiley & Sons Ltd.
- Atkins, D., & Norman, J. (1998). Mycotoxins and food safety. *Nutrition and Food Science*, 5, 260 - 266.
- Aycicek, H., Aksoy, A., & Saygi, S. (2005). Determination of aflatoxin levels in some dairy and food products which consumed in Ankara, Turkey. *Food Control*, 16, 263 - 266.
- Bacaloni, A., Cavaliere, C., Cucci, F., Foglia, P., Samperi, R., & Laganà, A. (2008). Determination of aflatoxins in hazelnuts by various sample preparation methods and liquid chromatography–tandem mass spectrometry. *Journal of Chromatography A*, 1179(2), 182-189.
- Battilani, P., Barbano, C., & Logrieco, A. (2008). Chapter 1 - Risk Assessment and Safety Evaluation of Mycotoxins in Fruits. In B.-G. Rivka & P. Nachman (Eds.), *Mycotoxins in Fruits and Vegetables* (pp. 1-26). San Diego: Academic Press.
- Beltrán, E., Ibáñez, M., Sancho, J. V., Cortés, M. Á., Yusà, V., & Hernández, F. (2011). UHPLC–MS/MS highly sensitive determination of aflatoxins, the aflatoxin metabolite M1 and ochratoxin A in baby food and milk. *Food Chemistry*, 126, 737-744.
- Bennett, J. W., & Klich, M. (2003). Mycotoxins. *Clinical Microbiology Reviews*, 16(3), 497–516.

- Berthiller, F., Schuhmacher, R., Buttinger, G., & Krska, R. (2005). Rapid simultaneous determination of major type A- and B-trichothecenes as well as zearalenone in maize by high performance liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A*, 1062(2), 209-216.
- Bradburn, N., Coker, R. D., Jewers, K., & Tomlins, K. I. (1990). Evaluation of the ability of different concentrations of aqueous acetone, aqueous methanol and aqueous acetone: Methanol (1:1) to extract aflatoxin from naturally contaminated maize. *Chromatographia*, 29(9-10), 435-440.
- Cahill, L. M., Kruger, S. C., McAlice, B. T., Ramsey, C. S., Prioli, R., & Kohn, B. (1999). Quantification of deoxynivalenol in wheat using an immunoaffinity column and liquid chromatography. *Journal of Chromatography A*, 859(1), 23-28.
- Castells, M., Ramos, A. J., Sanchis, V., & Marin, S. (2007). Distribution of total aflatoxins in milled fractions of hulled rice. *Journal of Agricultural and Food Chemistry*, 55, 2760-2764.
- Cavaliere, C., D'Ascenzo, G., Foglia, P., Pastorini, E., Samperi, R., & Laganà, A. (2005). Determination of type B trichothecenes and macrocyclic lactone mycotoxins in field contaminated maize. *Food Chemistry*, 92(3), 559-568.
- Cavaliere, C., Foglia, P., Guarino, C., Nazzari, M., Samperi, R., & Laganà, A. (2007). Determination of aflatoxins in olive oil by liquid chromatography-tandem mass spectrometry. *Analytica Chimica Acta*, 596(1), 141-148.
- Christensen, H. R., Yu, F.-Y., & Chu, F. S. (2000). Development of a polyclonal antibody-based sensitive enzyme-linked immunosorbent assay for fumonisin B4. *Journal of Agricultural and Food Chemistry* 48, 1977-1984.
- Chun, H. S., Choi, E. H., Chang, H.-J., Choi, S.-W., & Eremin, S. A. (2009). A fluorescence polarization immunoassay for the detection of zearalenone in corn. *Analytica Chimica Acta*, 639(1-2), 83-89.
- Cigić, I. K. j., & Prosen, H. (2009). An Overview of Conventional and Emerging Analytical Methods for the Determination of Mycotoxins. *international Journal of Molecular Sciences*, 10, 62 -115.
- Clifford, A. A. (1998). Introduction to supercritical fluid extraction in analytical science. In E. D. Ramsey (Ed.), *Analytical supercritical fluid extraction technique*. New York, USA: Springer.
- Cole, R. J., & Cox, R. H. (1981). *Handbook of toxic fungal metabolites*. New York, N.Y.: Academic Press.

- European Commission (EC). (2006a). *Laying down the methods of sampling and analysis for the official control of the levels of mycotoxins in foodstuffs*, NO 401/2006 of 23 February 2006 .
- European Commission (EC). (2006b). *Setting maximum levels for certain contaminants in foodstuffs*, No 1881/2006 of 19 December 2006.
- Correia, S. L., Hotza, D., & Segadães, A. M. (2004). Simultaneous optimization of linear firing shrinkage and water absorption of triaxial ceramic bodies using experiments design. *Ceramics International*, 30(6), 917-922.
- Coulombe, R. A. (1993). Symposium: Biological action of mycotoxins. *Journal of Dairy Science*, 76, 880-891.
- Countryman, S., Huq, S., & Mathews, T. (2009). Rapid, High Resolution Analysis of Aflatoxin Extracts from Peanut Butter Using Kinetex™ Core-Shell Technology and Strata® SPE. *Phenomenex*.
- D'Mello, J. P. F., & Macdonald, A. M. C. (1997). Mycotoxins. *Animal Feed Science and Technology*, 69, 155 - 166.
- Dall'Asta, C., Sforza, S., Galaverna, G., Dossena, A., & Marchelli, R. (2004). Simultaneous detection of type A and type B trichothecenes in cereals by liquid chromatography-electrospray ionization mass spectrometry using NaCl as cationization agent. *Journal of Chromatography A*, 1054(1-2), 389-395.
- Dean, J., R. (1998). *Extraction methods for environmental analyis*. New York: John Wiley.
- Desmarchelier, A., Oberson, J.-M., Tella, P., Gremaud, E., Seefelder, W., & Mottier, P. (2010). Development and comparison of two multiresidue methods for the analysis of 17 mycotoxins in cereals by liquid chromatography electrospray ionization tandem mass spectrometry. *Journal of Agricultural and Food Chemistry*, 58(13), 7510-7519.
- Diener, U. L., Cole, R. J., Sanders, T. H., Payne, G. A., Lee, L. S., & Klich, M. A. (1987). Epidemiology of aflatoxin formation by *Aspergillus flavus*. *Annual Review of Phytopathology*, 25, 249 - 270.
- Elbert, D., Czapiewski, K., Bujara, I., Kunze, J., & Giger, A. (2008). Simultaneous Analysis of 10 Mycotoxins in Crude Extracts of Different Types of Grains by LC/MS/MS. *Applied Biosystems*.
- Fallah, A. A., Jafari, T., Fallah, A., & Rahnama, M. (2009). Determination of aflatoxin M1 levels in Iranian white and cream cheese. *Food and Chemical Toxicology*, 47(8), 1872-1875.

- Food and Agriculture Organization (FAO). (2004). *Worldwide Regulations for Mycotoxins in Food and Feed in 2003*. FAO Food and Nutrition Paper 81, Food and Agriculture Organization of the United Nations, Rome, Italy.
- Food and Agriculture Organization (FAO). (2001). *Safety evaluation of certain mycotoxins in food*. FAO Food and Nutrition Paper 74 / Who Food Additives Series 47, prepared by the fifty-sixth meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA), Rome, 705.
- Flajs, D., Domijan, A. M., Ivic, D., Cvjetkovic, B., & Peraica, M. (2009). ELISA and HPLC analysis of ochratoxin A in red wines of Croatia. *Food Control*, 20(6), 590-592.
- Frenich, A. G., Romero-González, R., Gómez-Pérez, M. L., & Vidal, J. L. M. (2011). Multi-mycotoxin analysis in eggs using a QuEChERS-based extraction procedure and ultra-high-pressure liquid chromatography coupled to triple quadrupole mass spectrometry. *Journal of Chromatography A*, 1218(28), 4349-4356.
- Frenich, A. G., Vidal, J. L. M., Romero-González, R., & Aguilera-Luiz, M. d. M. (2009). Simple and high-throughput method for the multimycotoxin analysis in cereals and related foods by ultra-high performance liquid chromatography/tandem mass spectrometry. *Food Chemistry*, 117(4), 705-712.
- Ghali, R., Hmaissia-khlifa, K., Ghorbel, H., Maaroufi, K., & Hedili, A. (2009). HPLC determination of ochratoxin A in high consumption Tunisian foods. *Food Control*, 20(8), 716-720.
- Hayashi, Y., & Yoshizawa, T. (2005). Analysis of cyclopiazonic acid in corn and rice by a newly developed method. *Food Chemistry*, 93(2), 215-221.
- Herzallah, S. M. (2009). Determination of aflatoxins in eggs, milk, meat and meat products using HPLC fluorescent and UV detectors. *Food Chemistry*, 114(3), 1141-1146.
- Hinojo, M. J., Medina, A., Valle-Algarra, F. M., Gimeno-Adelantado, J. V., Jiménez, M., & Mateo, R. (2006). Fumonisin production in rice cultures of *Fusarium verticillioides* under different incubation conditions using an optimized analytical method. *Food Microbiology*, 23(2), 119-127.
- Holcomb, M., Thompson Jr, H. C., Cooper, W. M., & Hopper, M. L. (1996). SFE extraction of aflatoxins (B1, B2, G1, and G2) from corn and analysis by HPLC. *Journal of Supercritical Fluids*, 9(2), 118-121.
- International Agency for Research on Cancer (IARC). (1976) *Cadmium, Nickel, Some Epoxides, Miscellaneous Industrial Chemicals and General Considerations on Volatiles Anaesthetics*. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man, Volume 11, Lyon, France.

- International Agency for Research on Cancer (IARC). (1993) *Some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins*. IARC monographs on the evaluation of carcinogenic risks to humans. Volume 56, Lyon, France.
- Ibáñez-Vea, M., Lizarraga, E., & González-Peñas, E. (2011). Simultaneous determination of type-A and type-B trichothecenes in barley samples by GC-MS. *Food Control*, 22(8), 1428-1434.
- Jaimez, J., Fente, C. A., Vazquez, B. I., Franco, C. M., Cepeda, A., G. Mahuzier, et al. (2000). Application of the assay of aflatoxins by liquid chromatography with fluorescence detection in food analysis. *Journal of Chromatography A*, 882, 1-10.
- Jelen, H. H., & Wasowicz, E. (2008). Determination of trichothecenes in wheat grain without sample cleanup using comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry. *Journal of Chromatography A*, 1215(1-2), 203-207.
- JMP®. (2010). JMP® Design of Experiments (Version 9.0.0): SAS Institute Inc.
- Josephs, R. D., Krska, R., Grasserbauer, M., & Broekaert, J. A. C. (1998). Determination of trichothecene mycotoxins in wheat by use of supercritical fluid extraction and high-performance liquid chromatography with diode array detection or gas chromatography with electron capture detection. *Journal of Chromatography A*, 795(2), 297-304.
- Juan, C., Gonzalez, L., Soriano, J. M., Molto, J. C., & Manes, J. (2005). Accelerated Solvent Extraction of Ochratoxin A from Rice Samples. *Journal of Agricultural and Food Chemistry*, 53(24), 9348-9351.
- Kabak, B. (2009). Ochratoxin A in cereal-derived products in Turkey: Occurrence and exposure assessment. *Food and Chemical Toxicology* 47, 348 - 352.
- Kaniou-Grigoriadou, I., Eleftheriadou, A., Mouratidou, T., & Katikou, P. (2005). Determination of aflatoxin M1 in ewe's milk samples and the produced curd and Feta cheese. *Food Control*, 16(3), 257-261.
- Koch, P. (2004). State of the art of trichothecenes analysis. *Toxicology Letters*, 153(1), 109-112.
- Köppen, R., Koch, M., Siegel, D., Merkel, S., Maul, R., & Nehls, I. (2010). Determination of mycotoxins in foods: Current state of analytical methods and limitations. *Applied Microbiology and Biotechnology*, 86, 1595-1612.
- Krska, R. (1998). Performance of modern sample preparation techniques in the analysis of Fusarium mycotoxins in cereals. *Journal of Chromatography A*, 815(1), 49-57.

- Langseth, W., & Rundberget, T. (1998). Instrumental methods for determination of nonmacrocyclic trichothecenes in cereals, foodstuffs and cultures. *Journal of Chromatography A*, 815(1), 103-121.
- Lattanzio, V. M. T., Pascale, M., & Visconti, A. (2009). Current analytical methods for trichothecene mycotoxins in cereals. *TrAC, Trends in Analytical Chemistry*, 28(6), 758-768.
- Leong, Y.-H., Ismail, N., Latif, A. A., & Ahmad, R. (2010). Aflatoxin occurrence in nuts and commercial nutty products in Malaysia. *Food Control*, 21(3), 334-338.
- Lin, L., Zhang, J., Wang, P., Wang, Y., & Chen, J. (1998). Thin-layer chromatography of mycotoxins and comparison with other chromatographic methods. *Journal of Chromatography A*, 815(1), 3-20.
- Liu, Z., Gao, J., & Yu, J. (2006). Aflatoxins in stored maize and rice grains in Liaoning Province, China. *Journal of Stored Products Research*, 42(4), 468-479.
- Maragos, C. (2004). Emerging technologies for mycotoxin detection. *Journal of Toxicology - Toxin Reviews*, 23(2&3), 317-344.
- Maragos, C. (2009). Fluorescence polarization immunoassay of mycotoxins: A review. *Toxins*, 1, 196-207.
- Maragos, C. M. (2006). Measurement of Aflatoxins Using Capillary Electrophoresis. In M. W. Trucksess & A. E. Pohland (Eds.), *Mycotoxin Protocols* (Vol. 157, pp. 51-58). Totowa, NJ: Humana Press.
- Maragos, C. M., & Platner, R. D. (2002). Rapid fluorescence polarization immunoassay for the mycotoxin deoxynivalenol in wheat. *Journal of Agricultural and Food Chemistry*, 50, 1827-1832.
- Micheli, L., Grecco, R., Badea, M., Moscone, D., & Palleschi, G. (2005). An electrochemical immunosensor for aflatoxin M1 determination in milk using screen-printed electrodes. *Biosensors and Bioelectronics*, 21(4), 588-596.
- Monbaliu, S., Van Poucke, C., Van Peteghem, C., Van Poucke, K., Heungens, K., & De Saeger, S. (2009). Development of a multi-mycotoxin liquid chromatography/tandem mass spectrometry method for sweet pepper analysis. *Rapid Communications in Mass Spectrometry*, 23, 3 - 11.
- Móricz, Á. M., Fatér, Z., Otta, K. H., Tyihák, E., & Mincsovics, E. (2007). Overpressured layer chromatographic determination of aflatoxin B1, B2, G1 and G2 in red paprika. *Microchemical Journal*, 85(1), 140-144.

- Murphy, P. A., Hendrich, S., Landgren, C., & Bryant, C. M. (2006). Food Mycotoxins: An Update. *Journal of Food Science*, 71(5).
- Nasir, M. S., & Jolley, M. E. (2002). Development of a fluorescence polarization assay for the determination of aflatoxins in grains. *Journal of Agricultural and Food Chemistry*, 50, 3116-3121.
- Neuhof, T., Ganzauer, N., Koch, M., & Nehls, I. (2009). A comparison of chromatographic methods for the determination of deoxynivalenol in wheat. *Chromatographia*, 69(11-12), 1457-1462.
- Nilufer, D., & Boyacioglu, D. (2002). Comparative study of three different methods for the determination of aflatoxins in Tahini. *Journal of Agricultural and Food Chemistry*, 50, 3375-3379.
- Nonaka, Y., Saito, K., Hanioka, N., Narimatsu, S., & Kataoka, H. (2009). Determination of aflatoxins in food samples by automated on-line in-tube solid-phase microextraction coupled with liquid chromatography-mass spectrometry. *Journal of Chromatography A*, 1216(20), 4416-4422.
- Osnaya, L. G., Castillo, J. M. S. d., Cortés, J. C. M., & Vinuesa, J. M. (2006). Extraction and analysis of ochratoxin A in bread using pressurised liquid extraction and liquid chromatography. *Journal of Chromatography A*, 1113(1-2), 32-36.
- Peraica, M., Radić, B., Lucić, A., & Pavlović, M. (1999). Toxic effects of mycotoxins in humans. *Bulletin of the World Health Organization*, 77(9), 754 - 766.
- Pérez-Torrado, E., Blesa, J., Moltó, J. C., & Font, G. (2010). Pressurized liquid extraction followed by liquid chromatography-mass spectrometry for determination of zearalenone in cereal flours. *Food Control*, 21(4), 399-402.
- Pesavento, M., Domagala, S., Baldini, E., & Cucca, L. (1997). Characterization of an enzyme linked immunosorbent assay for Aflatoxin B1 based on commercial reagents. *Talanta*, 45(1), 91-104.
- Peter M, S. (1993). Recent developments in methods of analysis for mycotoxins in foodstuffs. *TrAC Trends in Analytical Chemistry*, 12(9), 382-386.
- Quan, Y., Zhang, Y., Wang, S., Lee, N., & Kennedy, I. R. (2006). A rapid and sensitive chemiluminescence enzyme-linked immunosorbent assay for the determination of fumonisin B1 in food samples. *Analytica Chimica Acta*, 580(1), 1-8.
- Radova, Z., Holadova, K., & Hajslova, J. (1998). Comparison of two clean-up principles for determination of trichothecenes in grain extract. *Journal of Chromatography A*, 829, 259-267.
- Rahmani, A., Jinap, S., & Soleimany, F. (2009). Qualitative and Quantitative Analysis of Mycotoxins. *Comprehensive Reviews in Food Science and Food Safety*, 8(3), 202-251.

- Rasmussen, R. R., Storm, I. M. L. D., Rasmussen, P. H., Smedsgaard, J., & Nielsen, K. F. (2010). Multi-mycotoxin analysis of maize silage by LC-MS/MS. *Applied Microbiology and Biotechnology*, 397, 765 -776.
- Reddy, K. R. N., Reddy, C. S., & Muralidharan, K. (2009). Detection of Aspergillus spp. and aflatoxin B1 in rice in India. *Food Microbiology*, 26(1), 27-31.
- Reinsch, M., Töpfer, A., Lehmann, A., Nehls, I., & Panne, U. (2007). Determination of ochratoxin A in beer by LC-MS/MS ion trap detection. *Food Chemistry*, 100(1), 312-317.
- Richard, J. L. (2007). Influence of harvest and storage conditions on trichothecenes levels in various cereals. *International Journal of Food Microbiology*, 119, 3 -10.
- Richard, J. L., Payne, G. A., Desjardins, E. A., Maragos, C. M., Norred, W. P., Pestka, J. J., et al. (2003). *Mycotoxins: risks in plant, animal, and human systems*. Ames, Iowa (USA): Council for Agricultural Science and Technology
- Richter, B. E., Jones, B. A., Ezzell, J. L., & Porter, N. L. (1996). Accelerated solvent extraction: A technique for sample preparation. *Analytical Chemistry*, 68, 1033-1039.
- Roch, O. G., Blunden, G., Coker, R. D., & Nawaz, S. (1995). The validation of a solid phase clean-up procedure for the analysis of aflatoxins in groundnut cake using HPLC. *Food Chemistry*, 52, 93-98.
- Romero-González, R., Garrido Frenich, A., Martínez Vidal, J. L., Prestes, O. D., & Grió, S. L. (2011). Simultaneous determination of pesticides, biopesticides and mycotoxins in organic products applying a quick, easy, cheap, effective, rugged and safe extraction procedure and ultra-high performance liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A*, 1218(11), 1477-1485.
- Schollenberger, M., Lauber, U., Jara, H. T., Suchy, S., Drochner, W., & Müller, H. M. (1998). Determination of eight trichothecenes by gas chromatography-mass spectrometry after sample clean-up by a two-stage solid-phase extraction. *Journal of Chromatography A*, 815(1), 123-132.
- Sheibani, A., & Ghaziaskar, H. S. (2009). Pressurized fluid extraction for quantitative recovery of aflatoxins B1 and B2 from pistachio. *Food Control*, 20(2), 124-128.
- Shephard, G. S. (1998). Chromatographic determination of the fumonisin mycotoxins. *Journal of Chromatography A*, 815(1), 31-39.

- Silva, L., Fernández-Franzón, M., Font, G., Pena, A., Silveira, I., Lino, C., et al. (2009). Analysis of fumonisins in corn-based food by liquid chromatography with fluorescence and mass spectrometry detectors. *Food Chemistry*, 112(4), 1031-1037.
- Sospedra, I., Blesa, J., Soriano, J. M., & Mañes, J. (2010). Use of the modified quick easy cheap effective rugged and safe sample preparation approach for the simultaneous analysis of type A- and B-trichothecenes in wheat flour. *Journal of Chromatography A*, 1217(9), 1437-1440.
- Stanker, L. H., & Beier, R. C. (1996). Introduction to immunoassays for residue analysis: Concepts, formats, and Applications. In R. C. Beier, et al (Ed.), *Immunoassays for residue analysis*. Washinton DC: ACS.
- Stecher, G., Jarukamjorn, K., Zaborski, P., Bakry, R., Huck, C. W., & Bonn, G. K. (2007). Evaluation of extraction methods for the simultaneous analysis of simple and macrocyclic trichothecenes. *Talanta*, 73(2), 251-257.
- Stroka, J., Otterdijk, R. v., & Anklam, E. (2000). Immunoaffinity column clean-up prior to thin-layer chromatography for the determination of aflatoxins in various food matrices. *Journal of Chromatography A*, 904, 251–256
- Sulyok, M., Berthiller, F., Krska, R., & Schuhmacher, R. (2006). Development and validation of a liquid chromatography/tandem mass spectrometric method for the determination of 39 mycotoxins in wheat and maize. *Rapid Communications in Mass Spectrometry*, 20(18), 2649-2659.
- Sulyok, M., Krska, R., & Schuhmacher, R. (2010). Application of an LC-MS/MS based multi-mycotoxin method for the semi-quantitative determination of mycotoxins occurring in different types of food infected by moulds. *Food Chemistry*, 119(1), 408-416.
- Tan, Y., Chu, X., Shen, G.-L., & Yu, R.-Q. (2009). A signal-amplified electrochemical immunosensor for aflatoxin B1 determination in rice. *Analytical Biochemistry*, 387(1), 82-86.
- Tanaka, T., Yoneda, A., Inoue, S., Sugiura, Y., & Ueno, Y. (2000). Simultaneous determination of trichothecene mycotoxins and zearalenone in cereals by gas chromatography-mass spectrometry. *Journal of Chromatography A*, 882(1-2), 23-28.
- Tavčar-Kalcher, G., Vrtač, K., Pestevšek, U., & Vengušt, A. (2007). Validation of the procedure for the determination of aflatoxin B1 in animal liver using immunoaffinity columns and liquid chromatography with postcolumn derivatisation and fluorescence detection. *Food Control*, 18(4), 333-337.
- Thrane, U. (1989). Fusarium. In J. Chelowski (Ed.), *Mycotoxins, Taxonomy and Pathogenicity*. Amsterdam: Elsevier.

- Turner, N. W., Subrahmanyam, S., & Piletsky, S. A. (2009). Analytical methods for determination of mycotoxins: A review. *Analytica Chimica Acta*, 632(2), 168-180.
- Urraca, J. L., Marazuela, M. D., & Moreno-Bondi, M. C. (2004). Analysis for zearalenone and [alpha]-zearalenol in cereals and swine feed using accelerated solvent extraction and liquid chromatography with fluorescence detection. *Analytica Chimica Acta*, 524(1-2), 175-183.
- Valle-Algarra, F. M., Medina, A., Gimeno-Adelantado, J. V., Llorens, A., Jiménez, M., & Mateo, R. (2005). Comparative assessment of solid-phase extraction clean-up procedures, GC columns and perfluoroacetylation reagents for determination of type B trichothecenes in wheat by GC-ECD. *Talanta*, 66(1), 194-201.
- Var, I., Kabak, B., & Gök, F. (2007). Survey of aflatoxin B1 in helva, a traditional Turkish food, by TLC. *Food Control*, 18(1), 59-62.
- Ventura, M., Gomez, A., Anaya, I., Diaz, J., Broto, F., Agut, M., et al. (2004). Determination of aflatoxins B1, G1, B2 and G2 in medicinal herbs by liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A*, 1048, 25-29.
- Visconti, A., Lattanzio, V. M. T., Pascale, M., & Haidukowski, M. (2005). Analysis of T-2 and HT-2 toxins in cereal grains by immunoaffinity clean-up and liquid chromatography with fluorescence detection. *Journal of Chromatography A*, 1075(1-2), 151-158.
- Visconti, A., & Pascale, M. (1998). Determination of zearalenone in corn by means of immunoaffinity clean-up and high-performance liquid chromatography with fluorescence detection. *Journal of Chromatography A*, 815(1), 133-140.
- Whitlow, L. W., & Hagler Jr, W. M. (2002). Mycotoxins in feeds. *Feedstuffs*, 74, 10.
- World Health Organization (WHO). (2002). Safety evaluation of certain mycotoxins in food. *Fifty-sixth report of the Joint FAO/WHO Expert Committee on Food Additives, WHO Technical Report Series 906, World Health Organization, Geneva, Switzerland*, 1-74.
- Yiannikouris, A., & Jouany, J.-P. (2002). Mycotoxins in feeds and their fate in animals: a review. *Animal Resources*, 51 81 - 99.
- Yu, F.-Y., Chi, T.-F., Liu, B.-H., & Su, C.-C. (2005). Development of a sensitive enzyme-linked immunosorbent assay for the determination of ochratoxin A. *Journal of Agricultural and Food Chemistry*, 53, 6947-6953.

- Zachariasova, M., Lacina, O., Malachova, A., Kostelanska, M., Poustka, J., Godula, M., et al. (2010). Novel approaches in analysis of Fusarium mycotoxins in cereals employing ultra performance liquid chromatography coupled with high resolution mass spectrometry. *Analytica Chimica Acta*, 662(1), 51-61.
- Zhang, H., Wang, S., & Fang, G. (2011). Applications and recent developments of multi-analyte simultaneous analysis by enzyme-linked immunosorbent assays. *Journal of Immunological Methods*, 368(1-2), 1-23.
- Zheng, M. Z., Richard, J. L., & Binder, J. (2006). A review of rapid methods for the analysis of mycotoxins. *Mycopathologis*, 161, 261-273.
- Zinedine, A., Brera, C., Elakhdari, S., Catano, C., Debegnach, F., Angelini, S., et al. (2006). Natural occurrence of mycotoxins in cereals and spices commercialized in Morocco. *Food Control*, 17(11), 868-874.
- Zöllner, P., & Mayer-Helm, B. (2006). Trace mycotoxin analysis in complex biological and food matrices by liquid chromatography-atmospheric pressure ionisation mass spectrometry. *Journal of Chromatography A*, 1136(2), 123-169.
- Zougagh, M., & Ríos, Á. (2008). Supercritical fluid extraction of macrocyclic lactone mycotoxins in maize flour samples for rapid amperometric screening and alternative liquid chromatographic method for confirmation. *Journal of Chromatography A*, 1177(1), 50-57.