

CHAPTER TEN

REFERENCES

- Aciego Pietri, J. C., & Brookes, P. C. (2008). Relationships between soil pH and microbial properties in a UK arable soil. [doi: DOI: 10.1016/j.soilbio.2008.03.020]. *Soil Biology & Biochemistry*, 40(7), 1856-1861.
- Aislabie, J., Fraser, R., Duncan, S., & Farrell, R. L. (2001). Effects of oil spills on microbial heterotrophs in Antarctic soils. *Polar Biology*, 24(5), 308-313.
- Aislabie, J., Jordan, S., Ayton, J., Klassen, J. L., Barker, G. M., & Turner, S. (2009). Bacterial diversity associated with ornithogenic soil of the Ross Sea region, Antarctica. *Canadian Journal of Microbiology*, 55(1), 21-36.
- Aislabie, J. M., Chhour, K. L., Saul, D. J., Miyauchi, S., Ayton, J., Paetzold, R. F., et al. (2006). Dominant bacteria in soils of Marble Point and Wright Valley, Victoria Land, Antarctica. *Soil Biology and Biochemistry*, 38(10), 3041-3056.
- Aislabie, J. M., Jordan, S., & Barker, G. M. (2008). Relation between soil classification and bacterial diversity in soils of the Ross Sea region, Antarctica. *Geoderma*, 144(1-2), 9-20.
- An, D.-S., Lee, H.-G., Im, W.-T., Liu, Q.-M., & Lee, S.-T. (2007). *Segetibacter koreensis* gen. nov., sp. nov., a novel member of the phylum Bacteroidetes, isolated from the soil of a ginseng field in South Korea. *International Journal of Systematic and Evolutionary Microbiology*, 57(8), 1828-1833.
- Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, 26(1), 32-46.
- Anderson, M. J., Gorley, R. N., & Clarke, K. R. (2008). *PERMANOVA+ for PRIMER: Guide to software and statistical methods*. Plymouth, UK: PRIMER-E.
- André M.-F., & Hall, K. (2005). Honeycomb development on Alexander Island, glacial history of George VI Sound and palaeoclimatic implications (Two Step Cliffs/Mars Oasis, W Antarctica). *Geomorphology*, 65(1-2), 117-138.
- Arnold, R. J., Convey, P., Hughes, K. A., & Wynn-Williams, D. D. (2003). Seasonal periodicity of physical factors, inorganic nutrients and microalgae in Antarctic fellfields. *Polar Biology*, 26(6), 396-403.
- Ashelford, K. E., Chuzhanova, N. A., Fry, J. C., Jones, A. J., & Weightman, A. J. (2005). At least 1 in 20 16S rRNA sequence records currently held in public repositories is estimated to contain substantial anomalies. *Applied and Environmental Microbiology*, 71(12), 7724-7736.
- Ashelford, K. E., Chuzhanova, N. A., Fry, J. C., Jones, A. J., & Weightman, A. J. (2006). New screening software shows that most recent large 16S rRNA gene clone

- libraries contain chimeras. *Applied and Environmental Microbiology*, 72(9), 5734-5741.
- Bååth, E. (1996). Adaptation of soil bacterial communities to prevailing pH in different soils. *FEMS Microbiology Ecology*, 19(4), 227-237.
- Bailey, A. D., & Wynn-Williams, D. D. (1982). Soil microbiology studies at Signy Island, South Orkney Islands. *British Antarctic Survey Bulletin*, 51.
- Baker, J. H. (1970). Quantitative studies of yeasts and bacteria in a Signy Island peat. *British Antarctic Survey Bulletin*(23), 51-55.
- Bargagli, R. (2005). *Antarctic Ecosystems: Environmental Contamination, Climate Change, and Human Impact* (Vol. 175): Springer.
- Bargagli, R. (2008). Environmental contamination in Antarctic ecosystems. *Science of the Total Environment*, 400(1-3), 212-226.
- Bargagli, R., Brown, D. H., & Nelli, L. (1995). Metal biomonitoring with mosses: procedures for correcting for soil contamination. *Environmental Pollution*, 89(2), 169-175.
- Bargagli, R., Sanchez-Hernandez, J. C., Martella, L., & Monaci, F. (1998). Mercury, cadmium and lead accumulation in Antarctic mosses growing along nutrient and moisture gradients. *Polar Biology*, 19(5), 316-322.
- Barrett, J. E., Virginia, R. A., Hopkins, D. W., Aislabie, J., Bargagli, R., Bockheim, J. G., et al. (2006a). Terrestrial ecosystem processes of Victoria Land, Antarctica. *Soil Biology & Biochemistry*, 38(10), 3019-3034.
- Barrett, J. E., Virginia, R. A., Wall, D. H., Cary, S. C., Adams, B. J., Hacker, A. L., et al. (2006b). Co-variation in soil biodiversity and biogeochemistry in northern and southern Victoria Land, Antarctica. *Antarctic Science*, 18(4), 535-548.
- Barrett, J. E., Virginia, R. A., Wall, D. H., Parsons, A. N., Powers, L. E., & Burkins, M. B. (2004). Variation in biogeochemistry and soil biodiversity across spatial scales in a polar desert ecosystem. *Ecology*, 85(11), 3105-3118.
- Beare, M., Coleman, D., Crossley, D., Hendrix, P., & Odum, E. (1995). A hierarchical approach to evaluating the significance of soil biodiversity to biogeochemical cycling. *Plant and Soil*, 170(1), 5-22.
- Benson, D. A., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., & Sayers, E. W. (2010). GenBank. *Nucleic Acids Research*, 38, D46-D51.
- Beyer, L., & Bolter, M. (2000). Chemical and biological properties, formation, occurrence and classification of Spodic Cryosols in a terrestrial ecosystem of East Antarctica (Wilkes Land). *Catena*, 39(2), 95-119.

- Beyer, L., Pingpank, K., Wriedt, G., & Bolter, M. (2000). Soil formation in coastal continental Antarctica (Wilkes Land). *Geoderma*, 95(3-4), 283-304.
- Bokhorst, S., Huiskes, A., Convey, P., & Aerts, R. (2007a). The effect of environmental change on vascular plant and cryptogam communities from the Falkland Islands and the Maritime Antarctic. *BMC Ecology*, 7, 15.
- Bokhorst, S., Huiskes, A., Convey, P., & Aerts, R. (2007b). External nutrient inputs into terrestrial ecosystems of the Falkland Islands and the Maritime Antarctic region. *Polar Biology*, 30(10), 1315-1321.
- Bokhorst, S., Huiskes, A., Convey, P., van Bodegom, P. M., & Aerts, R. (2008). Climate change effects on soil arthropod communities from the Falkland Islands and the Maritime Antarctic. *Soil Biology & Biochemistry*, 40(7), 1547-1556.
- Bolter, M. (1992). Environmental-Conditions and Microbiological Properties from Soils and Lichens from Antarctica (Casey-Station, Wilkes-Land). *Polar Biology*, 11(8), 591-599.
- Bolter, M. (2004). Soil: An extreme habitat for microorganisms? *Pedosphere*, 14(2), 137-144.
- Bolter, M., Blume, H. P., Schneider, D., & Beyer, L. (1997). Soil properties and distributions of invertebrates and bacteria from King George Island (Arctowski Station), maritime Antarctic. *Polar Biology*, 18(5), 295-304.
- Bowman, J. P., McCammon, S. A., Gibson, J. A., Robertson, L., & Nichols, P. D. (2003). Prokaryotic metabolic activity and community structure in Antarctic continental shelf sediments. *Applied and Environmental Microbiology*, 69(5), 2448-2462.
- Bridge, P. D., & Newsham, K. K. (2009). Soil fungal community composition at Mars Oasis, a southern maritime Antarctic site, assessed by PCR amplification and cloning. *Fungal Ecology*, 2, 66-74.
- Brussaard, L., Behan-Pelletier, V. M., Bignell, D. E., Brown, V. K., Didden, W., Folgarait, P., et al. (1997). Biodiversity and ecosystem functioning in soil. *Ambio*, 26(8), 563-570.
- Caldwell, M. M., Bornman, J. F., Ballare, C. L., Flint, S. D., & Kulandaivelu, G. (2007). Terrestrial ecosystems, increased solar ultraviolet radiation, and interactions with other climate change factors. *Photochemical & Photobiological Sciences*, 6(3), 252-266.
- Cannone, N., Wagner, D., Hubberten, H. W., & Guglielmin, M. (2008). Biotic and abiotic factors influencing soil properties across a latitudinal gradient in Victoria Land, Antarctica. *Geoderma*, 144(1-2), 50-65.
- Caulkett, A. P., & Ellis-Evans, J. C. (1997). Chemistry of streams of Signy Island, maritime Antarctic: sources of major ions. *Antarctic Science*, 9(1), 3-11.

- Chong, C. W., Dunn, M. J., Convey, P., Tan, G. Y. A., Wong, R. C. S., & Tan, I. K. P. (2009a). Environmental influences on bacterial diversity of soils on Signy Island, maritime Antarctic. *Polar Biology*, 32(11), 1571-1582.
- Chong, C. W., Pearce, D. A., Convey, P., Tan, G. Y. A., Wong, R. C. S., & Tan, I. K. P. (2010). High levels of spatial heterogeneity in the biodiversity of soil prokaryotes on Signy Island, Antarctica. *Soil Biology & Biochemistry*, 42(4), 601-610.
- Chong, C. W., Tan, G. Y. A., Wong, R. C. S., Riddle, M. J., & Tan, I. K. P. (2009b). DGGE fingerprinting of bacteria in soils from eight ecologically different sites around Casey Station, Antarctica. *Polar Biology*, 32(6), 853-860.
- Chown, S. L., & Convey, P. (2007). Spatial and temporal variability across life's hierarchies in the terrestrial Antarctic. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 362(1488), 2307-2331.
- Chu, H., Fierer, N., Lauber, C. L., Caporaso, J. G., Knight, R., & Grogan, P. (2010). Soil bacterial diversity in the Arctic is not fundamentally different from that found in other biomes. *Environmental Microbiology*, 12(11), 2998-3006.
- Claridge, G. G. C., Campbell, I. B., Powell, H. K. J., Amin, Z. H., & Balks, M. R. (1995). Heavy metal contamination in some soils of the McMurdo Sound region, Antarctica. *Antarctic Science*, 7(1), 9-14.
- Clarke, K. R. (1993). Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology*, 18(1), 117-143.
- Cleveland, C., & Liptzin, D. (2007). C:N:P stoichiometry in soil: is there a "Redfield ratio" for the microbial biomass? *Biogeochemistry*, 85(3), 235-252.
- Coenye, T., Laevens, S., Willems, A., Ohlen, M., Hannant, W., Govan, J. R., et al. (2001). *Burkholderia fungorum* sp. nov. and *Burkholderia caledonica* sp. nov., two new species isolated from the environment, animals and human clinical samples. *Int J Syst Evol Microbiol*, 51(Pt 3), 1099-1107.
- Cole, J. R., Wang, Q., Cardenas, E., Fish, J., Chai, B., Farris, R. J., et al. (2009). The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Research*, 37, D141-145.
- Coleman, D. C., & Whitman, W. B. (2005). Linking species richness, biodiversity and ecosystem function in soil systems. *Pedobiologia*, 49(6), 479-497.
- Connon, S. A., & Giovannoni, S. J. (2002). High-throughput methods for culturing microorganisms in very-low-nutrient media yield diverse new marine isolates. *Applied and Environmental Microbiology*, 68(8), 3878-3885.
- Connon, S. A., Lester, E. D., Shafaat, H. S., Obenhuber, D. C., & Ponce, A. (2007). Bacterial diversity in hyperarid Atacama Desert soils. *Journal of Geophysical Research*, 112(G4), G04S17.

- Convey, P. (1996). The influence of environmental characteristics on life history attributes of Antarctic terrestrial biota. *Biological Reviews of the Cambridge Philosophical Society*, 71(2), 191-225.
- Convey, P. (2003). Maritime Antarctic climate change: Signals from terrestrial biology. In E. Domack, A. Leventer, A. Burnett, R. Bindshadler, P. Convey & M. Kirby (Eds.), *Antarctic Peninsula Climate Variability: Historical and Paleoenvironmental Perspectives*, (pp 145-158): American Geophysical Union.
- Convey, P. (2006). Antarctic Climate Change and its Influences on Terrestrial Ecosystems. In D. M. Bergstrom, P. Convey & A. H. L. Huiskes (Eds.), *Trends in Antarctic Terrestrial and Limnetic Ecosystems* (pp. 253-272): Springer Netherlands.
- Convey, P. (2007). Influences on and origins of terrestrial biodiversity of the sub-Antarctic islands. *Papers and Proceedings of the Royal Society of Tasmania* 141, 83-93.
- Convey, P. (2010). Terrestrial biodiversity in Antarctica - Recent advances and future challenges. *Polar Science*, 4(2), 135-147.
- Convey, P., & Smith, R. I. L. (1997). The terrestrial arthropod fauna and its habitats in northern Marguerite Bay and Alexander Island, maritime Antarctic. *Antarctic Science*, 9(1), 12-26.
- Costello, E. K., Halloy, S. R. P., Reed, S. C., Sowell, P., & Schmidt, S. K. (2009). Fumarole-Supported Islands of Biodiversity within a Hyperarid, High-Elevation Landscape on Socompa Volcano, Puna de Atacama, Andes. *Applied and Environmental Microbiology*, 75(3), 735-747.
- Cowan, D. A., Russell, N. J., Mamais, A., & Sheppard, D. M. (2002). Antarctic Dry Valley mineral soils contain unexpectedly high levels of microbial biomass. *Extremophiles*, 6(5), 431-436.
- Culman, S. W., Bukowski, R., Gauch, H. G., Cadillo-Quiroz, H., & Buckley, D. H. (2009). T-REX: software for the processing and analysis of T-RFLP data. *BMC Bioinformatics*, 10, 171.
- Dar, S. A., Kuenen, J. G., & Muyzer, G. (2005). Nested PCR-denaturing gradient gel electrophoresis approach to determine the diversity of sulfate-reducing bacteria in complex microbial communities. *Applied and Environmental Microbiology*, 71(5), 2325-2330.
- Davey, M. C., & Rothery, P. (1993). Primary colonization by microalgae in relation to spatial variation in edaphic factors on Antarctic fell-field soils. *Journal of Ecology*, 81, 335-343.
- de la Torre, J. R., Goebel, B. M., Friedmann, E. I., & Pace, N. R. (2003). Microbial diversity of cryptoendolithic communities from the McMurdo Dry Valleys, Antarctica. *Applied and Environmental Microbiology*, 69(7), 3858-3867.

- Deheyn, D. D., Gendreau, P., Baldwin, R. J., & Latz, M. I. (2005). Evidence for enhanced bioavailability of trace elements in the marine ecosystem of Deception Island, a volcano in Antarctica. [doi: DOI: 10.1016/j.marenvres.2004.08.001]. *Marine Environmental Research*, 60(1), 1-33.
- Delille, D. (2000). Response of Antarctic Soil Bacterial Assemblages to Contamination by Diesel Fuel and Crude Oil. *Microbial Ecology*, 40(2), 159-168.
- Delille, D., Coulon, F., & Pelletier, E. (2007). Long-term changes of bacterial abundance, hydrocarbon concentration and toxicity during a biostimulation treatment of oil-amended organic and mineral sub-Antarctic soils. *Polar Biology*, 30(7), 925-933.
- Delille, D., & Delille, E. (2000). Distribution of Enteric Bacteria in Antarctic Seawater Surrounding the Dumont d'Urville Permanent Station (Ad die Land). [doi: DOI: 10.1016/S0025-326X(00)00077-1]. *Marine Pollution Bulletin*, 40(10), 869-872.
- Dewar, G. J. (1970). *The geology of Adelaide Island*.
- Dick, L. K., & Field, K. G. (2004). Rapid Estimation of Numbers of Fecal Bacteroidetes by Use of a Quantitative PCR Assay for 16S rRNA Genes. *Applied and Environmental Microbiology*, 70(9), 5695-5697.
- Doran, P. T., Priscu, J. C., Lyons, W. B., Walsh, J. E., Fountain, A. G., McKnight, D. M., et al. (2002). Antarctic climate cooling and terrestrial ecosystem response. *Nature*, 415(6871), 517-520.
- Dorigo, U., Volatier, L., & Humbert, J. F. (2005). Molecular approaches to the assessment of biodiversity in aquatic microbial communities. *Water Research*, 39(11), 2207-2218.
- dos Santos, I. R., Silva-Filho, E. V., Schaefer, C., Maria Sella, S., Silva, C. A., Gomes, V., et al. (2006). Baseline mercury and zinc concentrations in terrestrial and coastal organisms of Admiralty Bay, Antarctica. *Environmental Pollution*, 140(2), 304-311.
- Dunbar, J., Ticknor, L. O., & Kuske, C. R. (2000). Assessment of microbial diversity in four southwestern United States soils by 16S rRNA gene terminal restriction fragment analysis. *Applied and Environmental Microbiology*, 66(7), 2943-2950.
- Engelen, A., Convey, P., Hodgson, D. A., Worland, M. R., & Ott, S. (2008). Soil properties of an Antarctic inland site: implications for ecosystem development. *Polar Biology*, 31(12), 1453-1460.
- Enwall, K., & Hallin, S. (2009). Comparison of T-RFLP and DGGE techniques to assess denitrifier community composition in soil. *Letters in Applied Microbiology*, 48(1), 145-148.
- Ettema, C. H., & Wardle, D. A. (2002). Spatial soil ecology. *Trends in Ecology & Evolution*, 17(4), 177-183.

- Excoffier, L., Laval, G., & Schneider, S. (2005). Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online*, 1, 47-50.
- Faith, D. P. (1992). Conservation evaluation and phylogenetic diversity. *Biological Conservation*, 61(1), 1-10.
- Farmer, J. G., Eades, L. J., & Graham, M. C. (1999). The lead content and isotopic composition of british coals and their implications for past and present releases of lead to the UK environment. *Environmental Geochemistry and Health*, 21(3), 257-272.
- Fell, J. W., Scorzetti, G., Connell, L., & Craig, S. (2006). Biodiversity of micro-eukaryotes in Antarctic Dry Valley soils with <5% soil moisture. [doi: DOI: 10.1016/j.soilbio.2006.01.014]. *Soil Biology & Biochemistry*, 38(10), 3107-3119.
- Ferreira, A. C., Nobre, M. F., Moore, E., Rainey, F. A., Battista, J. R., & da Costa, M. S. (1999). Characterization and radiation resistance of new isolates of *Rubrobacter radiotolerans* and *Rubrobacter xylanophilus*. *Extremophiles*, 3(4), 235-238.
- Fierer, N., Bradford, M. A., & Jackson, R. B. (2007). Toward an ecological classification of soil bacteria. *Ecology*, 88(6), 1354-1364.
- Fierer, N., & Jackson, R. B. (2006). The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences of the United States of America*, 103(3), 626-631.
- Flint, H. J., Duncan, S. H., Scott, K. P., & Louis, P. (2007). Interactions and competition within the microbial community of the human colon: links between diet and health. *Environmental Microbiology*, 9(5), 1101-1111.
- Foti, M. J., Sorokin, D. Y., Zacharova, E. E., Pimenov, N. V., Kuenen, J. G., & Muyzer, G. (2008). Bacterial diversity and activity along a salinity gradient in soda lakes of the Kulunda Steppe (Altai, Russia). *Extremophiles*, 12(1), 133-145.
- Franklin, R. B., & Mills, A. L. (2003). Multi-scale variation in spatial heterogeneity for microbial community structure in an eastern Virginia agricultural field. *FEMS Microbiology Ecology*, 44(3), 335-346.
- Franzmann, P. D. (1996). Examination of Antarctic prokaryotic diversity through molecular comparisons. *Biodiversity and Conservation*, 5(11), 1295-1305.
- Franzmann, P. D., & Dobson, S. J. (1993). The phylogeny of bacteria from a modern Antarctic refuge. *Antarctic Science*, 5(3), 267-270.
- Fraschetti, S., Terlizzi, A., & Benedetti-Cecchi, L. (2005). Patterns of distribution of marine assemblages from rocky shores: evidence of relevant scales of variation. *Marine Ecology Progress Series*, 296, 13-29.

- Frenot, Y., Chown, S. L., Whinam, J., Selkirk, P. M., Convey, P., Skotnicki, M., et al. (2005). Biological invasions in the Antarctic: extent, impacts and implications. *Biological Reviews*, 80(1), 45-72.
- Gadd, G. M., & Griffiths, A. J. (1977). Microorganisms and heavy metal toxicity. *Microbial Ecology*, 4(4), 303-317.
- Gafan, G. P., Lucas, V. S., Roberts, G. J., Petrie, A., Wilson, M., & Spratt, D. A. (2005). Statistical analyses of complex denaturing gradient gel electrophoresis profiles. *J Clinical Microbiology*, 43(8), 3971-3978.
- Gillan, D. C., Danis, B., Pernet, P., Joly, G., & Dubois, P. (2005). Structure of sediment-associated microbial communities along a heavy-metal contamination gradient in the marine environment. *Applied and Environmental Microbiology*, 71(2), 679-690.
- Green, J., & Bohannan, B. J. M. (2006). Spatial scaling of microbial biodiversity. *Trends in Ecology & Evolution*, 21(9), 501-507.
- Green, K., & Williams, R. (1986). Observations on food remains in faeces of elephant, leopard and crabeater seals. *Polar Biology*, 6(1), 43-45.
- Grundmann, G. L. (2004). Spatial scales of soil bacterial diversity--the size of a clone. *FEMS Microbiology Ecology*, 48(2), 119-127.
- Guglielmin, M., Evans, C. J. E., & Cannone, N. (2008). Active layer thermal regime under different vegetation conditions in permafrost areas. A case study at Signy Island (Maritime Antarctica). *Geoderma*, 144(1-2), 73-85.
- Gurtler, V., & Stanisich, V. A. (1996). New approaches to typing and identification of bacteria using the 16s-23s rDNA spacer region. *Microbiology-Sgm*, 142, 3-16.
- Hamady, M., Lozupone, C., & Knight, R. (2010). Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data. *ISME Journal*, 4(1), 17-27.
- Hartman, W. H., Richardson, C. J., Vilgalys, R., & Bruland, G. L. (2008). Environmental and anthropogenic controls over bacterial communities in wetland soils. *Proceedings of the National Academy of Sciences of the United States of America*, 105(46), 17842-17847.
- Hartmann, M., & Widmer, F. (2006). Community structure analyses are more sensitive to differences in soil bacterial communities than anonymous diversity indices. *Appl Environmental Microbiology*, 72(12), 7804-7812.
- Head, I. M., Saunders, J. R., & Pickup, R. W. (1998). Microbial Evolution, Diversity, and Ecology: A Decade of Ribosomal RNA Analysis of Uncultivated Microorganisms. *Microbial Ecology*, 35(1), 1-21.

- Hill, G. T., Mitkowski, N. A., Aldrich-Wolfe, L., Emele, L. R., Jurkonie, D. D., Ficke, A., et al. (2000). Methods for assessing the composition and diversity of soil microbial communities.. *Applied Soil Ecology*, 15(1), 25-36.
- Hinojosa, M. B., Carreira, J. A., Garc ía-Ru íz, R., & Dick, R. P. (2005). Microbial Response to Heavy Metal-Polluted Soils. *Journal of Environmental Quality*, 34(5), 1789-1800.
- Hogg, I. D., Cary, S. C., Convey, P., Newsham, K. K., O'Donnell, A. G., Adams, B. J., et al. (2006). Biotic interactions in Antarctic terrestrial ecosystems: Are they a factor? *Soil Biology & Biochemistry*, 38(10), 3035-3040.
- Holdgate, M. W. (1977). Terrestrial Ecosystems in the Antarctic. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, 279(963), 5-25.
- Honda, K., Yamamoto, Y., & Tatsukawa, R. (1987). Distribution of heavy metals in Antarctic marine ecosystem. *Proceeding of NIPR Symposium on Polar Biology*, 1, 184-197.
- Hopkins, D. W., Sparrow, A. D., Shillam, L. L., English, L. C., Dennis, P. G., Novis, P., et al. (2008). Enzymatic activities and microbial communities in an Antarctic dry valley soil: Responses to C and N supplementation. *Soil Biology & Biochemistry*, 40(9), 2130-2136.
- Horner-Devine, M. C., & Bohannan, B. J. (2006). Phylogenetic clustering and overdispersion in bacterial communities. *Ecology*, 87(7 Suppl), S100-108.
- Horner-Devine, M. C., Carney, K. M., & Bohannan, B. J. (2004a). An ecological perspective on bacterial biodiversity. *Proc Biol Sci*, 271(1535), 113-122.
- Horner-Devine, M. C., Lage, M., Hughes, J. B., & Bohannan, B. J. (2004b). A taxa-area relationship for bacteria. *Nature*, 432(7018), 750-753.
- Hughes, K. A., & Lawley, B. (2003). A novel Antarctic microbial endolithic community within gypsum crusts. *Environmental Microbiology*, 5(7), 555-565.
- Hughes, K. A., McCartney, H. A., Lachlan-Cope, T. A., & Pearce, D. A. (2004). A preliminary study of airborne microbial biodiversity over peninsular Antarctica. *Cellular and Molecular Biology*, 50(5), 537-542.
- Janssen, P. H. (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Applied and Environmental Microbiology*, 72(3), 1719-1728.
- Johnson, M., Zaretskaya, I., Raytselis, Y., Merezuk, Y., McGinnis, S., & Madden, T. L. (2008). NCBI BLAST: a better web interface. *Nucleic Acids Research*, 36(Web Server issue), W5-9.
- Kaplan, C. W., & Kitts, C. L. (2003). Variation between observed and true Terminal Restriction Fragment length is dependent on true TRF length and purine content. *Journal of Microbiological Methods*, 54(1), 121-125.

- Kim, J.-S., Sparovek, G., Longo, R. M., De Melo, W. J., & Crowley, D. (2007). Bacterial diversity of terra preta and pristine forest soil from the Western Amazon. [doi: DOI: 10.1016/j.soilbio.2006.08.010]. *Soil Biology & Biochemistry*, 39(2), 684-690.
- Kirk, J. L., Beaudette, L. A., Hart, M., Moutoglis, P., Khironomos, J. N., Lee, H., et al. (2004). Methods of studying soil microbial diversity. *Journal of Microbiological Methods*, 58(2), 169-188.
- Klappenbach, J. A., Dunbar, J. M., & Schmidt, T. M. (2000). rRNA operon copy number reflects ecological strategies of bacteria. *Applied and Environmental Microbiology*, 66(4), 1328-1333.
- Konopka, A., Zakharova, T., Bischoff, M., Oliver, L., Nakatsu, C., & Turco, R. F. (1999). Microbial biomass and activity in lead-contaminated soil. *Applied and Environmental Microbiology*, 65(5), 2256-2259.
- Kowalchuk, G. A., Drigo, B., Yergeau, E., & Veen, J. A. (2006). Assessing Bacterial and Fungal Community Structure in Soil Using Ribosomal RNA and Other Structural Gene Markers. In P. Nannipieri & K. Smalla (Eds.), *Nucleic Acids and Proteins in Soil* (Vol. 8, pp. 159-188): Springer Berlin Heidelberg.
- Kumar, S., Tamura, K., & Nei, M. (2004). MEGA3: Integrated software for Molecular Evolutionary Genetics Analysis and sequence alignment. *Briefings in Bioinformatics*, 5(2), 150-163.
- Kunito, T., Saeki, K., Oyaizu, H., & Matsumoto, S. (1999). Influences of copper forms on the toxicity to microorganisms in soils. *Ecotoxicol Environ Saf*, 44(2), 174-181.
- Lawley, B., Ripley, S., Bridge, P., & Convey, P. (2004). Molecular analysis of geographic patterns of eukaryotic diversity in Antarctic soils. *Applied and Environmental Microbiology*, 70(10), 5963-5972.
- Li, S., Xiao, X., Yin, X., & Wang, F. (2006). Bacterial community along a historic lake sediment core of Ardley Island, west Antarctica. *Extremophiles*, 10(5), 461-467.
- Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J. P., Hector, A., et al. (2001). Ecology - Biodiversity and ecosystem functioning: Current knowledge and future challenges. *Science*, 294(5543), 804-808.
- Lozupone, C., Hamady, M., & Knight, R. (2006). UniFrac--an online tool for comparing microbial community diversity in a phylogenetic context. *BMC Bioinformatics*, 7, 371.
- Lozupone, C. A., & Knight, R. (2008). Species divergence and the measurement of microbial diversity. *FEMS Microbiology Reviews*, 32(4), 557-578.
- Lynnes, A. S., Reid, K., & Croxall, J. P. (2004). Diet and reproductive success of Adélie and chinstrap penguins: linking response of predators to prey population dynamics. *Polar Biology*, 27(9), 544-554.

- Malandrino, M., Abollino, O., Buoso, S., Casalino, C. E., Gasparon, M., Giacomino, A., et al. (2009). Geochemical characterisation of Antarctic soils and lacustrine sediments from Terra Nova Bay. *Microchemical Journal*, 92(1), 21-31.
- Malosso, E., Waite, I. S., English, L., Hopkins, D. W., & O'Donnell, A. G. (2006). Fungal diversity in maritime Antarctic soils determined using a combination of culture isolation, molecular fingerprinting and cloning techniques. *Polar Biology*, 29(7), 552-561.
- Männistö, M. K., Tirola, M., & Häggblom, M. M. (2007). Bacterial communities in Arctic fjelds of Finnish Lapland are stable but highly pH-dependent. *FEMS Microbiology Ecology*, 59(2), 452-465.
- Margesin, R., Sproer, C., Schumann, P., & Schinner, F. (2003). *Pedobacter cryoconitis* sp. nov., a facultative psychrophile from alpine glacier cryoconite. *International Journal of Systematic and Evolutionary Microbiology*, 53(5), 1291-1296.
- Martin, A. P. (2002). Phylogenetic approaches for describing and comparing the diversity of microbial communities. *Applied and Environmental Microbiology*, 68(8), 3673-3682.
- Martiny, J. B., Bohannan, B. J., Brown, J. H., Colwell, R. K., Fuhrman, J. A., Green, J. L., et al. (2006). Microbial biogeography: putting microorganisms on the map. *Nature Reviews Microbiology*, 4(2), 102-112.
- Maslen, N. R., & Convey, P. (2006). Nematode diversity and distribution in the southern maritime Antarctic - clues to history? *Soil Biology & Biochemistry*, 38(10), 3141-3151.
- Melick, D. R., Hovenden, M. J., & Seppelt, R. D. (1994). Phytogeography of bryophyte and lichen vegetation in the Windmill Islands, Wilkes Land, Continental Antarctica. *Plant Ecology*, 111(1), 71-87.
- Mengoni, A., Giuntini, E., & Bazzicalupo, M. (2007). Application of Terminal-Restriction Fragment Length Polymorphism for Molecular Analysis of Soil Bacterial Communities. In A. Varma & R. Oelmüller (Eds.), *Advanced Techniques in Soil Microbiology* (Vol. 11, pp. 295-305): Springer Berlin Heidelberg.
- Michel, R. F. M., Schaefer, C. E. G. R., Dias, L. E., Simas, F. N. B., de Melo Benites, V., & de Sá Mendonça, E. (2006). Ornithogenic Gelisols (Cryosols) from Maritime Antarctica. *Soil Science Society of America Journal*, 70(4), 1370-1376.
- Milius, N. (2000). The birds of Rothera, Adelaide Island, Antarctic Peninsula. *Marine Ornithology*, 28, 63-67.
- Mizutani, H., & Wada, E. (1988). Nitrogen and carbon isotope ratios in seabird rookeries and their ecological implications. *Ecology*, 69, 340-349.

- Moncrieff, A. C. M., & Kelly, S. R. A. (1993). Lithostratigraphy of the uppermost Fossil Bluff Group (Early Cretaceous) of Alexander Island, Antarctica: history of an Albian regression. *Cretaceous Research*, 14(1), 1-15.
- Moosvi, S. A., McDonald, I. R., Pearce, D. A., Kelly, D. P., & Wood, A. P. (2005). Molecular detection and isolation from Antarctica of methylotrophic bacteria able to grow with methylated sulfur compounds. *Systematic and Applied Microbiology*, 28(6), 541-554.
- Mulder, E., & Deinema, M. (2006). The Genus *Haliscomenobacter*. In M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer & E. Stackebrandt (Eds.), *The Prokaryotes* (pp. 602-604): Springer New York.
- Muradian, R. (2001). Ecological thresholds: a survey. *Ecological Economics*, 38(1), 7-24.
- Muyzer, G. (1999). DGGE/TGGE a method for identifying genes from natural ecosystems. *Current Opinion in Microbiology*, 2(3), 317-322.
- Muyzer, G., de Waal, E. C., & Uitterlinden, A. G. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology*, 59(3), 695-700.
- Muyzer, G., & Smalla, K. (1998). Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology*, 73(1), 127-141.
- Nakatsu, C. H. (2007). Soil microbial community analysis using denaturing gradient gel electrophoresis. *Soil Science Society of America Journal*, 71(2), 562-571.
- Nakatsu, C. H., Torsvik, V., & Ovreas, L. (2000). Soil community analysis using DGGE of 16S rDNA polymerase chain reaction products. *Soil Science Society of America Journal*, 64(4), 1382-1388.
- Nannipieri, P., Ascher, J., Ceccherini, M. T., Landi, L., Pietramellara, G., & Renella, G. (2003). Microbial diversity and soil functions. *European Journal of Soil Science*, 54(4), 655-670.
- Newberry, C. J., Webster, G., Cragg, B. A., Parkes, R. J., Weightman, A. J., & Fry, J. C. (2004). Diversity of prokaryotes and methanogenesis in deep subsurface sediments from the Nankai Trough, Ocean Drilling Program Leg 190. *Environmental Microbiology*, 6(3), 274-287.
- Newsham, K. K., Pearce, D. A., & Bridge, P. D. (2010). Minimal influence of water and nutrient content on the bacterial community composition of a maritime Antarctic soil. *Microbiological Research*, 165(7), 523-530.

- Niederberger, T. D., McDonald, I. R., Hacker, A. L., Soo, R. M., Barrett, J. E., Wall, D. H., et al. (2008). Microbial community composition in soils of Northern Victoria Land, Antarctica. *Environmental Microbiology*, 10(7), 1713-1724.
- Nocker, A., Burr, M., & Camper, A. K. (2007). Genotypic microbial community profiling: A critical technical review. *Microbial Ecology*, 54(2), 276-289.
- Nunan, N., Wu, K., Young, I. M., Crawford, J. W., & Ritz, K. (2002). In situ spatial patterns of soil bacterial populations, mapped at multiple scales, in an arable soil. *Microbial Ecology*, 44(4), 296-305.
- Nygård, T., Lie, E., Røy, N., & Steinnes, E. (2001). Metal Dynamics in an Antarctic Food Chain. [doi: DOI: 10.1016/S0025-326X(00)00206-X]. *Marine Pollution Bulletin*, 42(7), 598-602.
- O'Donnell, A. G., & Göres, H. E. (1999). 16S rDNA methods in soil microbiology. [doi: DOI: 10.1016/S0958-1669(99)80039-1]. *Current Opinion in Biotechnology*, 10(3), 225-229.
- Oren, A. (2006). The Genera *Rhodothermus*, *Thermonema*, *Hymenobacter* and *Salinibacter*. In M. Dworkin, S. Falkow, E. Rosenberg, K.-H. Schleifer & E. Stackebrandt (Eds.), *The Prokaryotes* (pp. 712-738): Springer New York.
- Overmann, J., & van Gernerden, H. (2000). Microbial interactions involving sulfur bacteria: implications for the ecology and evolution of bacterial communities. *FEMS Microbiology Reviews*, 24(5), 591-599.
- Ovreas, L. (2000). Population and community level approaches for analysing microbial diversity in natural environments. *Ecology Letters*, 3(3), 236-251.
- Pearce, D. A. (2003). Bacterioplankton community structure in a maritime Antarctic oligotrophic lake during a period of holomixis, as determined by denaturing gradient gel electrophoresis (DGGE) and fluorescence in situ hybridization (FISH). *Microbial Ecology*, 46(1), 92-105.
- Pearce, D. A., Bridge, P. D., Hughes, K. A., Sattler, B., Psenner, R., & Russell, N. J. (2009). Microorganisms in the atmosphere over Antarctica. *FEMS Microbiology Ecology*, 69(2), 143-157.
- Pearce, D. A., Hughes, K. A., Lachlan-Cope, T., Harangozo, S. A., & Jones, A. E. (2010). Biodiversity of air-borne microorganisms at Halley station, Antarctica. *Extremophiles*, 14(2), 145-159.
- Pearce, D. A., van der Gast, C. J., Lawley, B., & Ellis-Evans, J. C. (2003). Bacterioplankton community diversity in a maritime Antarctic lake, determined by culture-dependent and culture-independent techniques. *FEMS Microbiology Ecology*, 45(1), 59-70.

- Peck, L. S., Clark, M. S., Clarke, A., Cockell, C. S., Convey, P., Detrich, H. W., et al. (2005). Genomics: applications to Antarctic ecosystems. *Polar Biology*, 28(5), 351-365.
- Pointing, S. B., Chan, Y., Lacap, D. C., Lau, M. C. Y., Jurgens, J. A., & Farrell, R. L. (2010). Highly specialized microbial diversity in hyper-arid polar desert (vol 106, pg 19964, 2009). *Proceedings of the National Academy of Sciences of the United States of America*, 107(3), 1254-1254.
- Potts, M. (1994). Desiccation tolerance of prokaryotes. *Microbiological Review*, 58(4), 755-805.
- Powell, S. M., Bowman, J. P., Snape, I., & Stark, J. S. (2003). Microbial community variation in pristine and polluted nearshore Antarctic sediments. *FEMS Microbiology Ecology*, 45(2), 135-145.
- Powell, S. M., Riddle, M. J., Snape, I., & Stark, J. S. (2005a). Location and DGGE methodology can influence interpretation of field experimental studies on the response to hydrocarbons by Antarctic benthic microbial community. *Antarctic Science*, 17(3), 353-360.
- Powell, S. M., Snape, I., Bowman, J. P., Thompson, B. A. W., Stark, J. S., McCammon, S. A., et al. (2005b). A comparison of the short term effects of diesel fuel and lubricant oils on Antarctic benthic microbial communities. [doi: DOI: 10.1016/j.jembe.2005.02.005]. *Journal of Experimental Marine Biology and Ecology*, 322(1), 53-65.
- Ramette, A. (2007). Multivariate analyses in microbial ecology. *FEMS Microbiology Ecology*, 62(2), 142-160.
- Redfield, A. C. (1958). The biological control of chemical factors in the environment. *American Scientist*, 46, 205-221.
- Reid, K. (1995). The diet of Antarctic fur seals (*Arctocephalus gazella* Peters 1875) during winter at South Georgia. *Antarctic Science*, 7, 241-249.
- Revill, A. T., Snape, I., Lucieer, A., & Guille, D. (2007). Constraints on transport and weathering of petroleum contamination at Casey Station, Antarctica. [doi: DOI: 10.1016/j.coldregions.2007.01.001]. *Cold Regions Science and Technology*, 48(2), 154-167.
- Riesenfeld, C. S., Schloss, P. D., & Handelsman, J. (2004). Metagenomics: genomic analysis of microbial communities. *Annual Review of Genetics*, 38, 525-552.
- Roser, D. J., Seppelt, R. D., & Ashbolt, N. (1993). Microbiology of ornithogenic soils from the windmill islands, budd coast, continental Antarctica: Microbial biomass distribution. [doi: DOI: 10.1016/0038-0717(93)90023-5]. *Soil Biology & Biochemistry*, 25(2), 165-175.

- Rosswall, T. (1982). Microbiological Regulation of the Biogeochemical Nitrogen-Cycle. *Plant and Soil*, 67(1-3), 15-34.
- Rothschild, L. J., & Mancinelli, R. L. (2001). Life in extreme environments. *Nature*, 409(6823), 1092-1101.
- Salam, A. K., & Helmke, P. A. (1998). The pH dependence of free ionic activities and total dissolved concentrations of copper and cadmium in soil solution. *Geoderma*, 83(3-4), 281-291.
- Santos, I. R., Silva-Filho, E. V., Schaefer, C. E. G. R., Albuquerque-Filho, M. R., & Campos, L. S. (2005). Heavy metal contamination in coastal sediments and soils near the Brazilian Antarctic Station, King George Island. *Marine Pollution Bulletin*, 50(2), 185-194.
- Sattin, S., Cleveland, C., Hood, E., Reed, S., King, A., Schmidt, S., et al. (2009). Functional shifts in unvegetated, perhumid, recently-deglaciated soils do not correlate with shifts in soil bacterial community composition. *The Journal of Microbiology*, 47(6), 673-681.
- Saul, D. J., Aislabie, J. M., Brown, C. E., Harris, L., & Foght, J. M. (2005). Hydrocarbon contamination changes the bacterial diversity of soil from around Scott Base, Antarctica. *FEMS Microbiology Ecology*, 53(1), 141-155.
- Schloss, P. D. (2008). Evaluating different approaches that test whether microbial communities have the same structure. *ISME Journal*, 2(3), 265-275.
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., et al. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology*, 75(23), 7537-7541.
- Schneegurt, M. A., Dore, S. Y., & Kulpa, C. F., Jr. (2003). Direct extraction of DNA from soils for studies in microbial ecology. *Current Issues in Molecular Biology*, 5(1), 1-8.
- Schütte, U., Abdo, Z., Bent, S., Shyu, C., Williams, C., Pierson, J., et al. (2008). Advances in the use of terminal restriction fragment length polymorphism (T-RFLP) analysis of 16S rRNA genes to characterize microbial communities. *Applied Microbiology and Biotechnology*, 80(3), 365-380.
- Scouller, R. C., Snape, I., Stark, J. S., & Gore, D. B. (2006). Evaluation of geochemical methods for discrimination of metal contamination in Antarctic marine sediments: A case study from Casey Station. *Chemosphere*, 65(2), 294-309.
- Sheppard, D. S., Claridge, G. G. C., & Campbell, I. B. (2000). Metal contamination of soils at Scott Base, Antarctica. *Applied Geochemistry*, 15(4), 513-530.
- Shigematsu, T., Yumihara, K., Ueda, Y., Numaguchi, M., Morimura, S., & Kida, K. (2003). *Delftia tsuruhatensis* sp. nov., a terephthalate-assimilating bacterium isolated from

activated sludge. *International Journal of Systematic and Evolutionary Microbiology*, 53(Pt 5), 1479-1483.

- Shravage, B. V., Dayananda, K. M., Patole, M. S., & Shouche, Y. S. (2007). Molecular microbial diversity of a soil sample and detection of ammonia oxidizers from Cape Evans, Mcmurdo Dry Valley, Antarctica. *Microbiological Research*, 162(1), 15-25.
- Shyu, C., Soule, T., Bent, S. J., Foster, J. A., & Forney, L. J. (2007). MiCA: a web-based tool for the analysis of microbial communities based on terminal-restriction fragment length polymorphisms of 16S and 18S rRNA genes. *Microbial Ecology*, 53(4), 562-570.
- Simas, F. N. B., Schaefer, C. E. G. R., Melo, V. F., Albuquerque-Filho, M. R., Michel, R. F. M., Pereira, V. V., et al. (2007). Ornithogenic cryosols from Maritime Antarctica: Phosphatization as a soil forming process. *Geoderma*, 138(3-4), 191-203.
- Smalla, K., Oros-Sichler, M., Milling, A., Heuer, H., Baumgarte, S., Becker, R., et al. (2007). Bacterial diversity of soils assessed by DGGE, T-RFLP and SSCP fingerprints of PCR-amplified 16S rRNA gene fragments: do the different methods provide similar results? *Journal of Microbiological Methods*, 69(3), 470-479.
- Smith, C. J., Danilowicz, B. S., Clear, A. K., Costello, F. J., Wilson, B., & Meijer, W. G. (2005). T-Align, a web-based tool for comparison of multiple terminal restriction fragment length polymorphism profiles. *FEMS Microbiology Ecology*, 54(3), 375-380.
- Smith, H. G., & Tearle, P. V. (1985). Aspects of microbial and protozoan abundances in Signy Islands fellfields. *British Antarctic Survey Bulletin*(68), 83-90.
- Smith, J. J., Tow, L. A., Stafford, W., Cary, C., & Cowan, D. A. (2006). Bacterial diversity in three different Antarctic cold desert mineral soils. *Microbial Ecology*, 51(4), 413-421.
- Smith, R. I. L. (1972). Vegetation of the South Orkney Islands with particular reference to Signy Island. *British Antarctic Survey Scientific Reports*, 68, 124.
- Smith, R. I. L. (1984). Terrestrial plant biology of the subantarctic and Antarctic. In R. M. Laws (Ed.), *Antarctic Ecology* (Vol. 1, pp. 61-162). London: Academic Press.
- Smith, R. I. L. (1988a). Bryophyte oases in ablation valleys on Alexander Island, Antarctica. *Bryologist*, 91, 45-50.
- Smith, R. I. L. (1988b). Destruction of antarctic terrestrial ecosystems by a rapidly increasing fur seal population. *Biological Conservation*, 45(1), 55-72.
- Smith, R. I. L. (1990). Signy Island as a paradigm of biological and environmental change in Antarctic terrestrial ecosystems. In K. R. Kerry & G. Hempel (Eds.), *Antarctic Ecosystems Ecological Change and Conservation* (pp. 32-50). Berlin: Springer-Verlag.

- Smith, V. R. (2005). Moisture, carbon and inorganic nutrient controls of soil respiration at a sub-Antarctic island. [doi: DOI: 10.1016/j.soilbio.2004.07.026]. *Soil Biology & Biochemistry*, 37(1), 81-91.
- Snape, I., Ferguson, S. H., Harvey, P. M., & Riddle, M. J. (2006). Investigation of evaporation and biodegradation of fuel spills in Antarctica: II--Extent of natural attenuation at Casey Station. *Chemosphere*, 63(1), 89-98.
- Snape, I., Scouller, R. C., Stark, S. C., Stark, J., Riddle, M. J., & Gore, D. B. (2004). Characterisation of the dilute HCl extraction method for the identification of metal contamination in Antarctic marine sediments. *Chemosphere*, 57(6), 491-504.
- Solaiman, Z., & Marschner, P. (2007). DGGE and RISA Protocols for Microbial Community Analysis in Soil. In A. Varma & R. Oelmüller (Eds.), *Advanced Techniques in Soil Microbiology* (Vol. 11, pp. 167-180): Springer Berlin Heidelberg.
- Stackebrandt, E., & Goebel, B. M. (1994). Taxonomic Note: A Place for DNA-DNA Reassociation and 16S rRNA Sequence Analysis in the Present Species Definition in Bacteriology. *International Journal of Systematic Bacteriology*, 44(4), 846-849.
- Stark, J. S. (2000). The distribution and abundance of soft-sediment macrobenthos around Casey Station, East Antarctica. *Polar Biology*, 23(12), 840-850.
- Stark, J. S., Riddle, M. J., Snape, I., & Scouller, R. C. (2003). Human impacts in Antarctic marine soft-sediment assemblages: correlations between multivariate biological patterns and environmental variables at Casey Station. [doi: DOI: 10.1016/S0272-7714(02)00291-3]. *Estuarine, Coastal and Shelf Science*, 56(3-4), 717-734.
- Stark, J. S., Snape, I., Riddle, M. J., & Stark, S. C. (2005). Constraints on spatial variability in soft-sediment communities affected by contamination from an Antarctic waste disposal site. *Marine Pollution Bulletin*, 50(3), 276-290.
- Sugden, D. E., & Clapperton, C. M. (1981). An ice-shelf moraine, George VI Sound, Antarctica. *Annals of Glaciology*, 2, 135-141.
- Tamura, K., Dudley, J., Nei, M., & Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, 24(8), 1596-1599.
- Tejedo, P., Justel, A., Benayas, J., Rico, E., Convey, P., & Quesada, A. (2009). Soil trampling in an Antarctic Specially Protected Area: tools to assess levels of human impact. *Antarctic Science*, 21(3), 229-236.
- Thangaraj, K., Kapley, A., & Purohit, H. J. (2008). Characterization of diverse *Acinetobacter* isolates for utilization of multiple aromatic compounds. [doi: DOI: 10.1016/j.biortech.2007.04.053]. *Bioresource Technology*, 99(7), 2488-2494.
- Tin, T., Fleming, Z. L., Hughes, K. A., Ainley, D. G., Convey, P., Moreno, C. A., et al. (2009). Impacts of local human activities on the Antarctic environment. *Antarctic Science*, 21(1), 3-33.

- Tindall, B. J. (2004). Prokaryotic diversity in the Antarctic: The tip of the iceberg. *Microbial Ecology*, 47(3), 271-283.
- Torsvik, V., & Ovreas, L. (2002). Microbial diversity and function in soil: from genes to ecosystems. *Current Opinion in Microbiology*, 5(3), 240-245.
- Torsvik, V., Ovreas, L., & Thingstad, T. F. (2002). Prokaryotic diversity - Magnitude, dynamics, and controlling factors. *Science*, 296(5570), 1064-1066.
- Tow, L. A., & Cowan, D. A. (2005). Dissemination and survival of non-indigenous bacterial genomes in pristine Antarctic environments. *Extremophiles*, 9(5), 385-389.
- Townsend, A. T., & Snape, I. (2008). Multiple Pb sources in marine sediments near the Australian Antarctic Station, Casey. *Science of the Total Environment*, 389(2-3), 466-474.
- Turner, J., Colwell, S. R., Marshall, G. J., Lachlan-Cope, T. A., Carleton, A. M., Jones, P. D., et al. (2005). Antarctic climate change during the last 50 years. *International Journal of Climatology*, 25(3), 279-294.
- Ugolini, F. C., & Bockheim, J. G. (2008). Antarctic soils and soil formation in a changing environment: A review. *Geoderma*, 144(1-2), 1-8.
- Usher, M. B., & Booth, R. G. (1986). Arthropod communities in a maritime Antarctic moss-turf habitat: multiple scales of pattern in the mites and Collembola. *Journal of Animal Ecology*, 55, 155-170.
- Vamosi, S. M., Heard, S. B., Vamosi, J. C., & Webb, C. O. (2009). Emerging patterns in the comparative analysis of phylogenetic community structure. *Molecular Ecology*, 18(4), 572-592.
- Vaughan, D. G., Marshall, G. J., Connolley, W. M., Parkinson, C., Mulvaney, R., Hodgson, D. A., et al. (2003). Recent rapid regional climate warming on the Antarctic Peninsula. *Climatic Change*, 60(3), 243-274.
- Vincent, W. F. (2000). Evolutionary origins of Antarctic microbiota: invasion, selection and endemism. *Antarctic Science*, 12(3), 374-385.
- Vyverman, W., Verleyen, E., Wilmotte, A., Hodgson, D. A., Willems, A., Peeters, K., et al. (2010). Evidence for widespread endemism among Antarctic micro-organisms. [doi: DOI: 10.1016/j.polar.2010.03.006]. *Polar Science*, 4(2), 103-113.
- Wall, D. H. (2005). Biodiversity and ecosystem functioning in terrestrial habitats of Antarctica. *Antarctic Science*, 17(4), 523-531.
- Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., et al. (2002). Ecological responses to recent climate change. *Nature*, 416(6879), 389-395.
- Ward, N. L., Challacombe, J. F., Janssen, P. H., Henrissat, B., Coutinho, P. M., Wu, M., et al. (2009). Three Genomes from the Phylum Acidobacteria Provide Insight into the

- Lifestyles of These Microorganisms in Soils. *Applied and Environmental Microbiology*, 75(7), 2046-2056.
- Wardle, D. A. (1998). Controls of temporal variability of the soil microbial biomass: A global-scale synthesis. *Soil Biology & Biochemistry*, 30(13), 1627-1637.
- Watanabe, K., Nelson, J., Harayama, S., & Kasai, H. (2001). ICB database: the gyrB database for identification and classification of bacteria. *Nucleic Acids Research*, 29(1), 344-345.
- Webb, C. O., Ackerly, D. D., & Kembel, S. W. (2008). Phylocom: software for the analysis of phylogenetic community structure and trait evolution. *Bioinformatics*, 24(18), 2098-2100.
- Webb, C. O., Ackerly, D. D., McPeck, M. A., & Donoghue, M. J. (2002). Phylogenies and community ecology. *Annual Review of Ecology and Systematics*, 33, 475-505.
- White, C., Sharman, A. K., & Gadd, G. M. (1998). An integrated microbial process for the bioremediation of soil contaminated with toxic metals. *Nature Biotechnology*, 16(6), 572-575.
- Wichern, J., Wichern, F., & Joergensen, R. G. (2006). Impact of salinity on soil microbial communities and the decomposition of maize in acidic soils. [doi: DOI: 10.1016/j.geoderma.2006.08.001]. *Geoderma*, 137(1-2), 100-108.
- Wiens, J. A. (1989). Spatial Scaling in Ecology. *Functional Ecology*, 3(4), 385-397.
- Wiens, J. J., Ackerly, D. D., Allen, A. P., Anacker, B. L., Buckley, L. B., Cornell, H. V., et al. (2010). Niche conservatism as an emerging principle in ecology and conservation biology. *Ecology Letters*, 13(10), 1310-1324.
- Williams, R. A. (1969). Bacteriological survey in Antarctica. *British Antarctic Survey Bulletin*, 19, 97-98.
- Wynn-Williams, D. D. (1990). Microbial colonization processes in Antarctic fellfield soils - an experimental overview. *Proceedings of the NIPR Symposium on Polar Biology*, 3, 164-178.
- Wynn-Williams, D. D. (1996). Antarctic microbial diversity: The basis of polar ecosystem processes. *Biodiversity and Conservation*, 5(11), 1271-1293.
- Xiao, X., Li, M., You, Z., & Wang, F. (2007). Bacterial communities inside and in the vicinity of the Chinese Great Wall Station, King George Island, Antarctica. *Antarctic Science*, 19, 11-16.
- Xie, C.-H., & Yokota, A. (2006). Reclassification of [Flavobacterium] ferrugineum as Terrimonas ferruginea gen. nov., comb. nov., and description of Terrimonas lutea sp. nov., isolated from soil. *International Journal of Systematic and Evolutionary Microbiology* 56(5), 1117-1121.

- Yamamoto, Y., Honda, K., & Tatsukawa, R. (1987). Heavy metal accumulation in Antarctic krill *Euphausia Superba*. *Proceeding of NIPR Symposium on Polar Biology*, 1, 198-204.
- Yergeau, E., Bokhorst, S., Huiskes, A. H. L., Boschker, H. T. S., Aerts, R., & Kowalchuk, G. A. (2007a). Size and structure of bacterial, fungal and nematode communities along an Antarctic environmental gradient. *FEMS Microbiology Ecology*, 59(2), 436-451.
- Yergeau, E., Newsham, K. K., Pearce, D. A., & Kowalchuk, G. A. (2007b). Patterns of bacterial diversity across a range of Antarctic terrestrial habitats. *Environmental Microbiology*, 9(11), 2670-2682.
- Yergeau, E., Schoondermark-Stolk, S. A., Brodie, E. L., Dejean, S., DeSantis, T. Z., Goncalves, O., et al. (2009). Environmental microarray analyses of Antarctic soil microbial communities. *ISME Journal*, 3(3), 340-351.
- Yin, X. B., Xia, L. J., Sun, L. G., Luo, H. H., & Wang, Y. H. (2008). Animal excrement: A potential biomonitor of heavy metal contamination in the marine environment. *Science of the Total Environment*, 399(1-3), 179-185.
- Yoon, M.-H., & Im, W.-T. (2007). *Flavisolibacter ginsengiterrae* gen. nov., sp. nov. and *Flavisolibacter ginsengisoli* sp. nov., isolated from ginseng cultivating soil. *Int J Syst Evol Microbiol*, 57(8), 1834-1839.
- Youssef, N. H., & Elshahed, M. S. (2008). Diversity rankings among bacterial lineages in soil. *ISME Journal*, 3(3), 305-313.
- Zdanowski, M. K., Weglenski, P., Golik, P., Sasin, J. M., Borsuk, P., Zmuda, M. J., et al. (2004). Bacterial diversity in Adelie penguin, *Pygoscelis adeliae*, guano: molecular and morpho-physiological approaches. *FEMS Microbiology Ecology*, 50(3), 163-173.