

REFERENCES

REFERENCES

- Abdullah, Z., Hussain, K., Zhari, I., Rasadah, M. A., Mazura, P., Jamaludin, F., *et al.* (2009). Evaluation of extracts of leaf of three *Ficus deltoidea* varieties for antioxidant activities and secondary metabolites. *Pharmacognosy Research* 1, 216-223.
- Abdullah-Soheimi, S. S., Lim, B.-K., Hashim, O. H., & Shui, A. S. (2010). Patients with ovarian carcinoma excrete different altered levels of urine CD59, kininogen-1 and fragments of inter-alpha-trypsin inhibitor heavy chain H4 and albumin. *Proteome Science*, 58, 8.
- Adam, Z., Hamid, M., Ismail, A., & Khamis, S. (2007). Effect of *Ficus deltoidea* aqueous extract on blood glucose level in normal and mild diabetic rats. *Malaysian Journal of Health Science*, 5, 9-16.
- Adam, Z., Hamid, M., Ismail, A., & Khamis, S. (2009). Effect of *Ficus deltoidea* extracts on hepatic basal and insulin-stimulated glucose uptake. *Journal of Biological Sciences*, 9(8), 796-803.
- Alderman, M., Cohen, H., Sealey, J., & Laragh, J. (2010). Pressor responses to antihypertensive drug types. *American Journal of Hypertension*, 23, 1031-1037.
- Aller, M. a.-A., Arias, J.-L., Cruz, A., & Arias, J. (2007). Inflammation: a way to understanding the evalution of portal hypertension. *Theoretical Biology and Medical Modelling*, 4(44), 1-25.
- Amacher, D. E., Adler, R., Herath, A., & Townsend, R. R. (2005). Use of proteomic methods to identify serum biomarkers associated with rat liver toxicity or hypertrophy. *Clinical Chemistry*, 15, 1796-1803.
- Aminudin, N., Chung, Y. S., Eue, S. C., Kang, I. N., & Lee, R. (2007). Blood glucose lowering effect of *Ficus deltoidea* aqueous extract. *Malaysian Journal Of Science*, 26(1), 73-78.
- Anderson, N. L., & Anderson, N. G. (1998). Proteome and proteomics: new technologies, new concepts, and new words. *Electrophoresis*, 19(11), 1853-1861.
- Appel, L. J., Moore, T. J., Obarzanek, E., Vollmer, W. M., Svetkey, L. P., Sacks, F. M., *et al.* (1997). A clinical trial of the effects of dietary patterns on blood pressure: DASH Collaborative Research Group. *New England Journal of Medicine*, 336, 1117-1124.
- Ayub, M. Y., Norazmir, M. N., Mamot, S., Jeeven, K., & Hadijah, H. (2010). Anti-hypertensive effect of pink guava (*Psidium guajava*) puree on spontaneous hypertensive rats. *International Food Research Journal*, 17, 89-96.
- Barley, L. H., & Barley, E. Z. (Eds.). (1976). *Hortus* (3rd ed.): Macmillan General Reference, NY.

- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodology)*, 57, 289-300.
- Blackstock, W. P., & Weir, M. P. (1999). Proteomics: quantitative and physical mapping of cellular proteins. *Trends in Biotechnology*, 17(3), 121-127.
- Boucher, R. A. J., & Genest, J. (1974). A new enzyme leading to the direct formation of Angiotensin II. *Circulation Research*, 34/35, 1202-1212.
- Brickell, C., & Zuk, J. D. (1997). The American Horticultural Society A-Z Encyclopedia of Garden Plants: DK Publishing Inc, New York.
- Buzio, C., Regolisti, G., Perazzoli, F., Mutti, A., Bergamaschi, E., & Borghetti, A. (1994). Renal effects of nifedipine and captopril in patients with essential hypertension and reduced renal reserve. *Hypertension*, 24, 763-769.
- Canby, C. A., Palmer, P. J., Wall, K. R., & Tomanek, R. J. (1989). Effects of captopril on left ventricular structure and function in SHR with established hypertension. *Basic Research in Cardiology*, 84, 306-318.
- Carretero, O. A., & Oparil, S. (2000). Essential Hypertension : Part I: Definition and Etiology. *Circulation*, 101, 329-335.
- Chae, C. U., Lee, R. T., Rifai, N., & Ridker, P. M. (2001). Blood pressure and inflammation in apparently healthy men. *Hypertension*, 38, 399-403.
- Chiellini, C., Santini, F., Marsili, A., Berti, P., Bertacca, A., Polesini, C., et al. (2004). Serum haptoglobin: A novel marker of adiposity in humans. *The Journal of Clinical Endocrinology & Metabolism*, 89(6), 2678-2683.
- Chobanian A.V., Bakris G. L., Black H. R., Cushman W. C., Green L. A., Izzo J. L., Jones D. W., Materson B. J., Oparil S., Wright J. T. Jr, Roccella E. J. (2003) Seventh report of the Joint National Committee on prevention, detection, evaluation and treatment of high blood pressure: the JNC 7 report. *JAMA*, 289, 2560–2570.
- Chobanian, A. V., Bakris G. L., Black H. R., Cushman W. C., Green A. L., Izzo J. L., et al. (2003). Seventh report of the Joint National Committee on prevention, detection, evaluation and treatment of high blood pressure *Hypertension*, 42, 1206-1252.
- Cilia, M., Fish, T., Yang, X., McLaughlin, M., Thannhauser, T. W., & Gray, S. (2009). A comparison of protein extraction methods suitable for gel-based proteomic studies of aphid proteins. *Journal of Biomolecular Techniques*, 20, 201-215.
- Committee, J. N. (1988). The 1988 report of the Joint National Committee on detection, evaluation and treatment of high blood pressure. *Arch Intern Med*, 148, 1023-1038.

- Connell, J. M. C., Whitworth, J. A., Davies, D. L., Lever, A. F., Richards, A. M., & Fraser, R. (1986). Effects of ACTH and cortisol administration on blood pressure, electrolyte metabolism, atrial natriuretic peptide and renal function in normal man. *Journal of Hypertension*, 5, 425–433.
- Conrad, C. H., Brooks, W. W., Hayes, J. A., Sen, S., Robinson, K. G., & Bing, O. H. (1995). Myocardial fibrosis and stiffness with hypertrophy and heart failure in the spontaneously hypertensive rat. *Circulation*, 91, 161-170.
- Cushman, D. W., & Cheung, H. S. (1971). Spectrometric assay and properties of the angiotensin converting enzyme from rat lung. *Biochemical Pharmacology*, 20, 1637-1648.
- Cutler, J. A. (1996). High blood pressure and end-organ damage. *Journal of Hypertension*, 14, S3-S6.
- D'Arniento, J., Dalal, S. S., & Chada, K. (1997). Tissue, temporal and inducible expression pattern of haptoglobin in mice. *Gene*, 195, 19-27.
- Davis, B. J. (1964). Disc electrophoresis. II. Method and application to human serum proteins. *Annals of the New York Academy of Science*, 121, 404-427.
- deKleijin, D. P. V., Smeets, M. B., Kemmeren, P. P. C. W., Lim, S. K., vanMiddelaar, B. J., Velema, E., *et al.* (2002). Acute-phase protein haptoglobin is a cell migration factor involved in arterial restructuring. *The FASEB Journal*, 16, 1123-1125.
- DiBianco R. (1986). Adverse reactions with angiotensin converting enzyme (ACE) inhibitors. *Journal of Medical Toxicology* 1, 122-141.
- Eggertsen, G. H. G., Shiels, B., Reed, D., & Fey, G. H. (1991). Sequence of rat alpha 1 macroglobulin, a broad range proteinase inhibitor from the alpha-macroglobulin-complement family. *Molecular Biology Medicine*, 8, 287-302.
- Elbl, G., & Wagner, H. (1991). A new method for the in vitro screening of inhibitors of angiotensin-converting enzyme (ACE), using the chromophore-labeled and fluorophore labeled substrate, dansyltriglycine. *Planta Medica*, 57(2), 137-141.
- Ely, D. L. (1995.). Organization of cardiovascular and neurohumoral responses to stress: implications for health and disease. *Annals of the New York Academy of Sciences Stress*, 771, 594-608.
- Engström, G., Hedblad B., Janzon L., & Lindgärde F. (2006). Fatality of acute coronary events in relation to hypertension and low-grade inflammation: a population-based cohort study. *Journal of Human Hypertension*, 20, 581-586.
- Engström, G., Janzon, L., Berglund, G., Lind, P., Stavenow, L., Hedblad, B., *et al.* (2002). Blood pressure increase and incidence of hypertension in relation to inflammation-sensitive plasma proteins. *Arteriosclerosis, thrombosis and vascular biology*, 22, 2054-2058.

- Engström, G., Lind, P., Hedblad, B., Stavenow, L., Janzon, L., & Lindgarde, F. (2002). Effect of cholesterol and inflammation- sensitive plasma proteins on incidence of myocardial infarction and stroke in men. *Circulation*, 105, 2632-2637.
- Fazliana, M. S., Shafii, K., Mazleha, M., Mujahir, H., & Hazilawati, H. (2008). Effects of *Ficus deltoidea* aqueous extract on hematological and biochemical parameters in rats. *Medical Journal of Malaysia*, 63, 103-104.
- Filipovsky J, D. P., Eschwege E, Richard JL, Rosselin G, Claude JR. (1996). The relationship of blood pressure with glucose, insulin, heart rate, free fatty acids and plasma cortisol levels according to the degree of obesity in middle-aged men. . *Journal of Hypertension*, 14, 229 –235.
- Folkow, B. (1987). Structure and function of the arteries in hypertension. *American Heart Journal*, 114, 938-948.
- Friedman, G. D., Selby, J. V., & Quesenberry, C. P. (1990). The leukocyte count: a predictor of hypertension. *Journal of Clinical Epidemiology*, 43, 907-911.
- Fukuda, N., Hu, W. Y., Satoh, C., Nakayama, M., Kishioka, H., Kubo, A., et al. (1999). Contribution of synthetic phenotype on the enhanced angiotensin II-generating system in vascular smooth muscle cells from spontaneously hypertensive rats. *Journal of Hypertension*, 17, 1099-1107.
- Fukuda, N., Satoh, C., Hu, W. Y., Soma, M., Kubo, A., Kishioka, H., et al. (1999). Production of angiotensin II by homogenous cultures of vascular smooth muscle cells from spontaneously hypertensive rats. *Arteriosclerosis, thrombosis and vascular biology*, 19, 1210-1217.
- Gettins, P. G. (2002). Serpin structure, mechanism and function. . *Chemical Reviews*, 102(12), 4751-4804.
- Görg, A., Boguth, G., Obermaier, C., Posch, A., & Weiss, W. (1995). Two-dimensional polyacrylamide gel electrophoresis with immobilized pH gradients in the first dimension (IPG-Dalt): the state of the art and the controversy of vertical vs horizontal systems. . *Electrophoresis*, 16, 1079-1086.
- Görg, A., Obermaier, C., Boguth, G., Harder, A., Scheibe, B., Wildgruber, R., et al. (2000). The current state of two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis*, 21, 1037-1053.
- Görg, A., Postel, W., Wese, J., Günther, S., Strahler, J. R., Hanash, S. M., et al. (1987). Elimination of point streaking on silver stained two-dimensional gels by addition of iodoacetamide to the equilibration buffer. *Electrophoresis*, 8, 122-124.
- Görg, A., Postel, W., & Westermeir, R. (1978). Ultrathin-layer isoelectric focusing in polyacrylamide gels on cellophane. . *Analytical Biochemistry*, 89, 60-70.
- Grise, C. B. R., Thibault, G., & Genest, J. (1981). Formation of Angiotensin II by tonin from purified human angiotensinogen. *Canadian Journal of Biochemistry*, 59, 250-255.

- Gros, R., Chorazyczewski, J., Meek, M. D., Benovic, J. L., Ferguson, S. S. G., & Feldman, R. D. (2000). G-Protein–Coupled Receptor Kinase Activity in Hypertension Increased Vascular and Lymphocyte G-Protein Receptor Kinase-2 Protein Expression. *Hypertension*, 35, 38-42.
- Gupta, A. K., Poulter, N. R., Dobson, J., Eldridge, S., Cappuccio, F. P., Caulfield, M., *et al.* (2010). Ethnic differences in blood pressure response to first and second-line antihypertensive therapies in patients randomized in the ASCOT Trial. *American Journal of Hypertension*, 23(9), 1023-1030.
- Hadrava, V., Tremblay, J., & Hamet, P. (1989). Abnormalities in growth characteristics of aortic smooth muscle cells in spontaneously hypertensive rats. *Hypertension*, 13, 589-597.
- Hakiman, M., & Maziah, M. (2009). Non enzymatic and enzymatic antioxidant activities in aqueous extract of *Ficus deltoidea* accessions. *Journal of Medicinal Plant Research*, 3(3), 120-131.
- Hanley, J. M., Haugen, T. H., & Heath, E. C. (1983). Biosynthesis and processing of rat haptoglobin. *Journal of Biological Chemistry*, 258, 7858-7869.
- Haugen, T. H., Hanley, J. M., & Heath, E. C. (1981). Haptoglobin: a novel mode of biosynthesis of a liver secretory glycoprotein. *Journal of Biological Chemistry*, 256, 1055-1057.
- Hoe, S.-Z., Kamaruddin, M. Y., & Lam, S.-K. (2007). Inhibition of angiotensin-converting enzyme activity by a partially purified fraction of *Gynura procumbens* in spontaneously hypertensive rats. *Medical Principles and Practice*, 16, 203-208.
- Høstmark, A. T., Tomten, S. E., & Berg, J. E. (2005). Serum albumin and blood pressure: a population-based, cross-sectional study. *Journal of Hypertension*, 23(4), 725-730.
- Hsu, F. L., Lin, Y. H., Lee, M. H., Lin, C. L., & Hou, W. C. (2002). Both dioscorin, the tuber storage protein of yam (*Dioscorea alata* cv. Tainong No. 1), and its peptic hydrolysates exhibited angiotensin converting enzyme inhibitory activities. *Journal of agricultural and food chemistry*, 21, 6109-6113.
- Hu, W. Y., Fukuda, N., & Kanmatsuse, K. (2002). Growth characteristics, angiotensin II-generation and microarray-determined gene expression in vascular smooth muscle cells from young spontaneously hypertensive rats. *Journal of Hypertension*, 20, 1323-1333.
- Hu, W. Y., Fukuda, N., Satoh, C., Jian, T., Kubo, A., Nakayama, N., *et al.* (2000). Phenotypic modulation by fibronectin enhances angiotensin II-generating system in cultured vascular smooth muscles cells. *Arteriosclerosis, thrombosis and vascular biology*, 20, 1500-1505.
- Hunt, S. C. (Ed.). (2003). *Genetics and family history of hypertension*.

- Hurst, P. L., & Lovell-Smith, C. J. (1981). Optimized assay for serum Angiotensin-Converting Enzyme Activity. *Clinical Chemistry*, 27, 2048-2052.
- Hutchens, T. W., & Yip, T. T. (1992). Synthetic metal-binding protein surface domains for metal ion-dependent interaction chromatography. II. Immobilization of synthetic metal-binding peptides from metal ion transport proteins as model bioactive protein surface domains. . *Journal of Chromatography*, 604, 133-141.
- Hutchens, T. W., & Yip, T. T. (1993). New desorption strategies for the mass spectrometric analysis of macromolecules. *Rapid Communications in Mass Spectrometry*, 7(7), 576-580.
- Ikeda, M., & Arakawa, K. (1984). Kininogenase activity of tonin. *Hypertension*, 6, 222-228.
- Ikeda, M., Sasaguri, M., Maruta, H., & Arakawa, K. (1988). Formation of angiotensin II by tonin-inhibitor complex. *Hypertension*, 11, 63-70.
- Inoue, S., Takamoto, N., Akahori, Y., Masumoto, A., Nakatsukasa, H., Msuyama, H., et al. (2009). Elevated level of serum retinol-binding protein 4 in pregnancy-induced hypertension. *Journal of Obstetrics and Gynaecology Research*, 35, 293-300.
- INTERSALT-Co-operative-research-group. (1988). Sodium, potassium, body mass, alcohol and blood pressure: the INTERSALT study. *Journal of Hypertension*, 6 (supp. 4), S584-S586.
- Issaq, H. J., Conrads, T. P., Prieto, D. A., Tirumalai, R., & D.Veenstra, T. (Eds.). (2003). *SELDI-TOF MS for Diagnostic Proteomics*.
- Iwahashi, M., Muragaki, Y., Ooshima, A., & Umesaki, N. (2002). Decreased type I collagen expression in human uterine cervix during pregnancy. *The Journal of Clinical Endocrinology & Metabolism* 88(5), 2231-2235.
- J Li, Z Zhang, Rosenzweig, J., Wang, Y. Y., & Chan, D. W. (2002). Proteomics and bioinformatics approaches for identification of serum biomarkers to detect breast cancer. *Clinical Chemistry*, 48, 1296-1304.
- James, P. (1997). Protein identification in the post-genome era: the rapid rise of proteomics. *Quartely Reviews of Biophysics*, 30, 279-331.
- Jia, H., Liu, J. W., Ufur, H., He, G. S., Liqian, H., & Chen, P. (2011). The antihypertensive effect of ethyl acetate extract from red raspberry fruit in hypertensive rats. *Pharmacognosy Magazine*, 7, 19-24.
- Jin, L., Chao, L., & Chao, J. (1999). Potassium supplement upregulates the expression of renal kallikrein and bradykinin B 2receptor in SHR. *American Journal of Physiology Renal Physiology*, 276, F476-F484.

- Jr, G. W., Cazares, L. H., Leung, S. M., Nasim, S., Adam, B. L., Yip, T. T., *et al.* (1999). ProteinChip® surface enhanced laser desorption/ionization (SELDI) mass spectrometry: a novel protein biochip technology for detection of prostate cancer biomarkers in complex protein mixtures. *Prostate Cancer and Prostatic Diseases*, 2, 264-276.
- Kalmovarin, N., Friedrichs, W. E., & O'Brien, H. V. (1991). Extrahepatic expression of plasma protein genes during inflammation. *Inflammation*, 15, 369-379.
- Karas, M., Bachmann, D., Bahr, U., & Hillenkamp, F. (1987). Matrix-assisted ultraviolet Laser desorption of non-volatile compounds. *International Journal of Mass Spectrometry and Ion Processes*, 78, 53-68.
- Kearney, P. M., Welthon, M., Reynolds, K., Mutner, P., Whelton, P. K., & He, J. (2005). Global burden of hypertension: analysis of worldwide data. . *Lancet*, 365, 217-223.
- Khan, N., & McAlister, F. A. (2006). Re-examining the efficacy of β-blockers for the treatment of hypertension: a meta-analysis.. *CMAJ*, 174, 1737-1742.
- Kim, D. W., Yokozawa, T., Hattori, M., Kadato, S., & Namba, T. (2000). Effects of aqueous extract extract of *Apocynum venetum* leaves on spontaneously hypertensive, renal hypertensive and NaCl-fed rats. *Journal of Ethnopharmacology*, 72, 53-59.
- Klose, J. (1975). Protein mapping by combined isoelectric focusing and electrophoresis of mouse tissues.A novel approach to testing for induced point mutation in mammals. *Humangenetik*, 26, 231-243.
- Kostis J. B., Wilson, A. C., Freudenberger, R. S., Cosgrove, N. M., Pressel, S. L., & Davis, B. R. (2005). Long-term effect of diuretic-based therapy on fatal outcomes in subjects with isolated systolic hypertension with and without diabetes. *American Journal of Cardiology*, 95, 29-35.
- Kostis, J. B., Davis, B. R., Cutler, J., Richard H. Grimm, J., Berge, K. G., Cohen, J. D., *et al.* (1997). Prevention of heart failure by antihypertensive drug treatment in older persons with isolated systolic hypertension. *JAMA*, 278, 212-216.
- Krakoff, L. R. (2005). Diuretics for Hypertension *Circulation*, 112, e127-e129.
- Krotkiewski, M., Mandroukas, K., Sjöström, L., Sullivan, L., Wetterqvist, H., & Björntorp, P. (1979). Effects of long-term physical training on body fat, metabolism, and blood pressure in obesity. *Metabolism*, 28, 650-658.
- Kushner, & Mackiewicz, A. (1993). The acute phase response: an overview. *Methods in Enzymology*, 163, 373-383.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227, 680-685.

- Leduc, M. S., Shimmin, L. C., Klos, K. L. E., Hanis, C., Boerwinkle, E., & Hixson, J. E. (2008). Comprehensive evaluation of apolipoprotein H gene (APOH) variation identifies novel associations with measures of lipid metabolism in GENOA. *Journal of Lipid Research*, 49, 2648-2656.
- Lee, N. S., Brewer, J. H. B., & Osborne, J. J. C. (1983). B2-glycoprotein 1: Molecular properties of an unusual Apolipoprotein, Apolipoprotein H. *The Journal of Biological Chemistry*, 258, 4765-4770.
- Li J., Zhang Z., Rosenzweig J., Wang Y. Y., Chan D. W. (2002). Proteomics and bioinformatics approaches for identification of serum biomarkers to detect breast cancer. *Clin. Chem.* 48 (8), 1296–304.
- Li, Q. L., Li, B. G., Zhang, Y., Gao, X. P., Li, C. Q., & Zhang, G. L. (2008). Three angiotensin-converting enzyme inhibitors from *Rabdosia coetsa*. Retrieved from <http://www.thefreelibrary.com/Three%20angiotensin-converting%20enzyme%20inhibitors%20from%20Rabdosia%20coetsa-a0180847640>
- Lin, A.-H., Fukuda, N., Jin, X.-Q., Yao, E.-H., Ueno, T., Endo, M., et al. (2004). Complement 3 is involved in the synthetic phenotype and exaggerated growth vascular smooth muscle cells from spontaneously hypertensive rats. *Hypertension*, 44, 42-47.
- Luft, F. C. (1998). Molecular genetics of human hypertension. *Journal of Hypertension*, 16, 1871-1878.
- Maghrani, M., Zeggagh, N.-A., Michel, H.-B., & Eddouks, M. (2005). Antihypertensive effect of *Lepidium sativum* L. in spontaneously hypertensive rats. *Journal of Ethnopharmacology*, 100, 193-197.
- Martiniuk, A. L. C., Lee, C. M. Y., Lawes, C. M. M., Ueshima, H., Suh, I., Lam, T. H., et al. (2007). Hypertension: its prevalence and population-attributable fraction for mortality from cardiovascular disease in the Asia-Pacific region. *Journal of Hypertension*, 25(1), 73-79.
- McEwen, B. S. (Ed.). (1980). *The brain as a target of endocrine hormones*. In *Neuroendocrinology*. (33-42 ed.). Massachusetts: Sinauer Association, Inc.
- McMahon, F. G., & Vargas, R. (1993). Can garlic lower blood pressure? A pilot study. *Pharmacotherapy*, 13(4), 406-407.
- Medzihradsky, K. F., Campbell, J. M., Baldzwin, M. A., Falick, A. M., Juhasz, P., Vestal, M. L., et al. (2000). The characteristics of peptide collision-induced dissociation using a high-performance MALDI-TOF/TOF tandem mass spectrometer. *Analytical Chemistry*, 72(552-558).
- Merrell, K., Southwick, K., Graves, S. W., Esplin, M. S., Lewis, N. E., & Thulin, C. D. (2004). Analysis of low-abundance,low-molecular-weight serum proteins using mass spectrometry. *Journal of Biomolecular Techniques*, 15, 238-248.

- Morganroth, M. L., Schoeneich, S. O., Ward, P. A., Horvath, S. J., & Glovsky, M. M. (1990). C3a55-57, a C-terminal peptide causes thromboxane-dependent pulmonary vascular constriction in isolated perfused rat lungs. *American Review of Respiratory Disease*, 141, 296-300.
- Motoi, H., & Kodama, T. (2003). Isolation and characteristic of Angiotensin II Converting Enzyme inhibitory peptides from wheatgliadin hydrolysate. *Nahrung*, 47, 352-356.
- Muscati, A., Antonelli, S., Bianchi, G., Cavrini, G., Dapporto, S., Ligabue, A., et al. (2007). Serum C3 is a stronger inflammatory marker of insulin resistance than C-reactive protein, leukocyte count and erythrocyte sedimentation rate. A comparison study in an elderly population *Diabetes Care*, 30, 2362-2368.
- Nakagawa, H., & Komorita, N. (1993). Complement component C3-derived neutrophil chemotactic factors purified from exudate of rat carrageenin-induced inflammation. *Biochem. biophys. res. commun*, 194, 1181-1187.
- Nakaya, Y., Schaefer, E. J., & H B Brewer, J. (1980). Activation of human post heparin lipoprotein lipase by apolipoprotein H (beta2-glycoprotein I). *Biochem. biophys. res. commun*, 95, 1168-1172.
- Nava, M., Quiroz, Y., Vaziri, N., & Rodríguez-Iturbe, B. (2003). Melatonin reduces renal interstitial inflammation and improves hypertension in spontaneously hypertensive rats. *American Journal of Physiology Renal Physiology*, 284, F447-454.
- Neuhoff, V., Arold, N., Tauber, D., & Ehrhardt, W. (1985). Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis*, 9, 255-262.
- Nicoletti, A., Mandet, C., Challah, M., Bariety, J., & Michel, J. B. (1996). Mediators of perivascular inflammation in the left ventricle of renovascular hypertensive rats. *Cardiovascular Research*, 31, 585-595.
- Nimpf, J., Bevers, E. M., Bomans, P. H. H., Till, U., Wurm, H., Kostner, G. M., et al. (1986). Prothrombinase activity of human platelets is inhibited by B₂Glycoprotein I. *Biochimica et Biophysica Acta*, 884, 142-149.
- Nogata, Y., Nagamine, T., Yanaka, M., & Ohta, H. (2009). Angiotensin I Converting Enzyme inhibitory peptides produced by autolysis reactions from wheat bran, *Journal of Nutritional Biochemistry*.
- Ocobs D. C., McConkey E. H., & Sammons D. W. (1981). Silver stains for proteins in polyacrylamide gels:a comparison of six methods. *Electrophoresis*, 2, 304-307.
- O'Farrel, P. H. (1975). High Resolution Two-Dimensional Electrophoresis of Proteins. *Journal of Biological Chemistry*, 250(10), 4007-4021.
- Ornstein, L. (1964). Disc electrophoresis. I. Background and theory. *Annals of the New York Academy of Science*, 121, 321-349.

- Papale, M., Pedicillo, M. C., Paolo, S. D., Thatcher, B. J., Muzio, L. L., Bufo, P., *et al.* (2008). Saliva analysis by surface-enhanced laser desorption/ionization time-of-flight mass spectrometry (SELDI-TOF/MS): from sample collection to data analysis. *Clinical Chemistry and Laboratory Medicine*, 46(1), 89-99.
- Pauletto, P., & Rattazzi, M. (2006). Inflammation and hypertension: the search for a link. *Nephrol Dial Transplant*, 21, 850-853.
- Pedroche, J., Yust, M. M., Megías, C., Lqari, H., Manuel Alaiz, Girón-Calle, J., *et al.* (2004). Utilisation of rapeseed protein isolates for production of peptides with angiotensin I-converting enzyme (ACE)-inhibitory activity. *Grasas y Aceites*, 55, 354-358.
- Persson, I. A., Persson, K., Hägg, S., & Andersson, R. G. (2010). Effects of green tea, black tea and Rooibos tea on angiotensin-converting enzyme and nitric oxide in healthy volunteers. *Public Health Nutrition*, 13, 730-737.
- Pesquero, J. L., Thibault, G., & Genest, J. (1982). Effects of substrate size on tonin activity. *Biochemical and Biophysical Research Communications*, 108, 1441-1446.
- Petrella, R. J. (1998). How effective is exercise training for the treatment of hypertension? *Clinical Journal of Sport Medicine*, 8, 224-231.
- Pinto, M. D. S., Kwon, Y.-I., Apostolidis, E., Lajolo, F. M., Genovese, M. I. E., & Shetty, K. (2008). Functionality of bioactive compounds in Brazilian Strawberry (*Fragaria ananassa* Duch.) cultivars: Evaluation of hyperglycemia and hypertension potential using in vitro models. *Journal of agricultural and food chemistry*, 56, 4386-4392.
- Pinto, Y. M., Paul, M., & Ganter, D. (1998). Lessons from rat models of hypertension: from Goldblatt to genetic engineering. *Cardiovascular Research*, 39, 77-88.
- Piubelli, C., Cecconi, D., Astner, H., Caldara, F., Tessari, M., Carboni, L., *et al.* (2005). Proteomic changes in rat serum, polymorphonuclear and mononuclear leukocytes after chronic nicotine administration. *Proteomics*, 5, 1382-1394.
- Polz, E., & Kostner, G. M. (1979). The binding of B₂-glycoprotein-I to human serum serum lipoproteins: distribution among density fractions. *FEBS Letter*, 102(183-186).
- Preuss, H. G., Echard, B., Polansky, M. M., & Anderson, R. (2006). Whole cinnamon and aqueous extracts ameliorate sucrose-induced blood pressure elevations in spontaneously hypertensive rats. *Journal of the American College of Nutrition*, 25, 144-150.
- Pryde P. G., Sedman A. B., Nugent C. E., & Barr M. Jr. (1993). Angiotensin - converting enzyme inhibitor fetopathy. *Journal of the American Society of Nephrology*, 3, 1575 - 1582.

- Quadro, L., Blaner, W. S., Salchow, D. J., Vogel, S., Piantedosi, R., P Gouras, et al. (1999). Impaired retinal function and vitamin A availability in mice lacking retinol-binding protein. *EMBO Journal*, 18, 4633-4644.
- Quadroni, M., & James, P. (1999). Proteomic and automation. *Electrophoresis*, 20, 664-677.
- Raij, L., Dallmasso, A. P., Staley, N. A., & Fish, A. J. (1989). Renal injury in DOCA-salt hypertensive C5-sufficient and deficient mice. *Kidney Int*, 36, 582-592.
- Sanchez, A., & Pettinger, W. A. (1981). Dietary sodium regulation of blood pressure and renal alpha 1 and alpha 2 receptors in WKY and SH rats. *Life Sciences*, 29, 2795-2802.
- Schousboe, I. (1985). B₂Glycoprotein I: A plasma inhibitor of contact activation of the intrinsic blood coagulation pathway. *Blood*, 66, 1086-1091.
- Seibert, V., Wiesner, A., Buschmann, T., & Meuer, J. (2004). Surface-enhanced lased desorption ionization time-of-flight mass spectrometry (SELDI-TOF-MS) and ProteinChip® technology in proteomics research. *Pathology Research and Practice*, 200, 83-94.
- Sen, S., Tarazi, R. C., Khairallah, P. A., & Bumpus, F. M. (1974). Cardiac hypertrophy in spontaneously hypertensive rats. *Circulation Research*, 35, 775-781.
- Sever, P. S., & Poulter, N. R. (1989). A hypothesis for the pathogenesis of essential hypertension: the initiating factors. *Journal of Hypertension*, 7(suppl 1), S9-S12.
- Shapiro, A. L., Vinuela, E., & Maizel, J. V., Jr., . (1967). Molecular weight estimation of polypeptide chains by electrophoresis in SDS-polyacrylamide gels. *Biochemical and Biophysical Research Communications*, 28, 815-820.
- Shevchenko, A., Tomas, H., Havlis, J., Olsen, J. V., & Mann, M. (2007). In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nature Protocols*, 1, 2856-2860.
- Shevchenko, A., Wilm, M., Vorm, A., & Mann, M. (1996). Mass spectrometric analysis of proteins from silver-stained polyacrylamide gels. *Analytical Chemistry*, 68, 850-858.
- Sieber, R., Bütkofer, U., Egger, C., Portmann, R., Walther, B., & Wechsler, D. (2010). ACE-inhibitory activity and ACE-inhibiting peptides in different cheese varieties. *Dairy Science Technology*, 90, 47-73.
- Silagy, C., & Neil, A. (1994). Garlic as a lipid lowering agent — a meta-analysis. *Journal of the Royal College of Physicians of London*, 28(1), 39-45.
- Silagy, C. A., & Neil, H. A. (1994). A meta-analysis of the effect of garlic on blood pressure. *Journal of Hypertension*, 12(4), 463-468.
- Siti-Fatimah-Zahra, M. A., Mahmood, A. A., Hapipah, M. A., Suzita, M. N., & Salmah, I. (2009). Anti-ulcerogenic activity of aqueous extract of *Ficus deltoidea* against

- ethanol-induced gastric mucosal injury in rats. *Research Journal of Medicinal Sciences*, 3(2), 42-46.
- Smeets, M. B., Sluijter, J. P. G., Donners, M. M. P. C., Velema, E., Heeneman, S., Pasterkamp, G., et al. (2003). Increased arterial expression of a glycosylated haptoglobin isoform after balloon dilation. *Cardiovascular Research*, 58, 689-695.
- Smeets, M. B., Sluijter, J. P. G., Donnersc, M. M. P. C., Velemaa, E., Heenemanc, S., Pasterkamp, G., et al. (2003). Increased arterial expression of a glycosylated haptoglobin isoform after balloon dilation. *Cardiovascular Research*, 58, 689-695.
- Solini, A., Santini, E., madec, S., Rossi, C., & Muscelli, E. (2009). Retinol-binding protein-4 in women with untreated essential hypertension. *American journal of Hypertension*, 22, 1001-1006.
- Sottrup-Jensen, L. (1989). Alpha macroglobulins: structure, shape and mechanism of proteinase complex formation. *Journal of Biological Chemistry*, 264, 11539-11542.
- Staessen, J. A., Thijs, L., Fagard, R., O'Brien, E. T., Clement, D., & Leeuw, P. W. d. (1999). Predicting cardiovascular risk using conventional vs ambulatory blood pressure in older patients with systolic hypertension. *JAMA*, 282, 539-546.
- Suetsuna, K., Maekawa, K., & Chen, J.-R. (2004). Antihypertensive effects of *Undaria pinnatifida* (wakame) peptide on blood pressure in spontaneously hypertensive rats. *Journal of Nutritional Biochemistry*, 15, 267-272.
- Sulaiman, M. R., Hussain, M. K., Zakaria, Z. A., Somchmad, M. N., Moin, S., Mohammad, A. S., et al. (2008). Evaluation of the antinociceptive activity of *Ficus deltoidea* aqueous extract. *Fitoterapia*, 79, 557-561.
- Takagi, M., Kimura, K., Atarashi, K., Ikeda, T., Matsuoka, H., Ishii, M., et al. (1990). Immunohistorical study of vascular lesions in severe hypertension induced by DOCA and salt administration to spontaneously hypertensive rats. *American Journal of Hypertension*, 3, 838-845.
- Takebayashi, K., Suetsugu, M., Wakabayashi, S., Aso, Y., & Inukai, T. (2007). Retinol Binding Protein-4 Levels and Clinical Features of Type 2 Diabetes Patients. *The Journal of Clinical Endocrinology & Metabolism*, 92, 2712-2719.
- Tanaka, K., Waki, H., Ido, Y., Akita, S., Yoshida, Y., & Yoshida, T. (1988). protein and polymer analyses up to m/z 100,000 by Laser Ionization Time-of-Flight Mass Spectrometry. *Rapid Commun Mass Spectrom*, 2, 151-153.
- Trotta, R., Donati, M. D., & Iacoviella, L. (2004). Trends in pharmacogenomics of drugs acting on hypertension. *Pharmacology Research*, 49, 351-356.
- Tsuneki, H., Ishizuka, M., Terasawa, M., Wu, J.-B., Sasaoka, T., & Kimura, I. (2004). Effect of green tea on blood glucose levels and serum proteomic patterns in

- diabetic (db/db) mice and on glucose metabolism in healthy humans. *BMC Pharmacology*, 4(18).
- Turner, S. T., Schwartz, G. L., Chapman, A. B., Beitelshes, A. L., Gums, J. G., Cooper-DeHoff, R. M., et al. (2010). Plasma renin activity predicts blood pressure responses to β -blocker and thiazide diuretic as monotherapy and add-on therapy for hypertension. *American Journal of Hypertension*, 23, 1014-1022.
- Vicennati, V., L. Ceroni, L. Gagliardi, et al. . (2002). Response of the hypothalamic-pituitary-adrenocortical axis to high-protein/fat and high carbohydrate meals in women with different obesity phenotypes. *The Journal of Clinical Endocrinology and Metabolism*, 87, 3984-3988.
- Wagner, H., Elbi, G., Lotter, H., & Guinea, M. (1991). Evaluation of natural products as inhibitors of agiotensin I-converting enzyme (ACE). *Pharm. Pharmacol. Letters* 1, 15-18.
- Wagner, H., & Elbl, G. (1992). ACE-inhibitory procyanidins from *Lespedeza-capitata*. *Planta Medica*, 58(3), 297-297.
- Wallerius, S., R. Rosmond, & T. Ljung, e. a. (2003). Rise in morning saliva cortisol is associated with abdominal obesity in men: a preliminary report. *Journal of Endocrinology Investigation*, 26, 616-619.
- Walter, S. V., & Hamet, P. (1989). Enhanced DNA synthesis in heart and kidney of newborn spontaneously hypertensive rats. *Hypertension*, 8, 520-525.
- Warmegard, B. M. N., & Johansson, S. (1992). cDNA cloning and sequencing of rat alpha 1-macroglobulin. *Biochemistry*, 31, 2346-2352.
- Warshafsky, S., Kamer, R. S., & Sivak, S. L. (1993). Effect of garlic on total serum cholesterol. A meta-analysis. *Ann Intern Med*, 119, 599-605.
- Wasinger, V. C., Cordwell, S. J., Cerpa-Poljak, A., Yan, J. X., Gooley, A. A., Wilkins, M. R., et al. (1995). Progress with gene-product mapping of the Mollicutes: *Mycoplasma genitalium*. *Electrophoresis*, 7, 1090-1094.
- Weber, K., & Osborn, M. (1969). The reliability of molecular weight determinations by dodecyl sulphate polyacrylamide gel electrophoresis. *The Journal of Biological Chemistry*, 244, 4406-4412.
- Weijenberg, M. P., Feskens, E. J., Souverijn, J. H., & Kromhout, D. (1997). Serum albumin, coronary heart disease risk, and mortality in an elderly cohort. *Epidemiology*, 8, 87-92.
- Westermeier, R. (2006). Sensitive, quantitative, and fast modifications for coomassie blue staining of polyacrylamide gels. *Proteomics*, 6, 61-64.
- Wilson, C. M. (1979). Staining of protein on gels: comparisons of dyes and procedures. *Methods in Enzymology*, 91, 236-247.

- Wink, K. (2001). Are beta-blockers efficacious as first-line therapy for hypertension in the elderly? *Journal of Clinical and Basic Cardiology*, 4, 235-238.
- Wright, J. T. J., Dunn, J. K., Cutler, J. A., Davis, B. R., Cushman, W. C., Ford, C. E., *et al.* (2005). Outcomes in hypertensive black and nonblack patients treated with chlorthalidone, amlodipine, and lisinopril. *JAMA*, 293, 1595-1608.
- Yates, J. R. (2000). Mass spectrometry: from genomics to proteomics. *Trends in Genetic*, 16, 5-8.
- Ylitalo, K., Pajukanta, P., Meri, S., Cantor, R. M., Mero-Matikainen, N., Vakkilainen, J., *et al.* (2001). Serum C3 but not plasma acylation-stimulating protein is elevated in Finnish patients with familial combined hyperlipidemia. *Journal of the American Heart Association*, 21, 838-843.