

CHAPTER 7

7.0 SUPPLEMENTARY

7.1 BIBLIOGRAPHY

Augsburger, L.L., Vuppala, M.K.(1997) Theory of granulation. In: Parikh, D.M. (Ed.). Handbook of pharmaceutical granulation technology. *Marcel Dekker*: New York 1997.7- 25.

Al-Omranm et al., (2002) M.F. Al-Omranm, S.A. Al-Suwayeh, A.M. El-Helw, S.I. SalehTaste masking of diclofenac sodium using microencapsulation *J. Microencapsul.*, 19 (1) (2002), pp. 45–52.

Beneke, E. C., Viljoen, M. A., & Hamman, H. J., (2009) Polymeric Plant-derived Excipients in Drug Delivery, 14,2602-2620.

Bhardwaj, R. T., Kanwar, M., Lal, R., and Gupta, A. (2000) Natural Gums and Modified Natural Gums as Sustained-Release Carriers. *Drug Development and Industrial Pharmacy*, 26(10);10251038.

Botha, S.A. and Lotter, A.P. (1990), Compatibility study between naproxen and tablet excipients using differential scanning calorimetry. *Drug Dev. Ind. Pharm.*, 16, 673-683.

Clas, S. D., Dalton, C. R., & Hancock, B. C. (1999) Differential scanning Calorimetry: Applications in Drug Development, PSTM Vol. 2.

Coviello, T.; Alhaique, F.; Dorigo, A.; Matricardi, P.; Grassi, M.(2007) Two galactomannans and scleroglucan as matrices for drug delivery: Preparation and release studies. *Eur. J.Pharm.Biopharm.*, 66; 200-209.

Chukwu et al., (1991) A. Chukwu, S.P. Agarwal, M.U. Adikwu Some properties of chloroquine phosphate and quinine hydrochloride microcapsules STP Pharm. Sci., 1 (1991), pp. 117–120.

Faure, A., York, P., Rowe, R.C. Process control and scale-up of pharmaceutical wet granulation processes: a review. *Eur J Pharm Biopharm* 52 (2001) 269 - 277.

Ford, J.L. (1993), The use of thermal analysis in preformulation compatibility testing, Acta of Symposium on Pharmacy and Thermal Analysis, Freiburg, p. 2.

Ford,J.L. and Timmins, P.(1989) in: M.H. Rubinstein (Ed.), Pharmaceutical Thermal Analysis, Ellis Horwood, Chichester , pp. 201–237.

Gabaude, C.M.D., Gautier, J.C., Saudemon, P. Validation of a new pertinent packing coefficient to estimate flow properties of pharmaceutical powders at a very early development stage, by comparison with mercury intrusion and classical flowability methods. *J Mater Sci* 36 (2001) 1763 - 1773.

Giron, D. (1986) Applications of thermal analysis in the pharmaceutical industry, *J. Pharm. Biomed. Anal.*, 4, 755– 770.

Guerin, E., Tchoreloff, P., Leclerc, B., Tanguy, D., Deleuil, M., Couarrazé, G.,Rheological characterization of pharmaceutical powders using tap testing, shear celland mercury porosimeter. *Int J Pharm* 189 (1999) 91 - 103.

Gu, L., Liew, C.V.,&Heng, P.W. (2004) Wet spheronization by rotary processing: amultistage single-pot process for producing spheroids. *Drug Dev Ind Pharm.*, 30: 111123.

Gupta, V.K., Hariharan, M., Weatley, T.A.,& Price, J.C. (2001) Controlledrelease tablets from carrageenans: effect of formulation, storage and dissolution factors. *Eur J Pharm Biopharm.* ;51:241–8.

Gupta, V.K.; Hariharan, M.; & Wheatley, T.A., & Price, J.C.(2001) Controlled-release tablets from carrageenans: effect of formulation, storage and dissolution factors. *Eur. J.Pharm. Biopharm.*, 51, 241-248.

Hashimoto et al., (2002) Y. Hashimoto, M. Tanaka, H. Kishimoto, H. Shizowa, K. Hasegawa, K. Matsuyama, T. Uchida Preparation, characterization and taste-masking properties of polyvinylacetate microspheres containing trimebutine *J. Pharm. Pharmacol.*, 54 (2002), pp. 1323–1328.

Kristensen, H.G., Schaefer, T. Granulation - a review of pharmaceutical wet-granulation. *Drug Dev Ind Pharm* 13 (1987) 803 - 872.

Leuenberger, H., Luy, B., & Struder, J. (1990) New development in the control of a moist agglomerationand pelletization process. *STP Pharma Sci.*, 6: 303-309.

Lin, S.Y., & Han, R.Y. (1992). *Pharmazie*, 47, 266-268.

Matsuda et al., 2001S Matsuda, H Hatano, K Kuramoto, A Tsutsumi Fluidization of ultrafine particles with high G *Journal of Chemical Engineering of Japan*, 34 (2001), pp. 121–125.

Mohamadnia, Z.; Zohuriaan-Mehr, M.J.; Kabiri, K.; Jamshidi, A.; &Mobedi, H.(2008) Ionically crosslinked carrageenan-alginate hydrogel beads. *J. Biomater. Sci. Polymer Edn.*, 19, 47-59.

Mura, P., Manderioli, G., Bramanti, G., Furlanetto, S.,& Pinzauti, S. (1994). Utilization of differential scanning calorimetry as a screening technique to determine the compatibility of ketoprofen with excipients, 119, 71-79.

Nerurkar, J.; Jun, H.W.; Price, J.C.; Park, M.O.(2005) Controlled-release matrix tablets of ibuprofen using cellulose ethers and carrageenans: effect of formulation factors on dissolution rates. *Eur. J.Pharm.Biopharm.*, 61, 56-68.

Picker, K.M. (1999) Matrix tablets of carrageenans. I. A compaction study. *Drug Dev. Ind. Pharm.*, 25, 329-337.

Robson et al., (1999) H.J. Robson, D.Q.M. Craig, D. Deutsch An investigation into the release of cefuroxime axetil from taste-masked stearic acid microspheres. Part 1. The influence of the dissolution medium on the drug release profile and the physical integrity of the microspheres *Int. J. Pharm.*, 190 (1999), pp. 183–192.

Rumpf, H. Grundlagen und Methoden des Granulierens. *Chem Ing Tech* 3 (1958)144 - 158.

Saurabh, S., & Garima, M. (2010). Fluid Bed Technology: Overview and Parameters for Process Selection. *International Journal of Pharmaceutical sciences and drug Research*, 2, 236-246.

Shimano et al., (1995) K. Shimano, O. Kondo, A. Miwa, Y. Higashi, S. Go to Evaluation of uniform-sized microcapsules using a vibration-nozzle method *Drug Dev. Ind. Pharm.*, 21 (1995), pp. 331–347

Sorenti (2010) Thermal characterization of diltiazem and carrageenan binary systems. *J. Therm. Anal. Calorim.*, 102; 337-342.

Tapia, C., Corbala'n V., Costa, E., Nella Gai M., Yazdani-Pedram M.(2005) Study of the release mechanism of diltiazem hydrochloride from matrices based on chitosan–alginate and chitosan–carrageenan mixtures. *Biomacromolecules*. 6:2389–95.

Turi, E.A., ed. (1997) Thermal Characterization of Polymeric Materials (2nd edn), *Academic Press*, San Diego, CA, USA

Vuppala, M. K., Parikh, D. M., & Bhagat, H. R. (1997) Application of powder layering technology and film coating for manufacture of sustained release pellets using a rotary fluid bed processor. *Drug Dev Ind Pharm.*, 23, 687-694.

Wilson et al., (1994) M.L. Wilson, W.A. Ritshel, A. Sakr Development of an ibuprofen lozenge for the elderly Pharm. Ind., 56 (1994), pp. 836–841.

Yajima et al., (1996) T. Yajima, A. Nogata, M. Demachi, N. Umeki, S. Itai, N. Yunoki, M. Remoto Particle design for taste-masking using a spray-congealing technique *Chem. Pharm. Bull.*, 44 (1996), pp. 187–191.

Ylirussi, J., Rasanen, E., Rantanen, J., &Mannermaa, J.P. (2004) The characterization of Fluidization Bahaviour Using a Novel Multichamber Microscale Fluid Bed. *J. PharmaSci.*, 3, 780-791.