

CHAPTER 7

REFERENCES

- Adhikari, K. N. and Campbell, C. G. (1998). *In vitro* germination and viability of buckwheat (*Fagopyrum esculentum* Moench) pollen. *Euphytica*, 102(1), 87–92.
- Agrawal, N., Dasaradhi, P. V. N., Mohmmmed, A., Malhotra, P., Bhatnagar, R. K. and Mukherjee, S. K. (2003). RNA interference: biology, mechanism, and applications. *Microbiology and Molecular Biology Reviews*, 67(4), 657.
- Akashi, H., Miyagishi, M. and Taira, K. (2001). Suppression of gene expression by RNA interference in cultured plant cells. *Antisense and Nucleic Acid Drug Development*, 11(6), 359–367.
- Al-Babili, S. and Beyer, P. (2005). Golden Rice-five years on the road-five years to go? *Trends in Plant Science*, 10(12), 565–573.
- Alonso, J. M. and Ecker, J. R. (2006). Moving forward in reverse: genetic technologies to enable genome-wide phenomic screens in *Arabidopsis*. *Nature Reviews Genetics*, 7(7), 524–536.
- Amiss, T. and Prenell, S. C. (2006). Nucleic Acid Blotting Techniques: Theory and Practice. In W. B. Coleman and G. J. Tsongalis (Eds.), *Molecular diagnostics: for the clinical laboratorian* (pp. 567): Humana Press.
- An, G., Jeong, D. H., Jung, K. H. and Lee, S. (2005). Reverse genetic approaches for functional genomics of rice. *Plant Molecular Biology*, 59(1), 111–123.
- Armstrong, T. T., Fitzjohn, R. G., Newstrom, L. E., Wilton, A. D. and Lee, W. G. (2005). Transgene escape: what potential for crop-wild hybridization? *Molecular Ecology*, 14(7), 2111–2132.

- Ashikari, M. and Matsuoka, M. (2002). Application of rice genomics to plant biology and breeding. *Botanical Bulletin of Academia Sinica*, 43, 1–11.
- Ashikari, M., Sakakibara, H., Lin, S., Yamamoto, T., Takashi, T., Nishimura, A., *et al.* (2005). Cytokinin oxidase regulates rice grain production. *Science*, 309(5735), 741.
- Asins, M. J. (2002). Present and future of quantitative trait locus analysis in plant breeding. *Plant Breeding*, 121(4), 281–291.
- Bagasra, O. and Prilliman, K. R. (2004). RNA interference: the molecular immune system. *Journal of Molecular Histology*, 35(6), 545–553.
- Bailey, J. M. and Whelan, W. J. (1961). Physical properties of starch. *Journal of Biological Chemistry*, 236(4), 969.
- Baulcombe, D. (2000). Unwinding RNA silencing. *Science*, 290(5), 1108–1109.
- Baorong, L., Zhiping, S. and Jiakuan, C. (2003). Can transgenic rice cause ecological risks through transgene escape? *Progress in Natural Science*, 13(1), 17–24.
- Bartel, B. and Bartel, D. P. (2003). MicroRNAs: at the root of plant development? *Plant Physiology*, 132(2), 709.
- Baskin, C. C. and Baskin, J. M. (2001). *Seeds: ecology, biogeography, and evolution of dormancy and germination*: Academic Press.
- Baulcombe, D. (2004). RNA silencing in plants. *Nature*, 431(7006), 356–363.
- Becraft, P. W. (2002). Receptor kinase signaling in plant development. *Annual Review of Cell and Developmental Biology*, 18(6), 163–192.
- Bengtsson, S. (2006). *Evaluation of transgenic *Campanula carpatica* plants*. The Royal Veterinary and Agricultural University, Copenhagen.

- Bernacchi, D., Beck-Bunn, T., Eshed, Y., Lopez, J., Petiard, V., Uhlig, J., *et al.* (1998). Advanced backcross QTL analysis in tomato. I. Identification of QTLs for traits of agronomic importance from *Lycopersicon hirsutum*. *TAG Theoretical and Applied Genetics*, 97(3), 381–397.
- Bernstein, E., Caudy, A., Hammond, S. M. and Hannon, G. J. (2001). Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature*, 409(6818), 295–296.
- Bevan, M. (2002). Genomics and plant cells: application of genomics strategies to Arabidopsis cell biology. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 357(1422), 731.
- Bhat, S. R. and Srinivasan, S. (2002). Molecular and genetic analyses of transgenic plants: Considerations and approaches. *Plant Science*, 163(4), 673–681.
- Blair, M. W., Iriarte, G. and Beebe, S. (2006). QTL analysis of yield traits in an advanced backcross population derived from a cultivated Andean × wild common bean (*Phaseolus vulgaris* L.) cross. *TAG Theoretical and Applied Genetics*, 112(6), 1149–1163.
- Bots, M. and Mariani, C. (2005). *Pollen viability in the field*: Radboud Universiteit Nijmegen.
- Bouchez, D. and Hofte, H. (1998). Functional genomics in plants. *Plant Physiology*, 118(3), 725.
- Brand, U., Fletcher, J. C., Hobe, M., Meyerowitz, E. M. and Simon, R. (2000). Dependence of stem cell fate in Arabidopsis on a feedback loop regulated by CLV3 activity. *Science's STKE*, 289(5479), 617.
- Brar, D. S. and Khush, G. S. (2006). Cytogenetic manipulation and germplasm enhancement of rice (*Oryza sativa* L.). In R. J. Singhand P. Jauhar (Eds.), *Genetic resources, chromosome engineering, and crop improvement* (Vol. 2, pp. 115–158): CRC Press.

- Braun, D. M., Stone, J. M. and Walker, J. C. (1997). Interaction of the maize and Arabidopsis kinase interaction domains with a subset of receptor-like protein kinases: implications for transmembrane signaling in plants. *The Plant Journal*, 12(1), 83–95.
- Brondani, C., Rangel, P., Brondani, R. and Ferreira, M. (2002). QTL mapping and introgression of yield-related traits from *Oryza glumaepatula* to cultivated rice (*Oryza sativa*) using microsatellite markers. *TAG Theoretical and Applied Genetics*, 104(6), 1192–1203.
- Burlibaşa, M., Burlibaşa, L., Gavrilă, L. B., Gavrilă, V. R. and Gavrilă, L. (2008). MicroRNA a macro Revolution in Medical Biotechnologies. *Roumanian Biotechnological Letters*, 13(6), 3977-3983.
- Carthew, R. W. and Sontheimer, E. J. (2009). Origins and mechanisms of miRNAs and siRNAs. *Cell*, 136(4), 642–655.
- Cazonelli, C. I. and Velten, J. (2004). Analysis of RNA-mediated gene silencing using a new vector (pKNOCKOUT) and an in planta *Agrobacterium* transient expression system. *Plant molecular biology reporter*, 22(4), 347–359.
- Chapman, E. J. and Carrington, J. C. (2007). Specialization and evolution of endogenous small RNA pathways. *Nat Rev Genet*, 8, 884–896.
- Chapman, M. A. and Burke, J. M. (2006). Letting the gene out of the bottle: the population genetics of genetically modified crops. *New Phytologist*, 170(3), 429–443.
- Clark, S. E., Running, M. P. and Meyerowitz, E. M. (1993). CLAVATA1, a regulator of meristem and flower development in Arabidopsis. *Development*, 119(2), 397.
- Clark, S. E., Williams, R. W. and Meyerowitz, E. M. (1997). The CLAVATA1 gene encodes a putative receptor kinase that controls shoot and floral meristem size in Arabidopsis. *Cell*, 89(4), 575–585.

- Collard, B. C. Y., Jahufer, M. Z. Z., Brouwer, J. B. and Pang, E. C. K. (2005). An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: The basic concepts. *Euphytica*, 142(1), 169–196.
- Dawe, D. (2002). Golden rice: what role could it play in alleviation of vitamin A deficiency? *Food Policy*, 27(5-6), 541–560.
- Dillon, C. P., Sandy, P., Nencioni, A., Kissler, S., Rubinson, D. A. and Van Parijs, L. (2005). RNAi as an experimental and therapeutic tool to study and regulate physiological and disease processes. *Physiology*, 67(1), 147.
- Doebley, J., Stec, A. and Hubbard, L. (1997). The evolution of apical dominance in maize.
- Eastham, K. and Sweet, J. (2002). Genetically modified organism (GMOs): The significance of gene flow through pollen transfer. *Environmental Issue Report*, 52(8).
- Erlandson, M. A. and Theilmann, D. A. (2009). Molecular approaches to viruse characterization and detection. In S. P. Stock, I. Glazer, N. Boemare and J. Vandenberg (Eds.), *Insect Pathogens: Molecular Approaches and Techniques* (pp. 440).
- Falconer, D. (1965). *The inheritance of liability to certain diseases, estimated from the incidence among relatives* (Vol. 29).
- Fan, C., Xing, Y., Mao, H., Lu, T., Han, B., Xu, C., *et al.* (2006). GS3, a major QTL for grain length and weight and minor QTL for grain width and thickness in rice, encodes a putative transmembrane protein. *TAG Theoretical and Applied Genetics*, 112(6), 1164–1171.
- Finch-Savage, W. E. and Leubner-Metzger, G. (2006). Seed dormancy and the control of germination. *New Phytologist*, 171(3), 501–523.

- Fletcher, J. C., Brand, U., Running, M. P., Simon, R. and Meyerowitz, E. M. (1999). Signaling of cell fate decisions by CLAVATA3 in Arabidopsis shoot meristems. *Science*, 283(5409), 1911.
- Friedrichsen, D. M., Joazeiro, C. A. P., Li, J., Hunter, T. and Chory, J. (2000). Brassinosteroid-insensitive-1 is a ubiquitously expressed leucine-rich repeat receptor serine/threonine kinase. *Plant Physiology*, 123(4), 1247.
- Fu, Q., Zhang, P., Tan, L., Zhu, Z., Ma, D., Fu, Y., *et al.* (2010). Analysis of QTLs for yield-related traits in Yuanjiang common wild rice (*Oryza rufipogon* Griff.). *Journal of Genetics and Genomics*, 37(2), 147–157.
- Futuyma, D. J. (1998). *Evolutionary Biology* (Third ed.): Sinauer Associates.
- Ge, S., Sang, T., Lu, B. R. and Hong, D. Y. (1999). Phylogeny of rice genomes with emphasis on origins of allotetraploid species. *Proceedings of the National Academy of Sciences of the United States of America*, 96(25), 14400.
- German, M. A., Kandel-Kfir, M., Swarzberg, D., Matsevitz, T. and Granot, D. (2003). A rapid method for the analysis of zygosity in transgenic plants. *Plant Science*, 164(2), 183-187.
- Gigerenzer, G., Swijtink, Z. and Daston, L. (1990). *The empire of chance: How probability changed science and everyday life*: Cambridge University Press.
- Gilissen, L. J. W., Metz, P. L. J., Stiekema, W. J. and Nap, J. P. (1998). Review: Biosafety of *E. coli* -glucuronidase (GUS) in plants. *Transgenic Research*, 7(3), 157–163.
- Gregory, R. I., Chendrimada, T. P., Cooch, N. and Shiekhattar, R. (2005). Human RISC couples microRNA biogenesis and posttranscriptional gene silencing. *Cell*, 123(4), 631–640.

- Groot, M. H. M., van de Wiel, C. C. M., van Tienderen, P. H. and den Nijs, H. C. M. (2003). Hybridisation and introgression between crops and wild relatives. *Cogem Research*, 2.
- Gupta, P. K. (2002). Molecular markers and QTL analysis in crop plants. *Current Science*, 83(2), 113–114.
- Gupta, P. K. (2007). *Genetics Classical To Modern*: Rastogi Publications.
- Gur, A. and Zamir, D. (2004). Unused natural variation can lift yield barriers in plant breeding. *PLoS Biology*, 2, 1610–1615.
- Hails, R. S. and Morley, K. (2005). Genes invading new populations: a risk assessment perspective. *Trends in Ecology & Evolution*, 20(5), 245–252.
- Hames, B. D. (1995). Analysis of gene expression by blotting techniques. In S. J. Higgins (Ed.), *Gene Probes 2: A Practical Approach* (pp. 448): Oxford University Press.
- Hanafi, M. M., Hartinie, A., Shukor, J. and Mahmud, T. M. M. (2009). Upland Rice Varieties in Malaysia: Agronomic and Soil Physico-Chemical Characteristics. *Tropical Agricultural Science*, 32(2).
- Hannon, G. J. (2002). RNA interference. *Nature*, 418(6894), 244–251.
- Haygood, R., Ives, A. R. and Andow, D. A. (2004). Population genetics of transgene containment. *Ecology Letters*, 7(3), 213–220.
- Hey-Chi, H., Shian-Yang, P. and Chia-Tung, S. (1991). High quality of DNA retrieved for Southern blot hybridization from microwave-fixed, paraffin-embedded liver tissues. *Virological Methods*, 31(2-3), 251–261.
- Hieter, P. and Boguski, M. (1997). Functional genomics: it's all how you read it. *Science*, 278(5338), 601.

- Honda, M., Muramoto, Y., Kuzuguchi, T., Sawano, S., Machida, M. and Koyama, H. (2002). Determination of gene copy number and genotype of transgenic *Arabidopsis thaliana* by competitive PCR. *Journal of experimental botany*, 53(373), 1515.
- Initiative, A. G. (2000). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature*, 408(6814), 796–815.
- Jeong, S., Trotochaud, A. E. and Clark, S. E. (1999). The *Arabidopsis* CLAVATA2 gene encodes a receptor-like protein required for the stability of the CLAVATA1 receptor-like kinase. *The Plant Cell*, 11(10), 1925.
- Jungkind, D. and Kessler, H. H. (2002). Molecular Methods for Diagnosis of Infectious Disease. In A. L. Truant (Ed.), *Manual of Commercial Methods in Clinical Microbiology* (pp. 481): ASM Press.
- Kang, J. X. and Leaf, A. (2007). Why the omega-3 piggy should go to market. *Nature Biotechnology*, 25(5), 505–506.
- Kayes, J. M. and Clark, S. E. (1998). CLAVATA2, a regulator of meristem and organ development in *Arabidopsis*. *DEVELOPMENT-CAMBRIDGE-*, 125, 3843–3851.
- Khatun, S. and Flowers, T. J. (1995). The estimation of pollen viability in rice. *Journal of Experimental Botany*, 46(1), 151.
- Khush, G. S. (1997). Origin, dispersal, cultivation and variation of rice. *Plant Molecular Biology*, 35(1), 25–34.
- Kiani, G., Nematzadeh, G. A., Ghareyazie, B. and Sattari, M. (2009). Genetic analysis of cry1Ab gene in segregating populations of rice. *African Journal of Biotechnology*, 8(16), 3703–3707.

- Kihara, T., Zhao, C. R., Kobayashi, Y., Takita, E., Kawazu, T. and Koyama, H. (2006). Simple identification of transgenic Arabidopsis plants carrying a single copy of the integrated gene. *Bioscience, biotechnology, and biochemistry*, 70(7), 1780–1783.
- Kirk, R. E. (2007). *Statistics: an introduction*: Wadsworth Pub Co.
- Kusaba, M. (2004). RNA interference in crop plants. *Current Opinion in Biotechnology*, 15(2), 139–143.
- Kuvshinov, V., Koivu, K., Kanerva, A. and Pehu, E. (2001). Molecular control of transgene escape from genetically modified plants. *Plant Science*, 160(3), 517–522.
- Lander, E. S. and Botstein, D. (1989). Mapping Mendelian factors underlying quantitative traits using RFLP linkage maps. *Genetics*, 121(1), 185.
- Law, Y. S. (2011). *Molecular characterization and expression analysis of putative RPK1 and putative CLV1 in Malaysian indica rice (Oryza Sativa L.) CV. MR219 and Malaysian wild rice (Oryza rufipogon) IRGC105491*. University of Malaya, Kuala Lumpur.
- Lee, H.-S., Karunanandaa, B., McCubbin, A., Gilroy, S. and Kao, T.-h. (1996). PRK1, a receptor-like kinase of *Petunia inflata*, is essential for postmeiotic development of pollen *The Plant Journal*, 9(5), 613–624.
- Lehti-Shiu, M. D., Zou, C., Hanada, K. and Shiu, S. H. (2009). Evolutionary history and stress regulation of plant receptor-like kinase/pelle genes. *Plant Physiology*, 150(1), 12.
- Leister, D. (2005). *Plant Functional Genomics*: Food Products Press.
- Li, J. and Chory, J. (1997). A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. *Cell*, 90(5), 929–938.

- Lu, B. R. (1999). Taxonomy of the genus *Oryza* (Poaceae): historical perspective and current status. *International Rice Research Notes*, 24(3).
- Lu, B. R. (2008). Transgene escape from GM crops and potential biosafety consequences: an environmental perspective. *Collection of Biosafety Reviews*, 4, 66–141.
- Marri, P., Sarla, N., Reddy, L. and Siddiq, E. A. (2005). Identification and mapping of yield and yield related QTLs from an Indian accession of *Oryza rufipogon*. *BMC genetics*, 6(1), 33.
- Matsushima, S., Wada, G. and Matsuzaki, A. (1966). Analysis of yield-determining process and its application to yield-prediction and culture improvement of lowland rice. *Crop Science Society of Japan*, 34, 321–328.
- Matthew, L. (2004). RNAi for plant functional genomics. *Comparative and Functional Genomics*, 5(3), 240–244.
- Matzke, M., Matzke, A., Pruss, G. and Vance, V. (2001). RNA-based silencing strategies in plants. *Current Opinion in Genetics & Development*, 11, 221–227.
- McCouch, S. R., Sweeney, M., Li, J., Jiang, H., Thomson, M., Septiningsih, E., *et al.* (2007). Through the genetic bottleneck: *O. rufipogon* as a source of trait-enhancing alleles for *O. sativa*. *Euphytica*, 154(3), 317–339.
- McCough, S. R. and Doerge, R. W. (1995). QTL mapping in rice. *TRENDS in Genetics*, 11(12), 482–487.
- Meister, G. and Tuschl, T. (2004). Mechanisms of gene silencing by double-stranded RNA. *Nature*, 431, 343–349.
- Messeguer, J., Peñas, G., Ballester, J., Bas, M., Serra, J., Salvia, J., *et al.* (2006). Pollen-mediated gene flow in maize in real situations of coexistence. *Plant Biotechnol Journal*, 4, 633–645.

- Miki, D. and Shimamoto, K. (2004). Simple RNAi vectors for stable and transient suppression of gene function in rice. *Plant and cell physiology*, 45(4), 490.
- Miura, K., Ikeda, M., Matsubara, A., Song, X. J., Ito, M., Asano, K., *et al.* (2010). OsSPL14 promotes panicle branching and higher grain productivity in rice. *Nature genetics*, 42(6), 545–549.
- Moncada, P., Martinez, C. P., Borrero, J., Chatel, M., Gauch, J. H., Guimaraes, E., *et al.* (2001). Quantitative trait loci for yield and yield components in an *Oryza sativa* × *Oryza rufipogon* BC2F2 population evaluated in an upland environment. *TAG Theoretical and Applied Genetics*, 102(1), 41–52.
- Moncada, P., Martinez, C. P., Borrero, J., Chatel, M., Gauch Jr, H., Guimaraes, E., *et al.* (2001). Quantitative trait loci for yield and yield components in an *Oryza sativa* × *Oryza rufipogon* BC2F2 population evaluated in an upland environment. *TAG Theoretical and Applied Genetics*, 102(1), 41–52.
- Moore, G., Devos, K. M., Wang, Z. and Gale, M. D. (1995). Cereal genome evolution: grasses, line up and form a circle. *Current Biology*, 5(7), 737–739.
- Mu, J. H., Lee, H. S. and Kao, T. H. (1994). Characterization of a Pollen-Expressed Receptor-like Kinase Gene of *Petunia inflata* and the Activity of Its Encoded Kinase. *The Plant Cell*, 6, 709–721.
- Murray, M. G. and Thompson, W. F. (1980). Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Research*, 8(19), 4321.
- Napoli, C., Lemieux, C. and Jorgensen, R. (1990). Introduction of a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes in trans. *The Plant Cell*, 2(4), 279.
- Ohnishi, T., Yoshino, M., Yamakawa, H. and Kinoshita, T. (2011). The biotron breeding system: a rapid and reliable procedure for genetics studies and breeding in rice.

- Ori, N., Eshed, Y., Paran, I., Presting, G., Aviv, D., Tanksley, S., *et al.* (1997). The I2C family from the wilt disease resistance locus I2 belongs to the nucleotide binding, leucine-rich repeat superfamily of plant resistance genes. *The Plant Cell*, 9(4), 521. *Plant and Cell Physiology*, 52(7).
- Parker, G. S., Eckert, D. M. and Bass, B. L. (2006). RDE-4 preferentially binds long dsRNA and its dimerization is necessary for cleavage of dsRNA to siRNA. *RNA*, 12(5), 807–818.
- Pasquinelli, A. E., Hunter, S. and Bracht, J. (2005). MicroRNAs: a developing story. *Current Opinion in Genetics & Development*, 15(2), 200–205.
- Paterson, A. H., Lander, E. S., Hewitt, J. D., Peterson, S., Lincoln, S. E. and Tanksley, S. D. (1989). Resolution of quantitative traits into Mendelian factors by using a complete linkage map of restriction fragment length polymorphisms *Nature*, 335, 721–726.
- Paterson, A. H. and Press, C. R. C. (1998). *Molecular dissection of complex traits*: CRC Press.
- Peters, J. L., Cnudde, F. and Gerats, T. (2003). Forward genetics and map-based cloning approaches. *Trends in Plant Science*, 8(10), 484–491.
- Pinheiro, P. V., Faria, J. C., Nogueira, E. O. P. L. and Aragão, F. J. L. (2009). Transgene inheritances and genetic similarities of near isogenic lines of genetically modified common beans. *Pesquisa Agropecuária Brasileira*, 44(9), 1168–1176.
- Price, A. H. (2006). Believe it or not, QTLs are accurate! *Trends in Plant Science*, 11(5), 213–216.
- Puchta, H. and Hohn, B. (1996). From centiMorgans to base pairs: homologous recombination in plants. *Trends in Plant Science*, 1(10), 340–348.

- Robles, P., Quesada, V. and Hricová, A. (2010). Unravelling gene function through mutagenesis *Molecular Techniques in Crop Improvement*: Springer Verlag.
- Rodriguez-Riano, T. and Dafni, A. (2000). A new procedure to asses pollen viability. *Sexual Plant Reproduction*, 12(4), 241–244.
- Sakamoto, T. (2006). Phytohormones and rice crop yield: strategies and opportunities for genetic improvement. *Transgenic research*, 15(4), 399-404.
- Sakamoto, T. and Matsuoka, M. (2008). Identifying and exploiting grain yield genes in rice. *Current opinion in plant biology*, 11(2), 209–214.
- Sambrook, J., Maniatis, T. and Fritsch, E. F. (1982). *Molecular cloning: A laboratory manual* (Vol. 1): Cold Spring Harbor Laboratory.
- Sambrook, J. and Russell, D. W. (2001). *Molecular cloning: a laboratory manual* (Vol. 2): CSHL Press.
- Sang, T. and Ge, S. (2007). The puzzle of rice domestication. *Journal of Integrative Plant Biology*, 49(6), 760–768.
- Sato, F. (2005). RNAi and functional genomics. *Plant Biotechnology*, 22(5), 431–442.
- Sax, K. (1923). The association of size differences with seed-coat pattern and pigmentation in *Phaseolus vulgaris*. *Genetics*, 8(6), 552.
- Scattat, M., Klosgen, R. and Marques, J. (2004). A novel vector for efficient gene silencing in plants. *Plant Mol Biol Rep* 22, 145–153.
- Septiningsih, E. M., Prasetiyono, J., Lubis, E., Tai, T. H., Tjubaryat, T., Moeljopawiro, S., *et al.* (2003). Identification of quantitative trait loci for yield and yield components in an advanced backcross population derived from the *Oryza sativa* variety IR64 and the wild relative *O. rufipogon*. *TAG Theoretical and Applied Genetics*, 107(8), 1419–1432.

- Septiningsih, E. M., Trijatmiko, K. R., Moeljopawiro, S. and McCouch, S. R. (2003). Identification of quantitative trait loci for grain quality in an advanced backcross population derived from the *Oryza sativa* variety IR64 and the wild relative *O. rufipogon*. *TAG Theoretical and Applied Genetics*, 107(8), 1433–1441.
- Sharma, S. D., Tripathy, S. and Biswal, J. (2000). Origin of *O. sativa* and its ecotypes. In J. S. Nanda (Ed.), *Rice breeding and genetics: research priorities and challenges* (pp. 349–369): Science Publications.
- Shiu, S. H. and Bleecker, A. B. (2001). Plant receptor-like kinase gene family: diversity, function, and signaling. *Science's STKE*, 2001(113).
- Shiu, S. H. and Bleecker, A. B. (2003). Expansion of the receptor-like kinase/Pelle gene family and receptor-like proteins in *Arabidopsis*. *Plant Physiology*, 132(2), 530.
- Shiu, S. H., Karlowski, W. M., Pan, R., Tzeng, Y. H., Mayer, K. F. X. and Li, W. H. (2004). Comparative analysis of the receptor-like kinase family in *Arabidopsis* and rice. *The Plant Cell*, 16(5), 1220.
- Shivanna, K. R., Linskens, H. F. and Cresti, M. (1991). Pollen viability and pollen vigor. *Theoretical and Applied Genetics*, 81(1), 38–42.
- Smith-Zagone, M. J., Pulliam, J. F. and Farkas, D. H. (2007). Molecular Pathology Methods. In D. G. B. Leonard (Ed.), *Molecular Pathology in Clinical Practice* (pp. 600): Springer.
- Smith, N. A., Singh, S. P., Wang, M. B., Stoutjesdijk, P. A., Green, A. G. and Waterhouse, P. M. (2000). Gene expression: Total silencing by intron-spliced hairpin RNAs. *Nature*, 407(6802), 319–320.
- Snow, A. A., Andow, D. A., Gepts, P., Hallerman, E. M., Power, A., Tiedje, J. M., *et al.* (2005). Genetically engineered organisms and the environment: current status and recommendations. *Ecological Applications*, 15(2), 377–404.

- Snowdon, R. J. and Langsdorf, A. (1998). An Introduction to DNA Fingerprinting using RLFP and RAPD Techniques. In D. Tietz (Ed.), *Nucleic acid electrophoresis* (pp. 328): Springer.
- Sofi, P. and Rather, A. G. (2007). QTL analysis in rice improvement: concept, methodology and application. *Biotechnology*, 6(1), 1–13.
- Song, X. J., Huang, W., Shi, M., Zhu, M. Z. and Lin, H. X. (2007). A QTL for rice grain width and weight encodes a previously unknown RING-type E3 ubiquitin ligase. *Nature genetics*, 39(5), 623–630.
- Song, W. Y., Wang, G. L., Chen, L. L., Kim, H. S., Pi, L. Y., Holsten, T., *et al.* (1995). A receptor kinase-like protein encoded by the rice disease resistance gene, Xa21. *Science*, 29(2), 189–199.
- Song, Z. P., Lu, B., Zhu, Y. G., Chen, J. K. (2003). Gene flow from cultivated rice to the wild species *Oryza rufipogon* under experimental field conditions. *New phytologist*, 157(3), 657–665.
- Song, Z. P., Lu, B. R. and Chen, J. K. (2001). A study of pollen viability and longevity in *Oryza rufipogon*, *O. sativa*, and their hybrids. *International Rice Research Notes (Philippines)*.
- Swamy, B. P. and Sarla, N. (2008). Yield-enhancing quantitative trait loci (QTLs) from wild species. *Biotechnology advances*, 26(1), 106–120.
- Swapan, K. D. (2004). Rice Biotechnology: A Need for Developing Countries. *The Journal of Agrobiotechnology Management and Economics*, 7.
- Sweeney, M. and McCouch, S. R. (2007). The complex history of the domestication of rice. *Annals of Botany*.
- Takeda, T., Suwa, Y., Suzuki, M., Kitano, H., Ueguchi-Tanaka, M., Ashikari, M., *et al.* (2003). The OsTB1 gene negatively regulates lateral branching in rice. *The Plant Journal*, 33(3), 513–520.

- Tanksley, S. D. (1993). Mapping polygenes. *Annual Review of Genetics*, 27(1), 205–233.
- Tanksley, S. D. and McCouch, S. R. (1997). Seed banks and molecular maps: unlocking genetic potential from the wild. *Science*, 277(5329), 1063.
- Tanksley, S. D. and Nelson, J. C. (1996). Advanced backcross QTL analysis: a method for the simultaneous discovery and transfer of valuable QTLs from unadapted germplasm into elite breeding lines. *TAG Theoretical and Applied Genetics*, 92(2), 191–203.
- Tao, J., Zhang, L., Chong, K. and Wang, T. (2007). OsRAD21 3, an orthologue of yeast RAD21, is required for pollen development in *Oryza sativa*. *The Plant Journal*, 51(5), 919–930.
- Tenea, G. N. (2009). Exploring the world of RNA interference in plant functional genomics: a research tool for many biology phenomena. *Roumanian Biotechnological Letters*, 14(3), 4360–4364.
- Thakur, A. (2003). RNA interference revolution. *Electronic Journal of Biotechnology*, 6, 1–2.
- Thomson, M. J., Tai, T. H., McClung, A. M., Lai, X. H., Hinga, M. E., Lobos, K. B., *et al.* (2003). Mapping quantitative trait loci for yield, yield components and morphological traits in an advanced backcross population between *Oryza rufipogon* and the *Oryza sativa* cultivar Jefferson. *TAG Theoretical and Applied Genetics*, 107(3), 479–493.
- Tor, M., Lotze, M. T. and Holton, N. (2009). Receptor-mediated signalling in plants: molecular patterns and programmes. *Journal of Experimental Botany*, 1–10.

- Trotochaud, A. E., Hao, T., Wu, G., Yang, Z. and Clark, S. E. (1999). The CLAVATA1 receptor-like kinase requires CLAVATA3 for its assembly into a signaling complex that includes KAPP and a Rho-related protein. *The Plant Cell*, 11(3), 393.
- Trotochaud, A. E., Jeong, S. and Clark, S. E. (2000). CLAVATA3, a multimeric ligand for the CLAVATA1 receptor-kinase. *Science*, 289(5479), 613.
- Tumolo, A., Nguyen, Q. and Winteny, F. (1995). Detection of DNA on Membranes with Alkaline Phosphatase-Labeled Probes and Chemiluminescent CSPD Substrate. In L. Kricka (Ed.), *Nonisotopic Probing, Blotting, and Sequencing* (pp. 518): ACADEMIC PRESS.
- Vaughan, D. A., Morishima, H. and Kadowaki, K. (2003). Diversity in the Oryza genus. *Current opinion in plant biology*, 6(2), 139–146.
- Voinnet, O. (2001). RNA silencing as a plant immune system against viruses. *TRENDS in Genetics*, 17(8), 449–459.
- Voorhoeve, P. M., Sage, C. I., Schrier, M., Gillis, A. J. M., Stoop, H., Nagel, R., *et al.* (2006). A genetic screen implicates miRNA-372 and miRNA-373 as oncogenes in testicular germ cell tumors. *Cell*, 124(6), 1169–1181.
- Walker, J. C. and Zhang, R. (1990). Relationship of a putative receptor protein kinase from maize to the S-locus glycoproteins of Brassica. *Nature*, 345, 743–746.
- Wang, Z. Y., Ge, Y., Scott, M. and Spangenberg, G. (2004). Viability and longevity of pollen from transgenic and nontransgenic tall fescue (*Festuca arundinacea*)(Poaceae) plants. *American Journal of Botany*, 91(4), 523.
- Wang, P., Zhao, J., Rodrigo-Simón, A., Kian, W., Janmaat, A., Shelton, A. M., Ferré, J. and Myers, J. (2006). Mechanism of resistance to bacillus thuringiensis toxin Cry1Ac in a greenhouse population of the cabbage looper, *trichoplusia ni*. *Applied and Environmental Microbiology*, 73(4).

- Waterhouse, P. M. and Helliwell, C. A. (2003). Exploring plant genomes by RNA-induced gene silencing. *Nature Reviews Genetics*, 4(1), 29–38.
- Wesley, S. V., Helliwell, C. A., Smith, N. A., Wang, M. B., Rouse, D. T., Liu, Q., *et al.* (2001). Construct design for efficient, effective and high throughput gene silencing in plants. *The Plant Journal*, 27(6), 581–590.
- Wesley, S. V., Liu, Q., Wielopolska, A., Ellacott, G., Smith, N., Singh, S., *et al.* (2003). Custom knock-outs with hairpin RNA-mediated gene silencing. *Methods Mol Biol*, 236, 273–286.
- Wing, R. A., Kim, H. R., Goicoechea, J. L., Yu, Y., Kudrna, D., Zuccolo, A., *et al.* (2007). The Oryza map alignment project (OMAP): a new resource for comparative genomics studies within Oryza. In D. S. Brar, D. J. Mackill and B. Hardy (Eds.), *Rice Genetics V: Proceedings of the Fifth International Rice Genetics Symposium*: World Scientific Publishing Company.
- Wu, W., Ye, Q. and Min, H. (2004). Effect of straws from Bt-transgenic rice on selected biological activities in water-flooded soil. *European Journal of Soil Biology*, 40(1), 15–22.
- Xiao, J., Grandillo, S., Ahn, S. N., McCouch, S. R., Tanksley, S. D., Li, J., *et al.* (1996). Genes from wild rice improve yield. *Nature*, 384(6606), 223–224.
- Xiao, J., Li, J., Grandillo, S., Ahn, S. N., Yuan, L., Tanksley, S. D., *et al.* (1998a). Identification of Trait-Improving Quantitative Trait Loci Alleles From a Wild Rice Relative, *Oryza rufipogon*. *Genetics*, 150(2).
- Xiao, J., Li, J., Grandillo, S., Ahn, S. N., Yuan, L., Tanksley, S. D., *et al.* (1998b). Identification of trait-improving quantitative trait loci alleles from a wild rice relative, *Oryza rufipogon*. *Genetics*, 150(2), 899.
- Xie, Z., Johansen, L. K., Gustafson, A. M., Kasschau, K. D., Lellis, A. D., Zilberman, D., *et al.* (2004). Genetic and functional diversification of small RNA pathways in plants *PLoS Biol*, 2, E104.

- Xing, Y. and Zhang, Q. (2010). Genetic and molecular bases of rice yield. *Plant Biology*, 61(1), 421.
- Xu, F. Q., Li, X. R. and Ruan, Y. L. (2008). RNAi-mediated suppression of hexokinase gene OsHXK10 in rice leads to non-dehiscent anther and reduction of pollen germination. *Plant Science*, 175(5), 674–684.
- Yamamuro, C., Ihara, Y., Wu, X., Noguchi, T., Fujioka, S., Takatsuto, S., *et al.* (2000). Loss of function of a rice brassinosteroid insensitive1 homolog prevents internode elongation and bending of the lamina joint. *The Plant Cell*, 12(9), 1591.
- Yano, M. and Sasaki, T. (1997). Genetic and molecular dissection of quantitative traits in rice. *Plant Molecular Biology*, 35(1), 145–153.
- Yin, Z., Plader, W. and Malepszy, S. (2004). Transgene inheritance in plants. *Journal of applied genetics*, 45(2), 127.
- Yoon, D. B., Kang, K. H., Kim, H. J., Ju, H. G., Kwon, S. J., Suh, J. P., *et al.* (2006). Mapping quantitative trait loci for yield components and morphological traits in an advanced backcross population between *Oryza grandiglumis* and the *O. sativa japonica* cultivar Hwaseongbyeo. *TAG Theoretical and Applied Genetics*, 112(6), 1052-1062.
- Yu, L. P., Miller, A. K. and Clark, S. E. (2003). POLTERGEIST encodes a protein phosphatase 2C that regulates CLAVATA pathways controlling stem cell identity at Arabidopsis shoot and flower meristems. *Current Biology*, 13(3), 179–188.
- Yu, L. P., Simon, E. J., Trotochaud, A. E. and Clark, S. E. (2000). POLTERGEIST functions to regulate meristem development downstream of the CLAVATA loci. *Development*, 127(8), 1661–1670.

Zambon, R. A., Vakharia, V. N. and Wu, L. P. (2006). RNAi is an antiviral immune response against a dsRNA virus in *Drosophila melanogaster*. *Cellular Microbiology*, 8(5), 880–889.

Zha, X., Luo, X., Qian, X., He, G., Yang, M., Li, Y., *et al.* (2009). Over expression of the rice LRK1 gene improves quantitative yield components. *Plant biotechnology journal*, 7(7), 611–620.