

## REFERENCES

1. Adamson, E. A. S. (2010). Influence of historical landscapes, drainage evolution and ecological traits on patterns of genetic diversity in Southeast Asian freshwater snakehead fishes. Unpublished Doctor of Philosophy's thesis, Queensland University of Technology, Brisbane, Australia.
2. Adamson, E. A. S., Hurwood, D. A., & Mather, P. B. (2010). A reappraisal of the evolution of Asian snakehead fishes (Pisces, Channidae) using molecular data from multiple genes and fossil calibration. *Molecular Phylogenetics and Evolution*, 56, 707-717.
3. Alarcon, J. A., Magoulas, A., Georgakopoulos, T., Zouros, E., & Alvarez, M. C. (2004). Genetic comparison of wild and cultivated European populations of the gilthead sea bream (*Sparus aurata*). *Aquaculture*, 230, 65-80.
4. Ali, A. B. (1990). Some ecological aspects of fish populations in tropical ricefields. *Hydrobiologia*, 190, 251-222.
5. Ali, A. B. (1999). Aspects of the reproductive biology of female snakehead (*Channa striata* Bloch) obtained from irrigated rice agroecosystem, Malaysia. *Hydrobiologia*, 411, 71-77.
6. Allendorf, F. W., & Luikart, G. (2007). *Conservation and the Genetics of Populations*. Oxford, UK: Blackwell Publishing Ltd.
7. Ambak, M. A., Bolong, A. A., Ismail, P. & Tam, B. M. (2006). Genetic Variation of Snakehead Fish (*Channa striata*) Populations Using Random Amplified Polymorphic DNA. *Biotechnology*, 5(1), 104-110.
8. Amilhat, E., & Lorenzen, K. (2005). Habitat use, migration pattern and population dynamics of chevron snakehead *Channa striata* in a rainfed rice farming landscape. *Journal of Fish Biology*, 67, 23-34.

9. Amilhat, E., Lorenzen, K., Morales, E. J., Yakupitiyage, A., & Little, D. C. (2009). Fisheries production in Southeast Asian farmer managed aquatic systems (FMAS) I. Characterisation of systems. *Aquaculture*, *26*, 219-226.
10. Angers, B., Gharbi, K., & Estoup, A. (2002). Evidence of gene conversion events between paralogous sequences produced by tetraploidization in *Salmoninae* fish. *J. Mol. Evol.*, *54*, 501-510.
11. Apte, S., Star, B., & Gardner, J. P. A. (2003). A comparison of genetic diversity between cultured and wild populations, and a test for genetic introgression in the New Zealand greenshell mussel *Perna canaliculus* (Gmelin 1791). *Aquaculture*, *219*, 193-220.
12. Baie, S. H., & Sheikh, K. A. (2000). The wound healing properties of *Channa striatus*-cetrimide cream – tensile strength measurement. *Journal of Ethnopharmacology*, *71*, 93-100.
13. Balloux, F., & Lugon-Moulin, N. (2002). The estimation of population differentiation with microsatellite markers. *Molecular Ecology*, *11*, 155-165.
14. Beardmore, J. A., Mair, G. C., & Lewis, R. I. (1997). Biodiversity in aquatic systems in relation to aquaculture. *Aquaculture Research*, *28*, 829-839.
15. Bekkevold, D., Hansen, M. M., & Nielsen, E. E. (2006). Genetic impact of gadoid culture on wild fish populations: predictions, lessons from salmonids, and possibilities for minimizing adverse effects. *ICES Journal of Marine Science*, *63*, 198-208.
16. Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, *57*(1), 289-300.

17. Bolong, A. A., Ambak, M. A., Ismail, P., & Tam, B. M. (2007). Molecular data from the Cytochrome *b* for the phylogeny of Channidae (*Channa* sp.) in Malaysia. *Biotechnology*, 6(1), 22-27.
18. Boonyaratpalin, M., McCoy, E. W., & Chittapalapong, T. (1985). *Snakehead culture and its socio-economics in Thailand*. Establishment of Network of Aquaculture Centres in Asia. Thailand: FAO.
19. Botstein, D., White, R. L., Skolnick, M., & Davis, R. W. (1980). Construction of a Genetic Linkage Map in Man Using Restriction Fragment Length Polymorphisms. *Am. J. Hum. Genet.*, 32, 314-331.
20. Brookfield, J. F. Y. (1996). A simple new method for estimating null allele frequency from heterozygote deficiency. *Molecular Ecology*, 5, 453-455.
21. Chakraborty, R., De Andrade, M., Daiger, S. P., & Budowle, B. (1992). Apparent heterozygote deficiencies observed in DNA typing data and their implications in forensic applications. *Annals of Human Genetics*, 56, 45-57.
22. Chandra, S., & Banerjee, T. K. (2004). Histopathological analysis of the respiratory organs of *Channa striata* subjected to air exposure. *Veterinarski Arhiv*, 74(1), 37-52.
23. Chareontawee, K., Poompuang, S., Na-Nakorn, U., & Kamonrat, W. (2007). Genetic diversity of hatchery stocks of giant freshwater prawn (*Macrobrachium rosenbergii*) in Thailand. *Aquaculture*, 271, 121-129.
24. Chaudhry, S. (2009). *Channa striata*. In: *IUCN 2011.Red List of Threatened Species. Version 2011.1*. Retrieved from 29/04/2011, from <http://www.iucnredlist.org>.
25. Chistiakov, D. A., Hellemans, B., & Volckaert, F. A. M. (2006). Microsatellites and their genomic distribution, evolution, function and

- applications: A review with special reference to fish genetics. *Aquaculture*, 255, 1-29.
26. Courtenay, W. R., & Williams, J. D. (2004). *Snakeheads (Pisces, Channidae)– A biological synopsis and risk assessment*. Denver: U.S. Geological Survey.
  27. Crawford, N. (2010). SMOGD: software for the management of genetic diversity. *Molecular Ecology Resources*, 10, 556-557.
  28. Davis, G. P., & Hetzel, D. J. S. (2000). Integrating molecular technology with traditional approaches for genetic improvement in aquaculture species. *Aquaculture Research*, 31, 3-10.
  29. Dakin, E. E., & Avise, J. C. (2004). Microsatellite null alleles in parentage analysis. *Heredity*, 93, 504-509.
  30. DeWoody, J. A., & Avise, J. C. (2000). Microsatellite variation in marine, freshwater and anadromous fishes compared with other animals. *Journal of Fish Biology*, 56, 461-473.
  31. DeWoody, J., Nason, J. D., & Hipkins, V. D. (2006). Mitigating scoring errors in microsatellite data from wild populations. *Molecular Ecology Notes*, 6, 951-957.
  32. Diniz, D., & Bertollo, L. A. C. (2003). Karyotypic studies on *Hoplerythrinus unitaeniatus* (Pisces, Erythrinidae) populations. A biodiversity analysis. *Caryologia*, 56(3), 303-313.
  33. DOF (2011). *Channa striata*. In: *Fisheries Research Institute (FRI)*. Department of Fisheries Malaysia: Ministry of Agriculture & Agro-Based Industry Malaysia. Retrieved from 18/03/2011, from <http://www.dbserver.fri.gov.my>.
  34. DOF (2010). Malaysia: Estimated value and aquaculture production from all aquaculture system, 2000-2010. In: *Annual Fisheries Statistics, 2010*.

Department of Fisheries Malaysia: Ministry of Agriculture & Agro-Based Industry Malaysia. Retrieved from 29/12/2011, from [http://www.dof.gov.my/html/themes/moa\\_dof/documents/jld1\\_2010.pdf](http://www.dof.gov.my/html/themes/moa_dof/documents/jld1_2010.pdf).

35. Donsakul, T., & Magtoon, W. (1991). A chromosome study on five species of fishes (*Channa*, family Channidae), from Thailand. The Proceedings of the 29<sup>th</sup> Kasetsart University Annual Conference (Fisheries section) (pp. 561-574). Bangkok, Thailand: Kasetsart University.
36. Ellegren, H. (2004). Microsatellites: Simple sequences with complex evolution. *Nature Reviews Genetics*, 5, 435-445.
37. Ellegren, H. (2000). Microsatellite mutations in the germline: implications for evolutionary inference. *Trends Genet.*, 16, 551-558.
38. Evanno, G., Regnaut, S., & Goudet J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, 14, 2611-2620.
39. Excoffier, L. G. L., & Schneider, S. (2005). Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online*, 1, 47-50.
40. Excoffier, L., Estoup, A., & Cornuet, J. (2005). Bayesian Analysis of an Admixture Model With Mutations and Arbitrarily Linked Markers. *Genetics Society of America*, 169, 1727-1738.
41. Excoffier, L., Smouse, P., & Quattro, J. (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics*, 131, 479-491.
42. Falush, D., Stephens, M., & Pritchard, J. K. (2003). Inference of Population Structure Using Multilocus Genotype Data: Linked Loci and Correlated Allele Frequencies. *Genetics Society of America*, 164, 1567-1587.

43. FAO (2011a). *Channa striata* (Bloch 1793). In: *Species Fact Sheets*. Fisheries and Aquaculture Department: Food and Agricultural Organisation of the United Nations. Retrieved 20/05/2011, from <http://www.fao.org/fishery/species/3062/en>.
44. FAO (2011b). Aquaculture Resources. In: *Aquaculture topics and activities*. Fisheries and Aquaculture Department: Food and Agricultural Organisation of the United Nations. Retrieved 11/03/2011, from <http://www.fao.org/fishery/topic/13530/en>.
45. FAO (2011c). State of world aquaculture. In: *Aquaculture topics and activities*. Fisheries and Aquaculture Department: Food and Agricultural Organisation of the United Nations. Retrieved 11/03/2011, from <http://www.fao.org/fishery/topic/13540/en>.
46. FAO (2010). Contribution to the economy. In: *Sector Performance. National Aquaculture Sector Overview – Malaysia* Fisheries and Aquaculture Department: Food and Agricultural Organisation of the United Nations. Retrieved 12/11/2010, from [http://www.fao.org/fishery/countrysector/naso\\_malaysia/en](http://www.fao.org/fishery/countrysector/naso_malaysia/en).
47. Frankham, R., Ballou, J. D., & Briscoe, D. A. (2002). *A primer of conservation genetics*. New York: Cambridge University Press.
48. Froese, R., & Pauly, D. (Eds.) (2011). *Channa striata* (Bloch, 1793). *Snakehead murrel*. Retrieved 15/04/2011, from [www.fishbase.org](http://www.fishbase.org).
49. Gam, L-H., Leow, C-Y., & Saringat Baie. (2006). Proteomic analysis of snakehead fish (*Channa striata*) muscle tissue. *Malaysian Journal of Biochemistry and Molecular Biology*, 14, 25-32.

50. Gam, L-H., Leow, C-Y., & Saringat Baie. (2005). Amino acid composition of snakehead fish (*Channa striatus*) of various sizes obtained at different times of the year. *Malaysian Journal of Pharmaceutical Sciences*, 3(2), 19-30.
51. Garza, J. C., & Williamson, E. G. (2001). Detection of reduction in population size using data from microsatellite loci. *Molecular Ecology*, 10, 305-318.
52. Glaubitz, J. C. (2004). CONVERT: A user-friendly program to reformat diploid genotypic data for commonly used population genetic software packages. *Molecular Ecology Notes*, 4, 309-310.
53. Goudet, J. (2002). FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3.2). Available from <http://www.unil.ch/izea/software/fstat.html>.
54. Guo, S. W., & Thompson E. A. (1992). Performing the Exact Test of Hardy-Weinberg Proportion for Multiple Alleles. *Biometrics*, 48(2), 361-372.
55. Gupta, M. V., Acosta, B. O., Eknath, A. E., & Dey, M. M. (2000). Breakthrough in genetic improvement of tropical finfish through partnership between ICLARM, ASI and developing country NARS. Pools of Knowledge and Document Repository, Global Forum on Agricultural Research. Retrieved 30/06/2011, from [http://www.egfar.org/documents/4\\_lines/ Research\\_Partnerships/Genetic\\_Resources\\_Management/GRM\\_cases/3\\_8.pdf](http://www.egfar.org/documents/4_lines/Research_Partnerships/Genetic_Resources_Management/GRM_cases/3_8.pdf)
56. Gur-Arie, R., Cohen, C., Eitan, L., Shelef, E., Hallerman, E., & Kashi, Y. (2000). Abundance, non-random genomic distribution, nucleotide composition, and polymorphism of simple sequence repeats in *E. coli*. *Genome Res.*, 10, 61-70.
57. Ha, H. P., Nguyen, T. T. T., Poompuang, S., & Na-Nakorn, U. (2009). Microsatellites revealed no genetic differentiation between hatchery and

- contemporary wild populations of striped catfish, *Pangasianodon hypophthalmus* (Sauvage 1878) in Vietnam. *Aquaculture*, 291, 154-160.
58. Hara, M., Sekino, M., & Na-Nakorn, U. (1998). Genetic Differentiation of Natural Populations of the Snake-head Fish, *Channa striatus* in Thailand. *Fisheries Science*, 64(6), 882-885.
59. Hecht, T., & Pienaar, A. G. (1993). A review of cannibalism and its implications in fish larviculture. *J. World Aquacult. Soc.*, 24, 246-261.
60. Hoelzel, A. R. (1998). *Molecular genetic analysis of populations: a practical approach*. New York: Oxford University Press.
61. Hossain, M. K., Latifa, G. A., & Rahman, M. M. (2008). Observations on Induced Breeding of Snakehead Murrel, *Channa striatus* (Bloch, 1793). *Int. J. Sustain. Crop Prod.*, 3(5), 65-68.
62. Hundertmark, K. J., & Van Daele, L. J. (2010). Founder effect and bottleneck signatures in an introduced, insular population of elk. *Conserv. Genet.*, 11, 139-147.
63. Jakupciak, J. P., & Wells, R. D. (2000). Gene conversion (recombination) mediates expansions of CTG.CAG repeats. *Journal of Biological Chemistry*, 275, 4003-4013.
64. Jamaluddin, J. A. F., Pau, T. M., & Siti-Azizah, M. N. (2011). Genetic structure of the snakehead murrel, *Channa striata* (channidae) based on the cytochrome c oxidase subunit I gene: Influence of historical and geomorphological factors. *Genetics and Molecular Biology*, 34(1), 152-160.
65. Jost, L. (2008). GST and its relatives do not measure differentiation. *Molecular Ecology*, 17(18), 4015-4026.



66. Kalinowski, S. T., Taper, M. L., & Marshall T. C. (2007). Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology*, 16, 1099-1106.
67. Kumar, D., Marimuthu, K., Haniffa, M. A., & Sethuramalingam, T. A. (2008). Effect of different live feed on growth and survival of striped murrel *Channa striatus* larvae. *E. U. Journal of Fisheries & Aquatic Sciences*, 25(2), 105-110.
68. Kumar, V. (1995). Haplotypic differences and genetic variability revealed by mitochondria DNA in haruan, *Channa striatus* Bloch. Unpublished Degree of Bachelor of Science (Fishery)'s thesis, Universiti Putra Malaysia, Selangor, Malaysia.
69. Leberg, P. L. (2002). Estimating allelic richness: Effects of sample size and bottlenecks. *Molecular Ecology*, 11, 2445-2449.
70. Lee, P. G., & Ng, P. K. L. (1994). The systematics and ecology of snakeheads (Pisces: Channidae) in Peninsular Malaysia and Singapore. *Hydrobiologia*, 285, 59-74.
71. Lee, P. G., & Ng, P. K. L. (1991). The snakehead fishes of the Indo-Malayan Region. *Nature Malaysiana*, 16(4), 113-129.
72. Lewis, P. O., & Zaykin, D. (2001). Genetic Data Analysis: Computer program for the analysis of allelic data. Available from <http://lewis.eeb.uconn.edu/lewishome/software.html>.
73. Li, X., Musikasinthorn, P., & Kumazawa, Y. (2006). Molecular phylogenetic analyses of snakeheads (Perciformes: Channidae) using mitochondrial DNA sequences. *Ichthyol. Res.*, 53, 148-159.
74. Lind, C. E., Evans, B. S., Knauer, J., Taylor, J. J. U., & Jerry, D. R. (2009). Decreased genetic diversity and a reduced effective population size in cultured silver-lipped oysters (*Pinctada maxima*). *Aquaculture*, 286, 12-19.

75. Ling, S. W. (1977). *Aquaculture in Southeast Asia – a historical overview*. WA, U.S.A.: Washington University Press.
76. Liu, Z. J. (2007). Fish genomics and analytical genetic technologies, with examples of their potential applications in management of fish genetic resources. Policy paper for FAO's Commission on Genetic Resources for Food and Agriculture. In Bartley D., Harvey, B., & Pullin, R. (Eds.), *Workshop on Status and Trends in Aquatic Genetic Resources-A basis for international policy: FAO Fisheries Proceedings 5, FAO* (pp. 145-179). Rome.
77. Liu, Z. J., & Cordes, J. F. (2004). DNA marker technologies and their applications. *Aquaculture*, 238, 1-37.
78. Lundrigan, T. A., Reist, J. D., & Ferguson, M. M. (2005). Microsatellite genetic variation within and among Arctic charr (*Salvelinus alpinus*) from aquaculture and natural populations in North America. *Aquaculture*, 244, 63-75.
79. Magoulas, A. (1998). Application of molecular markers to aquaculture and broodstock management with special emphasis on microsatellite DNA. *Cahiers Options Mediterrannes*, 34, 153-168.
80. Mantel, N. (1967). The Detection of Disease Clustering and a Generalized Regression Approach. *Cancer Research*, 27(1), 209-220.
81. Marimuthu, K., & Haniffa, M. A. (2007). Embryonic and larval development of the striped snakehead *Channa striatus*. *Taiwania*, 52(1), 84-92.
82. Marshall, T. C., Slate, J., Kruuk, L. E., & Pemberton, J. M. (1998). Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular Ecology*, 7, 639-655.
83. Mat Jais, A. M., Abdul Rahim, M. H., Alias, R., & Muhammad N. (2009). Genetic marker for haruan *Channa striatus*. In: Proceedings of the 8<sup>th</sup> Malaysia

Congress on Genetics. Role of Genetics in Wealth Creation and Quality of Life (pp. 29). Genting Highland, Malaysia: Genetics Society of Malaysia.

84. Mat Jais, A. M. (2007a). Pharmacognosy and pharmacology of Haruan (*Channa striatus*), A medicinal fish with wound healing properties. *Bol. Latinoam. Caribe Plant. Med. Aromaticas*, 6(3), 52-60.
85. Mat Jais, A. M. (2007b). Molecular size of the bio-active components from Haruan *Channa striatus* extract. *Journal of Applied Sciences*, 7(15), 2198-2199.
86. Mat Jais, A. M., Matori, M. F., Kittakoop, P., & Suwanborirux, K. (1998). Fatty acid composition in mucus and roe of Haruan, *Channa striatus*, for wound healing. *Gen. Pharmacol.*, 30(4), 561-563.
87. Mat Jais, A. M., Dambisya, Y. M., Lee, T. L. (1997). Antinociceptive activity of *Channa striatus* (Haruan) extracts in mice. *J. Ethnopharmacol.*, 57, 125-130.
88. Mat Jais, A. M., McCulloch, R., & Croft, K. (1994). Fatty Acid and Amino Acid Composition in Haruan as a Potential Role in Wound Healing. *Gen. Pharmac.*, 25(5), 947-950.
89. Mat Jais, A. M. (1991). Haruan, *Channa striatus*, Farming in Backyard. Proceedings of the Third Asian Conference of Technology for Rural Development (pp. 91, 230-232).
90. Meirmans, P. G., & Hedrick, P. W. (2011). Assessing population structure:  $F_{ST}$  and related measures. *Molecular Ecology Resources*, 11, 5-18.
91. Mohsin, A. K., & Ambak, M. A. (1983). *Freshwater fishes of Peninsular Malaysia*. Serdang, Malaysia: University Pertanian Malaysia Press.
92. Muir, J. F., & Young, J. A. (1998). Aquaculture and marine fisheries: Will capture fisheries remain competitive?. *J. Northw. Atl. Fish. Sci.*, 23, 157-174.

93. Musikasinthorn, P. (2003). Channoidei (snakeheads). In M. Hutchins, A. Thoney, P. V. Loisel & N. Schlager (Eds.), *Grzimek's animal life encyclopedia, 2<sup>nd</sup> Edn., Vols, 4, 5. Fishes I-II* (pp. 437-447). Farmington Hills, MI: Gale Group.
94. Na-Nakorn, U., Kamonrat, W., & Ngamsiri, T. (2004). Genetic diversity of walking catfish, *Clarias macrocephalus*, in Thailand and evidence of genetic introgression from introduced farmed *C. gariepinus*. *Aquaculture*, 240, 145-163.
95. Naylor, R. L., Goldberg, R. J., Primavera, J. H., Kautsky, N., Beveridge, M. C. M., Clay, J., et al. (2000). *Effect of aquaculture on world fish supplies*. *Nature*, 405, 1017-1024.
96. Nei, M. (1978). Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics*, 89, 583-590.
97. Nei, M., & Chesser, R. K. (1983). Estimation of fixation indices and gene diversities. *Annals of Human Genetics*, 47(3), 253-259.
98. Nei, M., Maruyama, T., & Chakraborty, R. (1975). The Bottleneck Effect and Genetic Variability in Populations. *Evolution*, 29(1), 1-10.
99. Ng, M. Y. T., Ganabadi, S., & Loqman, M. Y. (2004). Effect of Orally Administered *Channa striatus* Extract Against Experimentally-Induced Osteoarthritis in Rabbits. *Intern. J. Appl. Res. Vet. Med.*, 2(3), 171-175.
100. Nguyen, T. T. T., Hurwood D., Mather, P., Na-Nakorn, U., Kamonrat, W., & Bartley, D. (2006). *Manual on applications of molecular tools in aquaculture and inland fisheries management, Part 2: Laboratory protocols and data analysis*. NACA monograph number 2. Bangkok, Thailand: FAO.
101. O'Connell, M., & Wright, J. M. (1997). Microsatellite DNA in fishes. *Reviews in Fish Biology and Fisheries*, 7, 331-363.

102. Okumus, I., & Ciftci, Y. (2003). Fish population genetics and molecular markers: II- Molecular markers and their applications in fisheries and aquaculture. *Turkish Journal of Fisheries and Aquatic Sciences*, 3, 51-79.
103. Pampoulie, C., Jorundsdottir, T. D., Steinarsson, A., Petursdottir, G., Stefansson, M. O., & Danielsdottir, A. K. (2006). Genetic comparison of experimental farmed strains and wild Icelandic populations of Atlantic cod (*Gadus morhua* L.). *Aquaculture*, 261, 556-564.
104. Parameswaran, S., & Murugesan, V. K. (1976). Breeding season and seed resources of murrels in swamps of Karnataka state. *Journal of the Inland Fisheries Society of India*, 8, 60-67.
105. Paripatananont, T. (2002). Snakehead and *Pangasius* Catfish. In C. D. Webster & C. E. Lim (Eds.), *Nutrient requirements and feeding of finfish for aquaculture* (pp. 396-401). Oxon, UK: CABI.
106. Petit, R. J., El Mousadik, A., & Pons, O. (1998). Identifying populations for conservation on the basis of genetic markers. *Conservation Biology*, 12, 844-855.
107. Poulsen, A., Griffiths, D., Nam, S., & Nguyen, T. T. (2008). Capture-based aquaculture of Pangasiid catfishes and snakeheads in the Mekong River Basin. In A. Lovatelli & P. F. Holthus (Eds.), *Capture-based aquaculture. Global review. FAO Fisheries Technical Paper. No. 508*. (pp. 69-91). Rome: FAO.
108. Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of Population Structure Using Multilocus Genotype Data. *Genetics Society of America*, 155, 945-959.
109. Qin, J. G., & Fast, A. W. (1998). Effects of temperature, size and density on culture performance of snakehead *Channa striatus* fed formulated feed. *Aquaculture Research*, 29, 299-303.

110. Qin, J. G., & Fast, A. W. (1996). Size and feed dependent cannibalism with juvenile snakehead *Channa striatus*. *Aquaculture*, 144, 313-320.
111. Qin, J. G., Fast, A. W., & Kal, A. T. (1997). Tolerance of snakehead *Channa striatus* to ammonia at different pH. *Journal of the World Aquaculture Society*, 28, 87-90.
112. Rahim, M. H. A., Rozila, A., & Mat Jais, A. M. (2009). The physical-chemical and morphological study of Haruan *Channa striatus* in Peninsular Malaysia. *Research Journal of Biological Sciences*, 4(9), 994-1009.
113. Rahman, S. A., Huah, T. S., Hassan, O., & Daud, N. M. (1995). Fatty acid composition of some Malaysian freshwater fish. *Food Chemistry*, 54, 45-49.
114. Raymond, M., & Rousset, F. (1995). GENEPOP Version 4.0: population genetics software for exact tests and ecumenicism. *J. Heredity*, 86, 248-249.
115. Riju, A., Rajesh, M. K., Sherin, F. P. T. P., Chandrasekar, A., Apshara, E. S., & Arunachalam, V. (2009). Mining of expressed sequence tag libraries of cacao for microsatellite markers using five computational tools. *Journal of Genetics*, 88(2), 217-225.
116. Saitou, N., & Nei, M. (1987). The Neighbour-joining Method: A New Method for Reconstructing Phylogenetic Trees. *Mol. Biol. Evol.*, 4(4), 406-425.
117. Salah, Y. S. M. (2010). Isolation and characterization of microsatellite loci from wild striped snakehead (*Channa striata*). Unpublished Master of Biotechnology's thesis, University of Malaya, Kuala Lumpur, Malaysia.
118. Samantaray, K., & Mohanty, S. S. (1997). Interactions of dietary levels of protein and energy on fingerling snakehead, *Channa striata*. *Aquaculture*, 156, 241-249.
119. Schlotterer, C. (2000). Evolutionary dynamics of microsatellite DNA. *Chromosoma*, 109, 365-371.

120. Schlotterer, C., & Zangerl, B. (1999). The use of imperfect microsatellites for DNA fingerprinting and population genetics. In J. T. Eppelen & T. Lubjuhn (Eds.), *DNA Profiling and DNA Fingerprinting* (pp. 153-165). Basel, Switzerland: Birkhauser.
121. Sekar, M., Suresh, E., Kumar, N. S., Nayak, S. K., & Balakrishna, C. (2009, April-June). Microsatellite DNA markers, a fisheries perspective, Part 1: The nature of microsatellites. *Aquaculture Asia Magazine*, 27-29.
122. Sekino, M., Hara, M., & Taniguchi, N. (2002). Loss of microsatellite and mitochondrial DNA variation in hatchery strains of Japanese flounder *Paralichthys olivaceus*. *Aquaculture* 213, 101-122.
123. Selkoe, K. A., & Toonen, R. J. (2006). Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. *Ecology Letters*, 9, 615-629.
124. Serapion, J., Kucuktas, H., Feng, J., & Liu, Z. (2004). Bioinformatic mining of Type I microsatellites from expressed sequence tags of channel catfish (*Ictalurus punctatus*). *Mar. Biotechnol.*, 6, 364-377.
125. Simmons, M., Mickett, K., Kucuktas, H., Li, P., Dunham, R., & Liu, Z. J. (2006). Comparison of domestic and wild channel catfish (*Ictalurus punctatus*) populations provides no evidence for genetic impact. *Aquaculture*, 252, 133-146.
126. Siriwardena, S. N. (2007). Freshwater fish seed resources and supply: Asia regional synthesis: Asia regional synthesis. In M. G. Bondad-Reantaso (Ed.), *Assessment of freshwater fish seed resources for sustainable aquaculture. FAO Fisheries Technical Paper, No. 501* (pp. 59-90). Rome: FAO.
127. Slatkin, M. (1995). A measure of population subdivision based on microsatellite allele frequencies. *Genetics*, 139, 457-462.

128. Slatkin, M. (1991). Inbreeding coefficients and coalescence times. *Genet. Res.*, 58, 167-175.
129. Slatkin, M. (1985). Rare alleles as indicators of gene flow. *Evolution*, 39, 53-65.
130. Somchit, M. N., Solihah, M. H., Israf, D. A., Ahmad, Z., Arifah, A. K., & Mat Jais, A. M. (2004). Anti-inflammatory activity of *Channa striatus*, *Channa micropeltes* and *Channa lucius* extracts: chronic inflammatory modulation. *Journal of Oriental Pharmacy and Experimental Medicine*, 4(2), 91-94.
131. Supiwong, W., Jearranai-prepame, P., & Tanomtong, A. (2009). A new report of karyotype in the chevron snakehead fish, *Channa striata* (Channidae, Pisces) from Northeast Thailand. *Cytologia*, 74(3), 1-6.
132. Tamura, K., Dudley, J., Nei, M., & Kumar, S. (2007). Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, 24, 1596-1599.
133. Tan, C. E., Chong, B. J., Sier, H. K., & Moulton, T. (1973). A report on paddy and paddy field fish production in Krian, Perak. *Ministry of Agriculture and Fisheries Malaysia, Bulletin 128*, 57.
134. Van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M., & Shipley, P. (2004). MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, 4, 535-538.
135. Varshney, R. K., Chabane, K., Hendre, P. S., Aggarwal, R. K., & Graner A. (2007). Comparative assessment of EST-SSR, EST-SNP and AFLP markers for evaluation of genetic diversity and conservation of genetic resources using wild, cultivated and elite barleys. *Plant Science*, 173, 638-649.
136. Verhoeven, K. J. F., Simonsen, K. L., & McIntyre, L. M. (2005). Implementing false discovery rate control: increasing your power. *OIKOS*, 108, 643-647.



137. Ward, R. D., & Grewe, P. M. (1994). Appraisal of molecular genetic techniques in fisheries. *Reviews in Fish Biology and Fisheries*, 4, 300-325.
138. Wee, K. L. (1982). Snakeheads – their biology and culture. In J. F. Muir & R. J. Roberts (Eds.), *Recent advances in aquaculture, Vol. 1.* (pp. 181-213). Colorado: Westview Press.
139. Wee, K. L. (1981). Snakehead (*Channa striatus*) farming in Thailand. UNDP/FAO Network of Aquaculture Centres in Asia, Bangkok.
140. Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the Analysis of Population Structure. *Evolution*, 38(6), 1358-1370.
141. Wright, S. (1951). The genetical structure of populations. *Annals of Eugenics*, 15, 323-354.
142. Xu, Z., Primavera, J. H., de la Pena, L. D., Pettit, P., Belak, J., & Alcivar-Warren, A. (2001). Genetic diversity of wild and cultured Black Tiger Shrimp (*Penaeus monodon*) in the Philippines using microsatellites. *Aquaculture*, 199, 13-40.
143. Yaakob, W. A. A., & Ali, A. B. (1992). Simple method for backyard production of snakehead (*Channa striata* Bloch) fry. *Naga*, 15(2), 22-23.
144. Yang, C., Zhu, X., & Sun, X. (2008). Development of microsatellite markers and their utilization in genetic diversity analysis of cultivated and wild populations of the mud carp (*Cirrhina molitorella*). *J. Genet. Genomics*, 35, 201-206.
145. Yeh, F. C., Yang, R. C., Boyle, T., Ye, Z. H., & Mao, J. X. (1997). POPGENE, the user-friendly shareware for population genetic analysis. University of Alberta, Canada, Molecular Biology and Biotechnology Centre. Available from <http://www.ualberta.ca/~fyeh/>.

146. Zakaria, Z. A., Mat Jais, A. M., Goh, Y. M., Sulaiman, M. R., & Somchit, M. N. (2007). Amino acid and fatty acid composition of an aqueous extract of *Channa striatus* (Haruan) that exhibits antinociceptive activity. *Clinical and Experimental Pharmacology and Physiology*, 34, 198-204.
147. Zuraini, A., Somchit, M. N., Solihah, M. H., Goh, Y. M., Arifah, A. K., Zakaria, M. S., et al. (2006). Fatty acid and amino acid composition of three local Malaysian *Channa* spp. fish. *Food Chemistry*, 97, 674-678.