

REFERENCES

- Abal, E.G. and Dennison, W.C. (1996). Seagrass depth range and water quality in southern Moreton Bay, Queensland, Australia. *Marine and Freshwater Research*, 47: 763—771.
- Annaletchumy, L., Japar Sidik, B., Muta Harah, Z. and Arshad, A. (2005). Morphology of *Halophila ovalis* (R.Br.) Hook. f. From Peninsular and East Malaysia. *Pertanika J. Trop. Agric. Sci.* 28 (1): 1-11. Universiti Putra Malaysia Press.
- Aronoff, S. (1989). *Geographic Information Systems: A Management Perspective*. WDL Publications, Ottawa, Canada. 294pp
- Aspinall, R.J. (1995). Geographic information systems: their use for environmental management and nature conservation. *Parks*, 5: 20-31.
- Bach, S.S., Borum, J., Fortes, M.D. and Duarte, C.M. (1998). Species composition and plant performance of mixed seagrass beds along a siltation gradient at Cape Bolinao, The Philippines. *Marine Ecology Progress Series*, 174: 247–256.
- Beer, S., Mtolera, M., Lyimo, T. and Bjork, M. (2006). The photosynthetic performance of the tropical seagrass *Halophila ovalis* in the upper intertidal. *Aquatic Botany*, 84: 367-371.
- Burrough, P. (1986). *Principles of Geographical Information Systems for Land Resource Assessment*. Oxford: Oxford University Press.
- Burrough, P.A. and McDonnell R.A. (1998). *Principles of Geographical Information Systems*. Oxford University Press. 352pp
- Caloz, R. and Collet, C. (1997). Geographic information systems (GIS) and remote sensing in aquatic botany: methodological aspects. *Aquatic Botany*, 58: 209-228.
- Chang, K. (2008). *Introduction to Geographic Information Systems*. McGraw-Hill International Edition. Fourth Edition. 450pp

- Constanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R.V., Paruelo, J., Raskin, R.G., Sutton, P. and van der Belt, M. (1997). The value of the world's ecosystem services and natural capital. *Nature*, 387: 253-260.
- Den Hartog, C. (1970). *The Sea-Grasses of the World*. North-Holland Publishing Co. 275pp.
- Den Hartog, C. and Polderman, P.J.G. (1975). Changes in the seagrass populations of the Dutch Waddenzee. *Aquatic Botany*, 1: 141–147.
- Dennison, W.C., Orth, R.J., Moore, K.A., Stevenson, J.C., Carter, V., Kollar, S., Bergstrom, P.W. and Batiuk, R.A. (1993). Assessing water quality with submersed aquatic vegetation. *Bioscience*, 43: 86–94.
- Department of Environment (DOE), Malaysia. (2006). *Malaysia Environmental Quality Report 2005*. <http://doe.gov.my/old/?q=en/content/environmental-quality-report-eqr-2005>
- Duarte, C.M. (1991). Seagrass depth limits. *Aquatic Botany*, 40: 363-377.
- Duarte, C.M. (1999). Seagrass ecology at the turn of the millennium: challenges for the new century. *Aquatic Botany*, 65: 7-20.
- Duarte, C.M. (2000). Marine biodiversity and ecosystem services: an elusive link. *Journal of Experimental Marine Biology and Ecology*, 250: 117-131.
- Duarte, C.M. (2010). Seagrass Meadows. *The Encyclopedia of Earth*. [http://www.eoearth.org/article/Seagrass meadows](http://www.eoearth.org/article/Seagrass%20meadows)
- Duarte, C.M. and Chiscano, C.L. (1999). Seagrass biomass and production: a reassessment. *Aquatic Botany*, 65: 159-174.
- Dunton, K. (1998). Research Issues for Texas. <http://www/tpwd.state.tx.us/conserves/coastal/seagrass/chpt1.html>.

- Environmental Systems Research Institute (ESRI). (2003). Spatial Data Standards and GIS Interoperability. <http://www.esri.com> (last accessed April 2008)
- Ferguson, R.L. and Korfmacher, K. (1997). Remote sensing and GIS analysis of seagrass meadows in North Carolina, USA. *Aquatic Botany*, 58: 241–258.
- Fortes, M.D. (1989). Seagrasses: A Resource Unknown in the ASEAN region. International Center for Living Aquatic Resource Management (ICLARM) Education Series 6, Philippines. 45 p
- Fourqurean, J.W., Boyer, J.N., Durako, M.J., Hefty, L.N. and Peterson, B.J. (2003). Forecasting Responses of Seagrass Distributions to Changing Water Quality Using Monitoring Data. *Ecological Applications*, 13(2): 474-489.
- Freeman, A.S., Short, F.T., Isnain, I., Razak, F.A. and Coles, R.G. (2008). Seagrass on the edge: Land-use practice threaten coastal seagrass communities in Sabah, Malaysia. *Biological Conservation*, 141: 2993-3005.
- Gacia, E., Duarte, C.M., Marba, N., Terrados, J., Kennedy, H., Fortes, M.D. and Tri, N.H. (2003). Sediment deposition and production in SE-Asia seagrass meadows. *Estuarine, Coastal and Shelf Science*, 56: 909-919.
- Gallegos, C. L. (1994). Refining habitat requirements of submersed aquatic vegetation: Role of optical models. *Estuaries*, 17:187-199.
- Gallegos, C. L. and Kenworthy, W. J. (1996). Seagrass depth limits in the Indian River Lagoon (Florida, USA): application of an optical water quality model. *Estuarine, Coastal and Shelf Science*, 42:267-288.
- Giesen, W.B.J.T., van Katwijk, M.M. and den Hartog, C. (1990). Eelgrass condition and turbidity in the Dutch Wadden Sea. *Aquatic Botany*, 37: 71-85.
- Hall, M.O., Durako, M.J., Fourqurean, J.W. and Zieman, J.C. (1999). Decadal changes in seagrass distribution and abundance in Florida Bay. *Estuaries*, 22: 445-459.
- Harrison, J. (1995). Finding the information. *Parks*, 5: 12-19.

- Harrison, P.G. (1989). Detrital processing in seagrass systems: A review of factors affecting decay rates, remineralisation and detritivory. *Aquatic Botany*, 23: 263-288.
- Hemminga, M.A. and Duarte, C.M. (2000). *Seagrass Ecology*. Cambridge University Press, Cambridge, England. 298pp
- Herbert, D.A. and Fourqurean, J.W. (2009). Phosphorus availability and salinity control productivity and demography of the seagrass *Thalassia testudinum* in Florida Bay. *Estuaries Coasts*, 32: 188-201.
- Hillman, K., McComb, A.J. and Walker, D.I. (1995). The distribution, biomass and primary production of the seagrass *Halophila ovalis* in the Swan/Canning Estuary, Western Australia. *Aquatic Botany*, 51: 1-54.
- ITLOS. (2003). Land Reclamation: Hearing of Malaysia's application request for provisional measures against Singapore at International Tribunal for Law of the Sea (ITLOS) from 25-27 September 2003. <http://www.itlos.org/>
- Jackson, E.L., Rowden, A.S., Attrill, M.J., Bossey, S. and Jones, M. (2001). The importance of seagrass beds as a habitat for fishery species. *Oceanography and Marine Biology, Annual Review* 39: 269-303.
- Japar Sidik, B. (1994). Status of seagrass resources in Malaysia. In: Wilkinson, C., Sudara, S. And Ming, C.L. (eds.). *Proceedings of the Third ASEAN-Australia Symposium on Living Coastal Resources*, 16-20 May 1994, Vol. 1, pp 283-289.
- Japar Sidik, B., Arshad, A., Hishamuddin, O. and Shamsul, B. (1995). *Seagrass species and distribution in Peninsular Malaysia*. Faculty of Fisheries and Marine Science. Universiti Pertanian Malaysia. 8pp
- Japar Sidik, B., Muta Harah, Z. and Arshad, A. (2006). Distribution and significance of Seagrass Ecosystems in Malaysia. *Aquatic Ecosystem Health & Management*, 9: 203-214.
- Japar Sidik, B., Muta Harah, Z., Mohd Pauzi, A. and Madhavan, S. (1999). *Halodule* species from Malaysia - distribution and morphological variation. *Aquatic Botany*, 65: 33-45.

- Kirkman, H. (1985). Community structure in seagrasses in southern Western Australia. *Aquatic Botany*, 21: 363-375.
- Kiswara, W., Huiskes, A.H.L. and Herman, P.M.J. (2005). Uptake and allocation of  $^{13}\text{C}$  by *Enhalus acoroides* at sites differing in light availability. *Aquatic Botany*, 81: 353-366.
- Koch, E. W. (2001). Beyond light: physical, geological, and geochemical parameters as possible submersed aquatic vegetation requirements. *Estuaries*, 24: 1-17.
- Koch, M.S., Schopmeyer, S.A., Kyhn-Hansen, C., Maddden, C.J. and Peters, J.S. (2007). Tropical seagrass species tolerance to hypersalinity stress. *Aquatic Botany*, 86: 14-24.
- Kuo, J and den Hartog, C. (2001). Seagrass taxonomy and identification key. In: Short, F.T. and Coles, R.G. (eds.). *Global Seagrass Research Methods*. Elsevier Science B.V., Amsterdam. pp 31-58.
- Larkum, A.W.D., McComb, A.J. and Shepherd, S.A. (eds.). (1989). *Biology of Seagrasses*. Aquatic Plants Studies 2. Elsevier, Amsterdam.
- Lathrop, R.G., Styles, R.M., Seitzinger S.P. and Bognar, J.A. (2001). Use of GIS Mapping and Modeling Approaches to Examine the Spatial Distribution of Seagrasses in Barnegat Bay, New Jersey. *Estuaries*, 24: 904-916.
- Lee Long, W.J., Mellors, J.E. and Coles, R.G. (1993). Seagrasses between Cape York and Hervey Bay, Queensland, Australia. *Australian Journal of Marine and Freshwater Research*, 44: 19-31.
- Lehmann, A. and Lachavanne, J.B. (1997). GIS and remote sensing in Aquatic Botany: Introduction. *Aquatic Botany* special issue, 58: 195-207.
- Lewis, M.A., Dantin, D.D., Chancy C.A., Abel, K.C. and Lewis, C.G. (2007). Florida seagrass habitat evaluation: A comparative survey for chemical quality. *Environmental Pollution*, 146: 206-218.
- Livingston, R.J., McGlynn, S.E. and Niu, X. (1998). Factors controlling seagrass growth in the gulf coastal system: Water and sediment quality and light. *Aquatic Botany*, 60: 135-159.

- Macinnis-Ng, C.M.O. and Ralph, P.J. (2002). Towards a more ecologically relevant assessment of the impact of heavy metals on the photosynthesis of the seagrass, *Zostera capricorni*. *Marine Pollution Bulletin*, 45: 100-106.
- Madden, C. J. and Kemp, W.M. (1996). Ecosystem model of an estuarine submersed plant community: Calibration and simulation of eutrophication responses. *Estuaries*, 19: 457-474.
- McKenzie, L.J. (2008). *Seagrass-Watch: Proceedings of a Workshop for Mapping and Monitoring Seagrass Habitats in North East Amhem Land, Northern Territory, 18-20 October 2008*. (Seagrass-Watch HQ, Cairns). 49 pp
- McKenzie, L.J. and Yoshida, R.L. (2009). *Seagrass-Watch: Proceedings of a Workshop for Monitoring Seagrass Habitats in Indonesia*. The Nature Conservancy, Coral Triangle Center, Sanur, Bali, 9<sup>th</sup> May 2009. (Seagrass-Watch HQ, Cairns). 56pp
- McKenzie, L.J., Finkbeiner, M.A. and Kirkman, H. (2001). Methods for mapping seagrass distribution. In: Short, F.T. and Coles, R.G. (eds.). *Global Seagrass Research Methods*. Elsevier Science B.V., Amsterdam. pp 101-121.
- Mellors, J., Waycott, M. and Marsh, H. (2005). Variation in biogeochemical parameters across intertidal seagrass meadows in the central Great Barrier Reef region. *Marine Pollution Bulletin*, 51: 335–342.
- Ministry of Science, Technology and Environment (MOSTE), Malaysia. (1998). *Malaysia National Policy on Biological Diversity*. <http://www.arbec.com.my/NBP.pdf>
- Norhadi, I. (1993). Preliminary Study of the Seagrass Flora of Sabah, Malaysia. *Pertanika J. Trop. Agric. Sci.*, 16 (2): 111-118.
- Orth, R.J., Carruthers, T.J.B., Dennison, W.C., Duarte, C.M., Fourqurean, J.W., Heck, K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Olyarnik, S., Short, F.T., Waycott, M. and Williams, S.L. (2006). A global crisis for seagrass ecosystems. *Bioscience*, 56: 987-996.
- Perez, M. and Romero, J. (1992). Photosynthetic response to light and temperature of the seagrass *Cymodocea nodosa* and the prediction of its seasonality. *Aquatic Botany*, 43: 51- 62.

- Phang, S.M. (1989). Seagrasses - A Neglected Natural Resource In Malaysia. In: *Proc. 12<sup>th</sup> Annual Seminar of the Malaysian Society of Marine Sciences*. pp 269-279.
- Phang, S.M. (2000). *Seagrasses of Malaysia*. University of Malaya Botanical Monographs No. 2. 60 pp
- Philip, R.C. and McRoy, C.P. (1990). *Seagrass Research Methods*. UNESCO, Paris. 210 pp
- Ralph, P.J. (1998). Photosynthetic response of laboratory-cultured *Halophila ovalis* to thermal stress. *Marine Ecology Progress Series*, 171: 123-130.
- Ralph, P.J. and Burchett, M.D. (1998). Photosynthetic response of *Halophila ovalis* to heavy metal stress. *Environmental Pollution*, 103: 91-101.
- Robbins, B.D. (1997). Quantifying temporal change in seagrass areal coverage: the use of GIS and low resolution aerial photography. *Aquatic Botany*, 58: 259-267.
- Ruiz, J.M. and Romero, J. (2003). Effects of disturbances caused by coastal constructions on spatial structure, growth dynamics and photosynthesis of the seagrass *Posidonia oceanica*. *Marine Pollution Bulletin*, 46: 1523-1533.
- Salem, B.B. (2003). Application of GIS to biodiversity monitoring. *Journal of Arid Environments*, 54: 91-114.
- Schlacher-Hoenlinger, M.A. and Schlacher, T.A. (1998). Accumulation, contamination, and seasonal variability of trace metals in the coastal zone-patterns in a seagrass meadow from the Mediterranean. *Marine Biology* 131: 401-410.
- Shepherd, S.A., McComb, A.J., Bulthuis, D.A., Neverauskas, V., Steffensen, D.A. and West, R. (1989). Decline of seagrasses. In: Larkum, A.H.D., McComb, A.J. and Shepherd, S.A. (eds.). *Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region*. Elsevier, Amsterdam. pp 346-393.
- Short, F.T. (1987). Effects of sediment nutrients on seagrasses: literature review and mesocosm experiment. *Aquatic Botany*, 27: 41-57.

- Short, F.T. and Neckles, H.A. (1999). The effects of global climate change on seagrasses. *Aquatic Botany*, 63: 169-196.
- Short, F.T. and Wyllie-Echeverria, S. (1996). Natural and human-induced disturbance of seagrasses. *Environmental Conservation*, 23: 17-27.
- Short, F.T., Burdick, D.M. and Kaldy, J.E. (1995). Mesocosm experiments quantify the effects of eutrophication on eelgrass, *Zostera marina*. *Limnology and Oceanography*, 40 (4): 740-749.
- Short, F.T., Coles, R.G. and Pergent-Martini, C. (2001). Global seagrass distribution. In: Short, F.T. and Coles, R.G. (eds.). *Global Seagrass Research Methods*. Elsevier Science B.V., Amsterdam. pp 5-30.
- Terrados, J, Agawin, N.S.R., Duarte, C.M., Fortes, M.D., Kamp-Nielsen, L. and Borum, J. (1999). Nutrient limitation of the tropical seagrass *Enhalus acoroides* (L.) Royle in Cape Bolinao, NW Philippines. *Aquatic Botany*, 65: 123-139.
- Terrados, J. and Duarte, C.M. (1999). Experimental evidence of reduced particle resuspension within a seagrass (*Posidonia oceanica* L.) meadow. *Journal of Experimental Marine Biology and Ecology*, 243: 45-53.
- Torre-Castro, M.D.L. and Ronnback, P. (2004). Links between humans and seagrasses - an example from tropical East Africa. *Ocean & Coastal Management*, 47: 361-387.
- United Nations Environment Programme (UNEP). (1992). *Convention on Biological Diversity*. <http://www.cbd.int/>
- Walker, D.I. and Cambridge, M.L. (1995). An experimental assessment of the temperature responses of two sympatric seagrasses, *Amphibolis antarctica* and *Amphibolis griffithii*, in relation to their biogeography. *Hydrobiologia*, 302: 63-70.
- Walker, D.I., Kendrick, G.A. and McComb, A.J. (1988). The distribution of seagrass species in Shark Bay, Western Australia, with notes on their ecology. *Aquatic Botany*, 30: 305-317.



Ward, D. H., Markon, C. J. and Douglas D. C. (1997). Distribution and stability of eelgrass beds at Izembek Lagoon, Alaska. *Aquatic Botany*, 58: 229-240.

<http://www.aseansec.org/cme/Marine%20Water%20Quality%20Criteria%20for%20the%20ASEAN%20Region.pdf>

<http://www.seagrasswatch.org>