

## Bibliography

Abcouwer SF. Basic molecular biological methods in surgical research: genetic library construction, screening, and DNA sequencing. *Surgical research*: 217-232

Barnwell P, Blanchard AN, Bryant JA, Smirnoff N, and Weir AF (1998). Isolation of DNA from highly mucilaginous succulent plant *Sedum telephium*. *Plant molecular biology reporter*, 16: 133-138.

Bewley JD (1997). Seed germination and dormancy. *The Plant Cell*, 9: 1055-1066.

Brown TA (2001). *Gene cloning and DNA analysis: an introduction* 4<sup>th</sup> edition. Blackwell Science, UK pp 156-178.

Chauthaiwale VM, Therwath A and Deshpande VV (1992). Bacteriophage lambda as a cloning vector. *American Society for Microbiology*, 56(4): 577-591.

Cornel Mulhardt, E.W. Beese M.D (2007). Cloning DNA fragments. *Molecular biology and genomics*: 105-129

Couch JA and Fritz PJ (1990). Isolation of DNA from plant high in polyphenolics. *Plant Molecular Biology Reporter*, 8: 8-12.

De la Cruz M, Ramirez F and Hernandez H (1997). DNA isolation and amplification from cacti. *Plant molecular biology reporter*, 15: 319–325.

Diadema K, Baumel A, Lebris M and Affre L (2003). Genomic DNA isolation and amplification from callus culture in succulent Plants, *Carpobrotus* Species (Aizoaceae). *Journal of plant molecular biology reporter*, 21: 173a–173e

Downham A and Collins P (2000). Colouring our food in the last and next millennium. *Journal of food science and technology*, 35: 5-22.

Falcon ADV (2002). The human genome and its importance in clinical medicine. *International Congress Series*, 1237: 3 –13

Hanania U, Velcheva M, Sahar N and Peryl A (2004). An improved method for isolating high quality DNA from *Vitis vinefera* nuclei. *Plant molecular biology reporter*, 22: 173-177.

Hanley ME, Byron BL, Meredith MF, Rafferty CM (2007). Plant structural traits and their role in anti-herbivore defence. *Journal of Perspectives in Plant Ecology, Evolution and Systematics*, 8: 157–178.

He RF, Wang Y, Shi Z, Ren X, Zhu L, Weng Q and He GC (2003). Construction of a genomic library of wild rice and agrobacterium-mediated transformation of large insert DNA linked to BPH resistance locus. *Gene*, 321:113–121.

- Koornneef M, Bentsink L and Hilhorst H (2002). Seed dormancy and germination. *Journal of current opinion in plant biology*, 5: 33-36
- Lambda FIX II/*Xho* I Partial Fill-In Vector Kit: *Xho* I/Gigapack III XL Packaging Extract (Stratagene) and Klenow Fill-In Kit (Stratagene).
- Le Bellec F, Vaillant F and Imbert E (2006). Pitahaya (*Hylocereus* spp.): a new fruit crop, a market with a future. *Fruits*, 61 (4): 237-250
- Lichtenzveig J, Abbo S, Nerd A, Tel-zur N and Mizrahi Y (2000). Cytology and mating systems in the climabing cacti *Hylocereus* and *Selenicereus*. *American Journal of Botany*, 87(7): 1058–1065.
- Lin Y, Chow T, Luo M, Kudrna D, Lin C, Wing RA and Hsing YC (2006). Two highly representative rice BAC libraries of japonica cv Tainung 67 suitable for rice structural and functional genomic research. *Plant Science*, 170: 889–896
- Lodhi MA, Ye G, Weeden NF, and Reisch BI (1994). A simple and efficient method for DNA extraction from grapevine cultivars and *Vitis* species. *Plant Molecular Biology Reporter*, 12(1): 6-13.
- Maliyakal EJ (1992). An efficient method for isolation of RNA and DNA from plants containing polyphenolics. *Nucleic Acids Research*, 20(9):2381.
- Mayer K and Mewes HW (2001). How can we deliver the large plant genome? Strategies and perspectives. *Current opinion in plant biology*, 5:173-177.
- Mei YQ and Song SQ (2008). Early morphological and physiological events occurring during germination of maize seeds. *Journal of agricultural sciences in china*, 7(8): 950-957
- Michiels A, Van den Ende W, Tukcern M, Van Riet L and Van Laere A (2003). Extraction of high-quality genomic DNA from latex-containing plants. *Analytical biochemistry*, 315(1): 85-89.
- Michelmore R (2000). Genomic approaches to plant disease resistance. *Current opinion in plant biology*, 3:125-131.
- Murray MG and Thompson WF (1980). Rapid isolation of high molecular weight plant DNA. *Nucleic acids research*, 8: 4321–4325.
- Nagesha HS, Yu M and Wang LF (1996). Application of linker-ligation-PCR for construction of phage display epitope libraries. *Journal of Virological Methods*, 60: 147-154.
- Nerd A, Gutman F and Mizrahi Y (1999). Ripening and postharvest behaviour of fruits of two *Hylocereus* species (Cactaceae). *Journal of postharvest biology and technology* 17: 39–45.

- Pearson TRH, Burslem DFRP, Mullins CE and Dalling JW (2003). Functional significance of photoblastic germination in neotropical pioneer trees: a seed's eye view. *Functional Ecology*, 17(3): 394-402.
- Puchooa D (2004). A simple, rapid and efficient method for the extraction of genomic DNA from lychee (*Litchi chinensis* Sonn.). *African journal of biotechnology*, 3(4): 253-255.
- Rahman S, Ohto M, Seraj ZI and Blumwald E (2007). Bacteriophage Lambda Genomic Library Construction of *Oryza sativa* L. var. Pokkali. *Plant Tissue Cult. & Biotech.* 17(2): 149-159
- Rajjou L and Debeaujon I (2008). Seed longevity: survival and maintenance of high germination ability of dry seeds. *Journal of comptes rendus biologiques*, 331(10): 796-805
- Rojas-arehchiga M and Vahzquez-yanes C (2000). Cactus seed germination: a review. *Journal of arid environments*, 44: 85-104.
- Salzman RA, Fujita T, Zhu-Sazlman K, Hasegawa PM and Bressan RA (1999). An improved RNA isolation method for plant tissues containing high levels of phenolic compounds or carbohydrates. *Plant molecular biology reporter*, 17: 11-17.
- Sambrook J, Fritsch EF and Maniatis T (1989). *Molecular cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Shan W and Hardham AR (2004). Construction of a bacterial artificial chromosome library, determination of genome size, and characterization of an Hsp70 gene family in *Phytophthora nicotianae*. *Fungal Genetics and Biology*, 41(3): 369-380
- Shashikant CS, Carr JL, Bhargava J, Bentley KL and Ruddle FH (1998). Recombinogenic targeting: a new approach to genomic analysis-a review. *Gene*, 223: 9-20.
- Sharma AD, Gill PK and Singh P (2002). DNA isolation from dry and fresh samples of polysaccharide-rich plants. *Plant Molecular Biology Reporter*, 20: 415a-415f
- Sheffield WP (1997). Concepts and techniques in molecular biology: an overview. *Transfusion Medicine Reviews*, 11(3): 209-223.
- Simao E, Socolowski F and Takaki M (2007). The epiphytic cactaceae *Hylocereus setaceus* (Salm-Dick ex DC) Ralf Bauer seed germination is controlled by light and temperature. *Journal of Brazilian archives of biology and technology*, 50(4): 655-662.
- Tel-Zur N, Abbo S, Myslabodski D and Mizrahi Y (1999). Modified CTAB procedure for DNA isolation from epiphytic cacti of the Genera *Hylocereus* and *Selenicereus* (Cactaceae). *Journal of plant molecular biology reporter*, 17: 249-254.

Tel-Zur N, Abbo S, Bar-Zvi D and Mizrahi Y (2004). Genetic relationships among *Hylocereus* and *Selenicereus* vine cacti (Cactaceae): evidence from hybridization and cytological studies. *Annals of Botany*, 94: 527–534

Tel-Zur N, Abbo S, and Mizrahi Y (2005). Cytogenetics of Semi Fertile Triploid and Aneuploid Intergeneric Vine Cacti Hybrids. *Journal of heredity*, 96(2): 1–8.

Van Buuren ML, Lanfranco L, Longato Sabina, Minerdi D, Harrison MJ and Bonfante P (1999). Construction and characterization of genomic libraries of two endomycorrhizal fungi: *Glomus versiforme* and *Gigaspora margarita*. *Mycological Research*, 103(8): 955-960

Wang Y, Cao Z, Hood D and Townsel JG (2003). Construction of genomic libraries in  $\lambda$ -vectors. 153-168

[www.doa.gov.my](http://www.doa.gov.my)

## Appendix

### Obtaining Genomic DNA for Construction of Genomic Library

Lee K.P. Daniel, Loo Z.X. and Chandran S.  
Institute of Biological Sciences  
University of Malaya  
Malaysia

**Keywords:** Genomic library, dragon fruit, restriction digest, high molecular weight DNA, Cactaceae

#### Abstract

The isolation of pure genomic DNA from the white and red dragon fruits, *Hylocereus undatus* and *Hylocereus polyrhizus* (Cactaceae) for the construction of a genomic library is described. These vine cacti originate from northern South America, Central America and Mexico, and are cultivated as new exotic fruit crops in SE Asia. Successful establishment of a genomic library serves as a platform for molecular studies, such as gene characterization and gene analysis, but intact and high molecular weight DNA of good quality can be difficult to obtain. Furthermore, presence of polysaccharides in DNA extracts interferes with downstream processes such as restriction digest and polymerase chain reaction (PCR). Genomic DNA can be extracted using various plant parts from 2–3-week-old seedling leaves, roots and stems. DNA quality is checked using a spectrophotometer and digested using *Bam*H1. Genomic DNA so isolated had the highest quality following restriction digestion analysis.

#### INTRODUCTION

Dragon fruits of the genera *Hylocereus* and *Selenicereus* which originate from northern South America, Central America, and Mexico, are currently being grown as new exotic fruit crops in SE Asia. The fruit ‘flesh’ largely includes a mucilaginous placenta in which is found thousands of small seeds (Le Bellec *et al.*, 2006).

At present, only three species are being cultivated on a commercial scale which is *Hylocereus undatus*, *Hylocereus polyrhizus* and *Selenicereus megalanthus* in Colombia, Israel, Vietnam and Nicaragua (N. Tel-Zur *et al.*, 2005). Vietnam is the biggest producer of the *Hylocereus* spp. However, only *H. undatus* and *H. polyrhizus* are being cultivated in small scale in Malaysia.

These three species are distinguished by peel and pulp colour. *H. polyrhizus* fruit has red peel and red pulp while *H. undatus* fruit has red peel and white pulp. *S. megalanthus* has yellow peel and white pulp. The red colouration is due to the presence of betalain pigments. Plantations of such non-native fruit species are affected by various diseases that lead to poor fruit yield, threatening the income of small-scale farmers who resort to application of pesticides and herbicides. This problem can be solved by establishing a genomic library that serve as a platform to study the gene interaction. The overall procedure involved in genomic library construction includes isolation of genomic DNA, generation of DNA fragments for cloning, packaging and transduction.

## MATERIALS AND METHODS

### Plant material

Red and white dragon fruits were obtained from a local market. Their seeds were separated from the pulp and air-dried. The seeds were allowed to germinate for about two weeks and seedling leaves were used for DNA extraction.

### Reagents and solutions

DNA isolation buffer [0.15M Sorbitol, 0.125M Tris-base, 0.05M EDTA,  $8.3 \times 10^{-3}$ M Sodium Bis-sulfide, 0.83M NaCl, 2% w/v hexadecyltrimethyl-ammonium bromide (CTAB), 0.83% v/v Sodium Sarkosyl, 1% w/v Polyvinylpyrrolidone (PVP) (MW 40,000), 1% v/v  $\beta$ -mercaptoethanol] Chloroform: isoamyl alcohol (24:1), 5M NaCl, Isopropanol, 70% ethanol, TE buffer [10mM Tris-HCl (pH 8.0), 1mM EDTA (pH 8.0)], RNase A (0.1 $\mu$ g/ $\mu$ l)

### DNA isolation

A total of 400 $\mu$ l of DNA isolation buffer was added into each 1.5-ml microfuge tube. Leaf fragments from 2-week-old seedlings were weighed and pulverized in liquid nitrogen. 0.20g of leaf sample was transferred into each buffer-added microfuge tube. Microfuge tubes were quick-spun. An additional 300 $\mu$ l of DNA isolation buffer was added. All tubes were gently flicked and inverted to thoroughly mix the sample and incubated at 65°C for 40 minutes (tubes were gently flicked and inverted at 10-minute intervals). 700 $\mu$ l of chloroform: isoamyl alcohol (24:1) was added to each microfuge tube and tubes were gently inverted until an emulsion formed. Samples were centrifuged at 10,000 rpm for 15 minutes. The aqueous phase was gently transferred into new 1.5ml microfuge tubes. The chloroform: isoamyl alcohol procedure was repeated. Half volume of 5M NaCl was added and microfuge tubes were gently inverted. Two volumes of cold isopropanol (-20°C) was added into each microfuge tube and tubes were gently inverted before being incubated on ice for 10 minutes. Samples were centrifuged at 10,000 rpm for 5 minutes. The supernatant was discarded. DNA pellets were washed twice with 500 $\mu$ l of cold 70% ethanol (-20°C). Samples were centrifuged at 10,000 rpm for 1 minute. The supernatant was discarded and the pellet washing procedure repeated. DNA pellets were air-dried at room temperature, and then dissolved in 30 to 50 $\mu$ l of TE buffer (depending on their solubility). RNase A was added to each sample (1 $\mu$ l/10 $\mu$ l DNA sample), which was then incubated at 37°C for 1 hour. After RNase A treatment, total DNA was run on a 0.8% agarose gel against a marker of known molecular weight, and stained with ethidium bromide for visualization (Figure 1). The samples were stored at -20°C.

### Analytical analysis

Extracted DNA samples were quantified using a spectrophotometer at  $A_{260\text{nm}}$  while the purity of the DNA was checked through  $A_{260\text{nm}}/A_{280\text{nm}}$  (Table 1).

### Partial digestion of the Genomic DNA

Partial digestion of the genomic DNA was carried out with the enzyme *Bam*H1 according to the Stockinger Lab 06/01/05 protocol. After digestion, the product was analyzed through 0.8% agarose gel electrophoresis (Figure 2).

## RESULTS AND DISCUSSION

There are several published works that have reported on the use of different plant parts for obtaining genomic DNA, such as callus tissue and roots. Roots were chosen as the source tissue due to the lower viscosity of the extracts relative to that from other plant parts. Leaf callus, essentially an unorganized and undifferentiated cell mass, lacks synthesis of secondary metabolites and so may be more easily dealt with (Diadema *et al.*, 2003 and N. Tel-Zur *et al.*, 1999).

White and red dragon fruits contain high amounts of polysaccharides in their tissues generally. The polysaccharides bind to the DNA, rendering it viscous and glutinous after precipitation of the DNA. This makes the DNA unsuitable for downstream applications such as PCR and restriction digest, as the applied enzyme cannot access the DNA (N. Tel-Zur *et al.*, 1999, Barnwell *et al.*, 1998 and Puchooa, 2004). However, DNA extraction with CTAB in high salt condition suppresses the co-precipitation of polysaccharides and DNA (Murray and Thompson, 1980; Lodhi *et al.*, 1994).

In plant cells, there is an abundance of secondary metabolites. To prevent the interaction of these compounds with DNA,  $\beta$ -mercaptoethanol and/or polyvinyl pyrrolidone (PVP) is often added to the buffer solution. PVP forms complexes with polyphenolics through hydrogen bonding and  $\beta$ -mercaptoethanol reduces oxidation of phenolic compounds (Micheils *et al.*, 2003, De la Cruz *et al.*, 1997 and Maliyakal, 1992).

High yield and quality of DNA from leaf extractions were attributed to active cell division, which produces high cell density and little or no synthesis of secondary metabolites. In addition, the extracted genomic DNA can be restricted using enzyme *Bam*H1.

Table 1 show that DNA extraction using leaves provides higher purity compared to other parts of the plants. Figure 1 also shows that DNA extraction using young leaves yielded a higher amount of DNA. However, when the same extraction protocol was carried out on stem tissue, the extract was viscous throughout the extraction process due to the presence of high amount of polysaccharides (N. Tel-Zur *et al.*, 1999). The root extraction carried out in this study follows the protocol by N. Tel-Zur *et al.*, 1999 and resulted in a brown pellet. This may be caused by the binding of secondary metabolites to the DNA which results in low yield of DNA. Furthermore, Figure 2 confirms that the DNA extract from leaves is of high quality, implying that the enzyme efficiently digested total DNA into smaller fragments.

## CONCLUSION

From the experiment conducted, young seedling leaves gave the best quality of DNA and yield. This was confirmed through spectrophotometer and gel electrophoresis assessments. The ratio of absorbance at 260nm to 280nm fall between 1.8-2.0 range and the extracted genomic DNA can be restricted.

## Literature cited

- Barnwell, P., Blanchard, A. N., Bryant, J. A., Smirnoff, N. and Weir, A. F. 1998. Isolation of DNA from the highly mucilaginous succulent plant *Sedum telephium*. J. Plant Mol. Biol. Rptr. 16: 133–138.
- De la Cruz, M., Ramirez, F. and Hernandez, H. 1997. DNA isolation and amplification from cacti. J. Plant Mol Biol. Rptr. 15: 319–325.
- Diadema, K., Baumel, A., Lebris, M. and Affre, L. 2003. Genomic DNA isolation and amplification from callus culture in succulent plants, *Carpobrotus* Species (Aizoaceae). J. Plant Mol. Biol. Rptr. 21: 173a–173e.
- Khan, S., Qureshi, M. I., Kamaluddin, Alam, T. and Abdin, M. Z. 2007. Protocol for isolation of genomic DNA from dry and fresh roots of medicinal plants suitable for RAPD and restriction digestion. J. African Biotechnology, 6(3): 175-178
- Le Bellec, F., Vaillant, F. and Imbert, E. 2006. Pitahaya (*Hylocereus* spp.): a new fruit crop, a market with a future. Fruits 61 (4): 237-250.
- Lodhi, M. A., Ye, G. N., Weeden, N.F. and Reisch, B. I. 1994. A simple and efficient method for DNA extraction from grapevine cultivars and *Vitis* species. J. Plant Mol. Biol. Rptr. 12: 6–13.
- Maliyakal, E. J. 1992. An efficient method for isolation of RNA and DNA from plants containing polyphenolics. J. Nucleic Acid Res. 20(9): 2381
- Michiels, A., Van den Ende, W., Tukcern, M., Van Riet, L. and Van Laere, A. 2003. Extraction of high-quality genomic DNA from latex-containing plants. J. Analytical Biochem. 315(1): 85-89.
- Murray, M. G. and Thompson, W.F. 1980. Rapid isolation of high molecular weight plant DNA. J. Nucleic Acids Res. 8(19): 4321–4325.
- Puchooa, D. 2004. A simple, rapid and efficient method for the extraction of genomic DNA from lychee (*Litchi chinensis* Sonn.). J. African Biotechnology, 3(4): 253-255.
- Stockinger lab protocol 06/01/05 (Bacteriophage  $\lambda$  plant genomic library construction) <http://www.oardc.ohiostate.edu/stockingerlab/Protocols/GenomicDNAPhageLibraryConstruction.pdf>.
- Tel-Zur, N., Abbo, S. and Mizrahi, Y. 2005. Cytogenetics of semi-fertile triploid and aneuploid intergeneric vine cacti hybrids. J. of Heredity, 96 (2):1-8.
- Tel-Zur N, Abbo, S., Myslabodski, D. and Mizrahi, Y. 1999. Modified CTAB procedure for DNA isolation from epiphytic cacti of the genera *Hylocereus* and *Selenicereus* (Cactaceae). J. Plant Mol. Biol. Rptr. 17: 249–254

## Tables

Table 1. DNA purity and DNA yield using different plant materials

Plant material	A <sub>260nm</sub>	A <sub>280nm</sub>	Ratio (A <sub>260nm</sub> / A <sub>280nm</sub> )	DNA yield (μg/g)
Stem	0.008	0.007	1.14	20.0
Root	0.023	0.018	1.28	57.5
Leaves ( White dragon fruit)	0.081	0.042	1.92	202.5
Leaves (Red dragon fruit)	0.093	0.049	1.90	232.5

## Figures




Fig. 1. Electrophoresis of genomic DNA extracted using different plant materials. A- Lamda DNA ; B- stem ; C- Root; D- Leaves *Hylocereus undatus* ; E- Leaves *Hylocereus polyrhizus* ; F = 1kb DNA ladder




Fig. 2. Electrophoresis of *Bam*H1 restricted genomic DNA extracted from *Hylocereus* leaves. A- Lambda DNA ; B- Lambda DNA digested with *Bam*H1; C- 1kb DNA ladder; D- Total DNA *Hylocereus* ; E- Total DNA *Hylocereus* digested with *Bam*H1