

## References

- Abdulla, M. A., Ahmed, I., Assawamakin, A., Bhak, J., Brahmachari, S. K., Calacal, G. C. (2009). Mapping human genetic diversity in Asia. *Science*, 326(5959), 1541-1545.
- Adeyemo, A., Chen, G., Zhou, J., Shriver, D., Doumatey, A., Huang, H. (2010). FTO genetic variation and association with obesity in West Africans and African Americans. *Diabetes*, 59(6), 1549-1554.
- Ahima, R. S. (2006). Metabolic actions of adipocyte hormones: focus on adiponectin. *Obesity (Silver Spring)*, 14 Suppl 1, 9S-15S.
- Ahima, R. S., & Flier, J. S. (2000). Adipose tissue as an endocrine organ. *Trends Endocrinol Metab*, 11(8), 327-332.
- Ahmad, S. M. S. a. R. (2004). Childhood obesity: contributing factors, consequences and intervention. *Malaysian Journal of Nutrition*, 10(1), 13-22.
- Al-Attar, S. A., Pollex, R. L., Ban, M. R., Young, T. K., Bjerregaard, P., Anand, S. S. (2008). Association between the FTO rs9939609 polymorphism and the metabolic syndrome in a non-Caucasian multi-ethnic sample. *Cardiovasc Diabetol*, 7, 5.
- Al-Daghri, N., Chetty, R., McTernan, P. G., Al-Rubean, K., Al-Attas, O., Jones, A. F. (2005). Serum resistin is associated with C-reactive protein & LDL cholesterol in type 2 diabetes and coronary artery disease in a Saudi population. *Cardiovasc Diabetol*, 4(1), 10.
- Albertson, D. G. (2006). Gene amplification in cancer. *Trends Genet*, 22(8), 447-455.
- Albu, J. B., Murphy, L., Frager, D. H., Johnson, J. A., & Pi-Sunyer, F. X. (1997). Visceral fat and race-dependent health risks in obese nondiabetic premenopausal women. *Diabetes*, 46(3), 456-462.
- Andreasen, C. H., Mogensen, M. S., Borch-Johnsen, K., Sandbaek, A., Lauritzen, T., Sorensen, T. I. (2008). Non-replication of genome-wide based associations between common variants in INSIG2 and PFKP and obesity in studies of 18,014 Danes. *PLoS One*, 3(8), e2872.
- Andreasen, C. H., Stender-Petersen, K. L., Mogensen, M. S., Torekov, S. S., Wegner, L., Andersen, G. (2008). Low physical activity accentuates the effect of the FTO rs9939609 polymorphism on body fat accumulation. *Diabetes*, 57(1), 95-101.
- Apal Sammy, Y. D., Ming, M. F., Rampal, S., Bulgiba, A., & Mohamed, Z. (2011). Gender-Dependent Association of a beta2- Adrenergic Gene Variant With Obesity Parameters in Malaysian Malays. *Asia Pac J Public Health*.
- Ardlie, K. G., Kruglyak, L., & Seielstad, M. (2002). Patterns of linkage disequilibrium in the human genome. *Nat Rev Genet*, 3(4), 299-309.

- Arita, Y., Kihara, S., Ouchi, N., Takahashi, M., Maeda, K., Miyagawa, J. (1999). Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun*, 257(1), 79-83.
- Asano, H., Izawa, H., Nagata, K., Nakatochi, M., Kobayashi, M., Hirashiki, A. (2010). Plasma resistin concentration determined by common variants in the resistin gene and associated with metabolic traits in an aged Japanese population. *Diabetologia*, 53(2), 234-246.
- Azmi MY, J. R., Siti Mariam A, Safiah MY, Fatimah S, Norimah AK4, Poh BK, Kandiah M, Zalilah MS, Wan Abdul Manan WM, Siti Haslinda MD & Tahir A. (2009). Body Mass Index (BMI) of Adults: Findings of the Malaysian Adult Nutrition Survey (MANS). *Mal J Nutr*, 15(2), 97 - 119.
- Azuma, K., Katsukawa, F., Oguchi, S., Murata, M., Yamazaki, H., Shimada, A. (2003). Correlation between serum resistin level and adiposity in obese individuals. *Obes Res*, 11(8), 997-1001.
- Bamba, V., & Rader, D. J. (2007). Obesity and atherogenic dyslipidemia. *Gastroenterology*, 132(6), 2181-2190.
- Banerji, M. A., Lebowitz, J., Chaiken, R. L., Gordon, D., Kral, J. G., & Lebovitz, H. E. (1997). Relationship of visceral adipose tissue and glucose disposal is independent of sex in black NIDDM subjects. *Am J Physiol*, 273(2 Pt 1), E425-432.
- Barber, J. C., Maloney, V., Hollox, E. J., Stuke-Sontheimer, A., du Bois, G., Daumiller, E. (2005). Duplications and copy number variants of 8p23.1 are cytogenetically indistinguishable but distinct at the molecular level. *Eur J Hum Genet*, 13(10), 1131-1136.
- Bardel, C., Danjean, V., Hugot, J. P., Darlu, P., & Genin, E. (2005). On the use of haplotype phylogeny to detect disease susceptibility loci. *BMC Genet*, 6, 24.
- Bauer, F., Elbers, C. C., Adan, R. A., Loos, R. J., Onland-Moret, N. C., Grobbee, D. E. (2009). Obesity genes identified in genome-wide association studies are associated with adiposity measures and potentially with nutrient-specific food preference. *Am J Clin Nutr*, 90(4), 951-959.
- Bell, C. G., Walley, A. J., & Froguel, P. (2005). The genetics of human obesity. *Nat Rev Genet*, 6(3), 221-234.
- Bell, J. (2004). Predicting disease using genomics. *Nature*, 429(6990), 453-456.
- Ben Ali, S., Kallel, A., Ftouhi, B., Sediri, Y., Feki, M., Slimane, H. (2009). Association of G-2548A LEP polymorphism with plasma leptin levels in Tunisian obese patients. *Clin Biochem*, 42(7-8), 584-588.
- Berentzen, T., Kring, S. I., Holst, C., Zimmermann, E., Jess, T., Hansen, T. (2008). Lack of association of fatness-related FTO gene variants with energy expenditure or physical activity. *J Clin Endocrinol Metab*, 93(7), 2904-2908.
- Berg, A. H., & Scherer, P. E. (2005). Adipose tissue, inflammation, and cardiovascular disease. *Circ Res*, 96(9), 939-949.

- Berndt, C., Casaroli-Marano, R. P., Vilaro, S., & Reina, M. (2001). Cloning and characterization of human syndecan-3. *J Cell Biochem*, 82(2), 246-259.
- Bernfield, M., Gotte, M., Park, P. W., Reizes, O., Fitzgerald, M. L., Lincecum, J. (1999). Functions of cell surface heparan sulfate proteoglycans. *Annu Rev Biochem*, 68, 729-777.
- Bird, A. P. (1996). The relationship of DNA methylation to cancer. *Cancer Surv*, 28, 87-101.
- Boes, E., Kollerits, B., Heid, I. M., Hunt, S. C., Pichler, M., Paulweber, B. (2008). INSIG2 polymorphism is neither associated with BMI nor with phenotypes of lipoprotein metabolism. *Obesity (Silver Spring)*, 16(4), 827-833.
- Boissel, S., Reish, O., Proulx, K., Kawagoe-Takaki, H., Sedgwick, B., Yeo, G. S. (2009). Loss-of-function mutation in the dioxygenase-encoding FTO gene causes severe growth retardation and multiple malformations. *Am J Hum Genet*, 85(1), 106-111.
- Bottner, A., Kratzsch, J., Muller, G., Kapellen, T. M., Bluher, S., Keller, E. (2004). Gender differences of adiponectin levels develop during the progression of puberty and are related to serum androgen levels. *J Clin Endocrinol Metab*, 89(8), 4053-4061.
- Bouchard, C., Tremblay, A., Despres, J. P., Nadeau, A., Lupien, P. J., Theriault, G. (1990). The response to long-term overfeeding in identical twins. *N Engl J Med*, 322(21), 1477-1482.
- Bravard, A., Lefai, E., Meugnier, E., Pesenti, S., Disse, E., Vouillarmet, J. (2011). FTO is increased in muscle during type 2 diabetes, and its overexpression in myotubes alters insulin signaling, enhances lipogenesis and ROS production, and induces mitochondrial dysfunction. *Diabetes*, 60(1), 258-268.
- Bray, G., & Bouchard, C. (1997). Genetics of human obesity: research directions. *FASEB J*, 11(12), 937-945.
- Bray, G. A. (1991). Treatment for obesity: a nutrient balance/nutrient partition approach. *Nutr Rev*, 49(2), 33-45.
- Bray, M. S., Krushkal, J., Li, L., Ferrell, R., Kardina, S., Sing, C. F. (2000). Positional genomic analysis identifies the beta(2)-adrenergic receptor gene as a susceptibility locus for human hypertension. *Circulation*, 101(25), 2877-2882.
- Breitfeld, J., Stumvoll, M., & Kovacs, P. (2012). Genetics of adiponectin. *Biochimie*.
- Bressler, J., Fornage, M., Hanis, C. L., Kao, W. H., Lewis, C. E., McPherson, R. (2009). The INSIG2 rs7566605 genetic variant does not play a major role in obesity in a sample of 24,722 individuals from four cohorts. *BMC Med Genet*, 10, 56.
- Brichard, S. M., Delporte, M. L., & Lambert, M. (2003). Adipocytokines in anorexia nervosa: a review focusing on leptin and adiponectin. *Horm Metab Res*, 35(6), 337-342.

- Briscoe, D., Stephens, J. C., & O'Brien, S. J. (1994). Linkage disequilibrium in admixed populations: applications in gene mapping. *J Hered*, 85(1), 59-63.
- Bronner, G., Sattler, A. M., Hinney, A., Soufi, M., Geller, F., Schafer, H. (2006). The 103I variant of the melanocortin 4 receptor is associated with low serum triglyceride levels. *J Clin Endocrinol Metab*, 91(2), 535-538.
- Brown, C. D., Higgins, M., Donato, K. A., Rohde, F. C., Garrison, R., Obarzanek, E. (2000). Body mass index and the prevalence of hypertension and dyslipidemia. *Obes Res*, 8(9), 605-619.
- Brown, M. S., & Goldstein, J. L. (1999). A proteolytic pathway that controls the cholesterol content of membranes, cells, and blood. *Proc Natl Acad Sci U S A*, 96(20), 11041-11048.
- Burnett, M. S., Lee, C. W., Kinnaird, T. D., Stabile, E., Durrani, S., Dullum, M. K. (2005). The potential role of resistin in atherogenesis. *Atherosclerosis*, 182(2), 241-248.
- Burstein, R., Prentice, A. M., Goldberg, G. R., Murgatroyd, P. R., Harding, M., & Coward, W. A. (1996). Metabolic fuel utilisation in obese women before and after weight loss. *Int J Obes Relat Metab Disord*, 20(3), 253-259.
- Calabro, P., Golia, E., Maddaloni, V., Malvezzi, M., Casillo, B., Marotta, C. (2009). Adipose tissue-mediated inflammation: the missing link between obesity and cardiovascular disease? *Intern Emerg Med*, 4(1), 25-34.
- Calabro, P., Limongelli, G., Pacileo, G., Di Salvo, G., Golino, P., & Calabro, R. (2008). The role of adiposity as a determinant of an inflammatory milieu. *J Cardiovasc Med (Hagerstown)*, 9(5), 450-460.
- Calle, E. E. (2007). Obesity and cancer. *BMJ*, 335(7630), 1107-1108.
- Calle, E. E., & Kaaks, R. (2004). Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer*, 4(8), 579-591.
- Cameron, A. J., Magliano, D. J., Shaw, J. E., Zimmet, P. Z., Carstensen, B., Alberti, K. G. (2012). The influence of hip circumference on the relationship between abdominal obesity and mortality. *Int J Epidemiol*.
- Campa, D., Husing, A., McKay, J. D., Sinilnikova, O., Vogel, U., Tjonneland, A. (2010). The INSIG2 rs7566605 polymorphism is not associated with body mass index and breast cancer risk. *BMC Cancer*, 10, 563.
- Cao, H., & Hegele, R. A. (2001). Single nucleotide polymorphisms of the resistin (RSTN) gene. *J Hum Genet*, 46(9), 553-555.
- Carey, D. J., Conner, K., Asundi, V. K., O'Mahony, D. J., Stahl, R. C., Showalter, L. (1997). cDNA cloning, genomic organization, and in vivo expression of rat N-syndecan. *J Biol Chem*, 272(5), 2873-2879.

- Cargill, M., Altshuler, D., Ireland, J., Sklar, P., Ardlie, K., Patil, N. (1999). Characterization of single-nucleotide polymorphisms in coding regions of human genes. *Nat Genet*, 22(3), 231-238.
- Caro, J. F., Sinha, M. K., Kolaczynski, J. W., Zhang, P. L., & Considine, R. V. (1996). Leptin: the tale of an obesity gene. *Diabetes*, 45(11), 1455-1462.
- Carr, M. C., & Brunzell, J. D. (2004). Abdominal obesity and dyslipidemia in the metabolic syndrome: importance of type 2 diabetes and familial combined hyperlipidemia in coronary artery disease risk. *J Clin Endocrinol Metab*, 89(6), 2601-2607.
- Carroll, L., Voisey, J., & van Daal, A. (2005). Gene polymorphisms and their effects in the melanocortin system. *Peptides*, 26(10), 1871-1885.
- Cauchi, S., Stutzmann, F., Cavalcanti-Proenca, C., Durand, E., Pouta, A., Hartikainen, A. L. (2009). Combined effects of MC4R and FTO common genetic variants on obesity in European general populations. *J Mol Med (Berl)*, 87(5), 537-546.
- Cavalli-Sforza, L. L. (1966). Population structure and human evolution. *Proc R Soc Lond B Biol Sci*, 164(995), 362-379.
- Cecil, J. E., Tavendale, R., Watt, P., Hetherington, M. M., & Palmer, C. N. (2008). An obesity-associated FTO gene variant and increased energy intake in children. *N Engl J Med*, 359(24), 2558-2566.
- Cha, S., Koo, I., Choi, S. M., Park, B. L., Kim, K. S., Kim, J. R. (2009). Association analyses of the INSIG2 polymorphism in the obesity and cholesterol levels of Korean populations. *BMC Med Genet*, 10, 96.
- Cha, S., Koo, I., Park, B. L., Jeong, S., Choi, S. M., Kim, K. S. (2009). Genetic Effects of FTO and MC4R Polymorphisms on Body Mass in Constitutional Types. *Evid Based Complement Alternat Med*.
- Cha, S. W., Choi, S. M., Kim, K. S., Park, B. L., Kim, J. R., Kim, J. Y. (2008). Replication of genetic effects of FTO polymorphisms on BMI in a Korean population. *Obesity (Silver Spring)*, 16(9), 2187-2189.
- Chambers, J. C., Elliott, P., Zabaneh, D., Zhang, W., Li, Y., Froguel, P. (2008). Common genetic variation near MC4R is associated with waist circumference and insulin resistance. *Nat Genet*, 40(6), 716-718.
- Chang, Y. C., Liu, P. H., Lee, W. J., Chang, T. J., Jiang, Y. D., Li, H. Y. (2008). Common variation in the fat mass and obesity-associated (FTO) gene confers risk of obesity and modulates BMI in the Chinese population. *Diabetes*, 57(8), 2245-2252.
- Chapman, N. H., & Wijsman, E. M. (1998). Genome screens using linkage disequilibrium tests: optimal marker characteristics and feasibility. *Am J Hum Genet*, 63(6), 1872-1885.

- Chauhan, G., Tabassum, R., Mahajan, A., Dwivedi, O. P., Mahendran, Y., Kaur, I. (2011). Common variants of FTO and the risk of obesity and type 2 diabetes in Indians. *J Hum Genet*, 56(10), 720-726.
- Chen, C. C., Li, T. C., Li, C. I., Liu, C. S., Wang, H. J., & Lin, C. C. (2005). Serum resistin level among healthy subjects: relationship to anthropometric and metabolic parameters. *Metabolism*, 54(4), 471-475.
- Chen, Y., Gotte, M., Liu, J., & Park, P. W. (2008). Microbial subversion of heparan sulfate proteoglycans. *Mol Cells*, 26(5), 415-426.
- Cheung, C. Y., Tso, A. W., Cheung, B. M., Xu, A., Ong, K. L., Fong, C. H. (2010). Obesity susceptibility genetic variants identified from recent genome-wide association studies: implications in a chinese population. *J Clin Endocrinol Metab*, 95(3), 1395-1403.
- Cho, Y. M., Youn, B. S., Chung, S. S., Kim, K. W., Lee, H. K., Yu, K. Y. (2004). Common genetic polymorphisms in the promoter of resistin gene are major determinants of plasma resistin concentrations in humans. *Diabetologia*, 47(3), 559-565.
- Choi, S. W., & Friso, S. (2010). Epigenetics: A New Bridge between Nutrition and Health. *Adv Nutr*, 1(1), 8-16.
- Chrysohoou, C., Panagiotakos, D. B., Pitsavos, C., Skoumas, I., Papademetriou, L., Economou, M. (2007). The implication of obesity on total antioxidant capacity in apparently healthy men and women: the ATTICA study. *Nutr Metab Cardiovasc Dis*, 17(8), 590-597.
- Chung, W. K., Patki, A., Matsuoka, N., Boyer, B. B., Liu, N., Musani, S. K. (2009). Analysis of 30 genes (355 SNPS) related to energy homeostasis for association with adiposity in European-American and Yup'ik Eskimo populations. *Hum Hered*, 67(3), 193-205.
- Church, C., Lee, S., Bagg, E. A., McTaggart, J. S., Deacon, R., Gerken, T. (2009). A mouse model for the metabolic effects of the human fat mass and obesity associated FTO gene. *PLoS Genet*, 5(8), e1000599.
- Clark, A. G. (2004). The role of haplotypes in candidate gene studies. *Genet Epidemiol*, 27(4), 321-333.
- Clement, K. (2006). Genetics of human obesity. *C R Biol*, 329(8), 608-622; discussion 653-605.
- Clement, K., Vaisse, C., Lahlou, N., Cabrol, S., Pelloux, V., Cassuto, D. (1998). A mutation in the human leptin receptor gene causes obesity and pituitary dysfunction. *Nature*, 392(6674), 398-401.

- Clifton, I. J., McDonough, M. A., Ehrismann, D., Kershaw, N. J., Granatino, N., & Schofield, C. J. (2006). Structural studies on 2-oxoglutarate oxygenases and related double-stranded beta-helix fold proteins. *J Inorg Biochem*, 100(4), 644-669.
- Combs, T. P., Berg, A. H., Rajala, M. W., Klebanov, S., Iyengar, P., Jimenez-Chillaron, J. C. (2003). Sexual differentiation, pregnancy, calorie restriction, and aging affect the adipocyte-specific secretory protein adiponectin. *Diabetes*, 52(2), 268-276.
- Considine, R. V., Sinha, M. K., Heiman, M. L., Kriauciunas, A., Stephens, T. W., Nyce, M. R. (1996). Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med*, 334(5), 292-295.
- Constantin, A., Costache, G., Sima, A. V., Glavce, C. S., Vladica, M., & Popov, D. L. (2010). Leptin G-2548A and leptin receptor Q223R gene polymorphisms are not associated with obesity in Romanian subjects. *Biochem Biophys Res Commun*, 391(1), 282-286.
- Coop, G., Pickrell, J. K., Novembre, J., Kudaravalli, S., Li, J., Absher, D. (2009). The role of geography in human adaptation. *PLoS Genet*, 5(6), e1000500.
- Corella, D., Lai, C. Q., Demissie, S., Cupples, L. A., Manning, A. K., Tucker, K. L. (2007). APOA5 gene variation modulates the effects of dietary fat intake on body mass index and obesity risk in the Framingham Heart Study. *J Mol Med (Berl)*, 85(2), 119-128.
- Corella, D., Peloso, G., Arnett, D. K., Demissie, S., Cupples, L. A., Tucker, K. (2009). APOA2, dietary fat, and body mass index: replication of a gene-diet interaction in 3 independent populations. *Arch Intern Med*, 169(20), 1897-1906.
- Costa, V., Casamassimi, A., & Ciccodicola, A. (2010). Nutritional genomics era: opportunities toward a genome-tailored nutritional regimen. *J Nutr Biochem*, 21(6), 457-467.
- Dallongeville, J., Helbecque, N., Cottel, D., Amouyel, P., & Meirhaeghe, A. (2003). The Gly16-->Arg16 and Gln27-->Glu27 polymorphisms of beta2-adrenergic receptor are associated with metabolic syndrome in men. *J Clin Endocrinol Metab*, 88(10), 4862-4866.
- Degawa-Yamauchi, M., Bovenkerk, J. E., Juliar, B. E., Watson, W., Kerr, K., Jones, R. (2003). Serum resistin (FIZZ3) protein is increased in obese humans. *J Clin Endocrinol Metab*, 88(11), 5452-5455.
- Deka, R., Xu, L., Pal, P., Toelue, P. T., Laumoli, T. S., Xi, H. (2009). A tagging SNP in INSIG2 is associated with obesity-related phenotypes among Samoans. *BMC Med Genet*, 10, 143.
- Devlin, B., & Risch, N. (1995). A comparison of linkage disequilibrium measures for fine-scale mapping. *Genomics*, 29(2), 311-322.

- Diez, J. J., & Iglesias, P. (2003). The role of the novel adipocyte-derived hormone adiponectin in human disease. *Eur J Endocrinol*, 148(3), 293-300.
- Dina, C., Meyre, D., Gallina, S., Durand, E., Korner, A., Jacobson, P. (2007). Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet*, 39(6), 724-726.
- Do, R., Bailey, S. D., Desbiens, K., Belisle, A., Montpetit, A., Bouchard, C. (2008). Genetic variants of FTO influence adiposity, insulin sensitivity, leptin levels, and resting metabolic rate in the Quebec Family Study. *Diabetes*, 57(4), 1147-1150.
- Donohoe, C. L., Pidgeon, G. P., Lysaght, J., & Reynolds, J. V. (2010). Obesity and gastrointestinal cancer. *Br J Surg*, 97(5), 628-642.
- Dorajoo, R., Blakemore, A. I., Sim, X., Ong, R. T., Ng, D. P., Seielstad, M. (2012). Replication of 13 obesity loci among Singaporean Chinese, Malay and Asian-Indian populations. *Int J Obes (Lond)*, 36(1), 159-163.
- Duarte, S. F., Francischetti, E. A., Genelhu, V. A., Cabello, P. H., & Pimentel, M. M. (2007). LEPR p.Q223R, beta3-AR p.W64R and LEP c.-2548G>A gene variants in obese Brazilian subjects. *Genet Mol Res*, 6(4), 1035-1043.
- Echwald, S. M., Sorensen, T. I., Tybjaerg-Hansen, A., Andersen, T., & Pedersen, O. (1998). Gln27Glu variant of the human beta2-adrenoreceptor gene is not associated with early-onset obesity in Danish men. *Diabetes*, 47(10), 1657-1658.
- Ehrenborg, E., Skogsberg, J., Ruotolo, G., Large, V., Eriksson, P., Arner, P. (2000). The Q/E27 polymorphism in the beta2-adrenoceptor gene is associated with increased body weight and dyslipoproteinaemia involving triglyceride-rich lipoproteins. *J Intern Med*, 247(6), 651-656.
- Engeli, S., & Sharma, A. M. (2000). Role of adipose tissue for cardiovascular-renal regulation in health and disease. *Horm Metab Res*, 32(11-12), 485-499.
- Fair, A. M., & Montgomery, K. (2009). Energy balance, physical activity, and cancer risk. *Methods Mol Biol*, 472, 57-88.
- Fantuzzi, G. (2005). Adipose tissue, adipokines, and inflammation. *J Allergy Clin Immunol*, 115(5), 911-919; quiz 920.
- Farooqi, I., Keogh, J. M., Yeo, G. S., Lank, E. J., Cheetham, T., & O'Rahilly, S. (2003). Clinical spectrum of obesity and mutations in the melanocortin 4 receptor gene. *N Engl J Med*, 348(12), 1085-1095.
- Farooqi, I. S., & O'Rahilly, S. (2004). Monogenic human obesity syndromes. *Recent Prog Horm Res*, 59, 409-424.
- Farooqi, I. S., & O'Rahilly, S. (2005). Monogenic obesity in humans. *Annu Rev Med*, 56, 443-458.



- Farooqi, I. S., & O'Rahilly, S. (2007). Genetic factors in human obesity. *Obes Rev*, 8 Suppl 1, 37-40.
- Fawcett, K. A., & Barroso, I. (2010). The genetics of obesity: FTO leads the way. *Trends Genet*, 26(6), 266-274.
- Feng, Y., Dong, H., Xiang, Q., Hong, X., Wilker, E., Zhang, Y. (2007). Lack of association between rs7566605 and obesity in a Chinese population. *Hum Genet*, 120(5), 743-745.
- Feramisco, J. D., Goldstein, J. L., & Brown, M. S. (2004). Membrane topology of human insig-1, a protein regulator of lipid synthesis. *J Biol Chem*, 279(9), 8487-8496.
- Feuk, L., Carson, A. R., & Scherer, S. W. (2006). Structural variation in the human genome. *Nat Rev Genet*, 7(2), 85-97.
- Feuk, L., Marshall, C. R., Wintle, R. F., & Scherer, S. W. (2006). Structural variants: changing the landscape of chromosomes and design of disease studies. *Hum Mol Genet*, 15 Spec No 1, R57-66.
- Filippi, E., Sentinelli, F., Trischitta, V., Romeo, S., Arca, M., Leonetti, F. (2004). Association of the human adiponectin gene and insulin resistance. *Eur J Hum Genet*, 12(3), 199-205.
- Fischer, J., Koch, L., Emmerling, C., Vierkotten, J., Peters, T., Bruning, J. C. (2009). Inactivation of the Fto gene protects from obesity. *Nature*, 458(7240), 894-898.
- Fischer, T. P., Burnard, P., Marty, B., Hilton, D. R., Furi, E., Palhol, F. (2009). Upper-mantle volatile chemistry at Oldoinyo Lengai volcano and the origin of carbonatites. *Nature*, 459(7243), 77-80.
- Flatt, J. P. (1995). McCollum Award Lecture, 1995: diet, lifestyle, and weight maintenance. *Am J Clin Nutr*, 62(4), 820-836.
- Flier, J. S. (2004). Obesity wars: molecular progress confronts an expanding epidemic. *Cell*, 116(2), 337-350.
- Ford, S., Jr., Bozian, R. C., & Knowles, H. C., Jr. (1968). Interactions of obesity, and glucose and insulin levels in hypertriglyceridemia. *Am J Clin Nutr*, 21(9), 904-910.
- Franks, P. W., Jablonski, K. A., Delahanty, L. M., McAteer, J. B., Kahn, S. E., Knowler, W. C. (2008). Assessing gene-treatment interactions at the FTO and INSIG2 loci on obesity-related traits in the Diabetes Prevention Program. *Diabetologia*, 51(12), 2214-2223.
- Frayling, T. M., Timpson, N. J., Weedon, M. N., Zeggini, E., Freathy, R. M., Lindgren, C. M. (2007). A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science*, 316(5826), 889-894.

- Frayn, K. N. (2002). Insulin resistance, impaired postprandial lipid metabolism and abdominal obesity. A deadly triad. *Med Princ Pract, 11 Suppl 2*, 31-40.
- Frazer, K. A., Ballinger, D. G., Cox, D. R., Hinds, D. A., Stuve, L. L., Gibbs, R. A. (2007). A second generation human haplotype map of over 3.1 million SNPs. *Nature, 449*(7164), 851-861.
- Freathy, R. M., Timpson, N. J., Lawlor, D. A., Pouta, A., Ben-Shlomo, Y., Ruukonen, A. (2008). Common variation in the FTO gene alters diabetes-related metabolic traits to the extent expected given its effect on BMI. *Diabetes, 57*(5), 1419-1426.
- Frederich, R. C., Lollmann, B., Hamann, A., Napolitano-Rosen, A., Kahn, B. B., Lowell, B. B. (1995). Expression of ob mRNA and its encoded protein in rodents. Impact of nutrition and obesity. *J Clin Invest, 96*(3), 1658-1663.
- Friedman, J. M., & Halaas, J. L. (1998). Leptin and the regulation of body weight in mammals. *Nature, 395*(6704), 763-770.
- Friedman, J. M., Leibel, R. L., Siegel, D. S., Walsh, J., & Bahary, N. (1991). Molecular mapping of the mouse ob mutation. *Genomics, 11*(4), 1054-1062.
- Gabriel, S., Ziaugra, L., & Tabbaa, D. (2009). SNP genotyping using the Sequenom MassARRAY iPLEX platform. *Curr Protoc Hum Genet, Chapter 2*, Unit 2 12.
- Gabriel, S. B., Schaffner, S. F., Nguyen, H., Moore, J. M., Roy, J., Blumenstiel, B. (2002). The structure of haplotype blocks in the human genome. *Science, 296*(5576), 2225-2229.
- Galic, S., Oakhill, J. S., & Steinberg, G. R. (2010). Adipose tissue as an endocrine organ. *Mol Cell Endocrinol, 316*(2), 129-139.
- Galletti, F., Iacone, R., Ragone, E., Russo, O., Della Valle, E., Siani, A. (2004). Lack of association between polymorphism in the beta2-adrenergic receptor gene, hypertension, and obesity in the Olivetti heart study. *Am J Hypertens, 17*(8), 718-720.
- Gantz, I., Konda, Y., Tashiro, T., Shimoto, Y., Miwa, H., Munzert, G. (1993). Molecular cloning of a novel melanocortin receptor. *J Biol Chem, 268*(11), 8246-8250.
- Gavrila, A., Peng, C. K., Chan, J. L., Mietus, J. E., Goldberger, A. L., & Mantzoros, C. S. (2003). Diurnal and ultradian dynamics of serum adiponectin in healthy men: comparison with leptin, circulating soluble leptin receptor, and cortisol patterns. *J Clin Endocrinol Metab, 88*(6), 2838-2843.
- Geffroy, S., De Vos, P., Staels, B., Duban, B., Auwerx, J., & de Martinville, B. (1995). Localization of the human OB gene (OBS) to chromosome 7q32 by fluorescence in situ hybridization. *Genomics, 28*(3), 603-604.

- Geller, F., Reichwald, K., Dempfle, A., Illig, T., Vollmert, C., Herpertz, S. (2004). Melanocortin-4 receptor gene variant I103 is negatively associated with obesity. *Am J Hum Genet*, 74(3), 572-581.
- Gerber, M., Boettner, A., Seidel, B., Lammert, A., Bar, J., Schuster, E. (2005). Serum resistin levels of obese and lean children and adolescents: biochemical analysis and clinical relevance. *J Clin Endocrinol Metab*, 90(8), 4503-4509.
- Gerken, T., Girard, C. A., Tung, Y. C., Webby, C. J., Saudek, V., Hewitson, K. S. (2007). The obesity-associated FTO gene encodes a 2-oxoglutarate-dependent nucleic acid demethylase. *Science*, 318(5855), 1469-1472.
- Goldstein, B. J., & Scalia, R. (2004). Adiponectin: A novel adipokine linking adipocytes and vascular function. *J Clin Endocrinol Metab*, 89(6), 2563-2568.
- Goldstein, J. L., DeBose-Boyd, R. A., & Brown, M. S. (2006). Protein sensors for membrane sterols. *Cell*, 124(1), 35-46.
- Gonzalez-Sanchez, J. L., Zabena, C., Martinez-Larrad, M. T., Martinez-Calatrava, M. J., Perez-Barba, M., & Serrano-Rios, M. (2009). Variant rs9939609 in the FTO gene is associated with obesity in an adult population from Spain. *Clin Endocrinol (Oxf)*, 70(3), 390-393.
- Gonzalez, J. R., Gonzalez-Carpio, M., Hernandez-Saez, R., Serrano Vargas, V., Torres Hidalgo, G., Rubio-Rodrigo, M. (2011). FTO Risk Haplotype Among Early Onset and Severe Obesity Cases in a Population of Western Spain. *Obesity (Silver Spring)*.
- Goodman, E., Dolan, L. M., Morrison, J. A., & Daniels, S. R. (2005). Factor analysis of clustered cardiovascular risks in adolescence: obesity is the predominant correlate of risk among youth. *Circulation*, 111(15), 1970-1977.
- Goossens, G. H., Petersen, L., Blaak, E. E., Hul, G., Arner, P., Astrup, A. (2009). Several obesity- and nutrient-related gene polymorphisms but not FTO and UCP variants modulate postabsorptive resting energy expenditure and fat-induced thermogenesis in obese individuals: the NUGENOB study. *Int J Obes (Lond)*, 33(6), 669-679.
- Gordi, T., & Khamis, H. (2004). Simple solution to a common statistical problem: interpreting multiple tests. *Clin Ther*, 26(5), 780-786.
- Grant, S. F., Li, M., Bradfield, J. P., Kim, C. E., Annaiah, K., Santa, E. (2008). Association analysis of the FTO gene with obesity in children of Caucasian and African ancestry reveals a common tagging SNP. *PLoS One*, 3(3), e1746.
- Green, S. A., Turki, J., Innis, M., & Liggett, S. B. (1994). Amino-terminal polymorphisms of the human beta 2-adrenergic receptor impart distinct agonist-promoted regulatory properties. *Biochemistry*, 33(32), 9414-9419.
- Grundy, S. M. (1998). Multifactorial causation of obesity: implications for prevention. *Am J Clin Nutr*, 67(3 Suppl), 563S-572S.

- Grundy, S. M., Brewer, H. B., Jr., Cleeman, J. I., Smith, S. C., Jr., & Lenfant, C. (2004). Definition of metabolic syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Circulation*, *109*(3), 433-438.
- Grunnet, L. G., Nilsson, E., Ling, C., Hansen, T., Pedersen, O., Groop, L. (2009). Regulation and function of FTO mRNA expression in human skeletal muscle and subcutaneous adipose tissue. *Diabetes*, *58*(10), 2402-2408.
- Gu, H. F. (2009). Biomarkers of adiponectin: plasma protein variation and genomic DNA polymorphisms. *Biomark Insights*, *4*, 123-133.
- Guilbert, J. J. (2003). The world health report 2002 - reducing risks, promoting healthy life. *Educ Health (Abingdon)*, *16*(2), 230.
- Haemer, M. A., Huang, T. T., & Daniels, S. R. (2009). The effect of neurohormonal factors, epigenetic factors, and gut microbiota on risk of obesity. *Prev Chronic Dis*, *6*(3), A96.
- Hakanen, M., Raitakari, O. T., Lehtimaki, T., Peltonen, N., Pahkala, K., Sillanmaki, L. (2009). FTO genotype is associated with body mass index after the age of seven years but not with energy intake or leisure-time physical activity. *J Clin Endocrinol Metab*, *94*(4), 1281-1287.
- Halaas, J. L., Gajiwala, K. S., Maffei, M., Cohen, S. L., Chait, B. T., Rabinowitz, D. (1995). Weight-reducing effects of the plasma protein encoded by the obese gene. *Science*, *269*(5223), 543-546.
- Hall, D. H., Rahman, T., Avery, P. J., & Keavney, B. (2006). INSIG-2 promoter polymorphism and obesity related phenotypes: association study in 1428 members of 248 families. *BMC Med Genet*, *7*, 83.
- Halushka, M. K., Fan, J. B., Bentley, K., Hsie, L., Shen, N., Weder, A. (1999). Patterns of single-nucleotide polymorphisms in candidate genes for blood-pressure homeostasis. *Nat Genet*, *22*(3), 239-247.
- Hampton, R. Y. (2002). Proteolysis and sterol regulation. *Annu Rev Cell Dev Biol*, *18*, 345-378.
- Hanchard, N., Elzein, A., Trafford, C., Rockett, K., Pinder, M., Jallow, M. (2007). Classical sickle beta-globin haplotypes exhibit a high degree of long-range haplotype similarity in African and Afro-Caribbean populations. *BMC Genet*, *8*, 52.
- Hancock, A. M., Witonsky, D. B., Gordon, A. S., Eshel, G., Pritchard, J. K., Coop, G. (2008). Adaptations to climate in candidate genes for common metabolic disorders. *PLoS Genet*, *4*(2), e32.
- Hara, K., Boutin, P., Mori, Y., Tobe, K., Dina, C., Yasuda, K. (2002). Genetic variation in the gene encoding adiponectin is associated with an increased risk of type 2 diabetes in the Japanese population. *Diabetes*, *51*(2), 536-540.

- Hardie, D. G. (2003). Minireview: the AMP-activated protein kinase cascade: the key sensor of cellular energy status. *Endocrinology*, *144*(12), 5179-5183.
- Hassanein, M. T., Lyon, H. N., Nguyen, T. T., Akyzbekova, E. L., Waters, K., Lettre, G. (2010). Fine mapping of the association with obesity at the FTO locus in African-derived populations. *Hum Mol Genet*, *19*(14), 2907-2916.
- Haupt, A., Thamer, C., Machann, J., Kirchhoff, K., Stefan, N., Tschritter, O. (2008). Impact of variation in the FTO gene on whole body fat distribution, ectopic fat, and weight loss. *Obesity (Silver Spring)*, *16*(8), 1969-1972.
- Haupt, A., Thamer, C., Staiger, H., Tschritter, O., Kirchhoff, K., Machicao, F. (2009). Variation in the FTO gene influences food intake but not energy expenditure. *Exp Clin Endocrinol Diabetes*, *117*(4), 194-197.
- Havel, P. J., Kasim-Karakas, S., Dubuc, G. R., Mueller, W., & Phinney, S. D. (1996). Gender differences in plasma leptin concentrations. *Nat Med*, *2*(9), 949-950.
- Havel, P. J., Kasim-Karakas, S., Mueller, W., Johnson, P. R., Gingerich, R. L., & Stern, J. S. (1996). Relationship of plasma leptin to plasma insulin and adiposity in normal weight and overweight women: effects of dietary fat content and sustained weight loss. *J Clin Endocrinol Metab*, *81*(12), 4406-4413.
- Hayakawa, T., Nagai, Y., Kahara, T., Yamashita, H., Takamura, T., Abe, T. (2000). Gln27Glu and Arg16Gly polymorphisms of the beta2-adrenergic receptor gene are not associated with obesity in Japanese men. *Metabolism*, *49*(9), 1215-1218.
- Hedenbro JL, C. M., Orho M. (1999). No association between human beta2-adrenoceptor gene polymorphism and obesity or type 2 diabetes *Diabetological* *41*(42), 98-101.
- Heid, I. M., Henneman, P., Hicks, A., Coassin, S., Winkler, T., Aulchenko, Y. S. (2010). Clear detection of ADIPOQ locus as the major gene for plasma adiponectin: results of genome-wide association analyses including 4659 European individuals. *Atherosclerosis*, *208*(2), 412-420.
- Heid, I. M., Vollmert, C., Hinney, A., Doring, A., Geller, F., Lowel, H. (2005). Association of the 103I MC4R allele with decreased body mass in 7937 participants of two population based surveys. *J Med Genet*, *42*(4), e21.
- Heid, I. M., Vollmert, C., Kronenberg, F., Huth, C., Ankerst, D. P., Luchner, A. (2008). Association of the MC4R V103I polymorphism with the metabolic syndrome: the KORA Study. *Obesity (Silver Spring)*, *16*(2), 369-376.
- Heid, I. M., Wagner, S. A., Gohlke, H., Iglseider, B., Mueller, J. C., Cip, P. (2006). Genetic architecture of the APM1 gene and its influence on adiponectin plasma levels and parameters of the metabolic syndrome in 1,727 healthy Caucasians. *Diabetes*, *55*(2), 375-384.

- Hennig, B. J., Fulford, A. J., Sirugo, G., Rayco-Solon, P., Hattersley, A. T., Frayling, T. M. (2009). FTO gene variation and measures of body mass in an African population. *BMC Med Genet*, 10, 21.
- Herbert, A., Gerry, N. P., McQueen, M. B., Heid, I. M., Pfeufer, A., Illig, T. (2006). A common genetic variant is associated with adult and childhood obesity. *Science*, 312(5771), 279-283.
- Hertel, J. K., Johansson, S., Raeder, H., Midthjell, K., Lyssenko, V., Groop, L. (2008). Genetic analysis of recently identified type 2 diabetes loci in 1,638 unselected patients with type 2 diabetes and 1,858 control participants from a Norwegian population-based cohort (the HUNT study). *Diabetologia*, 51(6), 971-977.
- Hinney, A., Nguyen, T. T., Scherag, A., Friedel, S., Bronner, G., Muller, T. D. (2007). Genome wide association (GWA) study for early onset extreme obesity supports the role of fat mass and obesity associated gene (FTO) variants. *PLoS One*, 2(12), e1361.
- Hinuy, H. M., Hirata, M. H., Forti, N., Diament, J., Sampaio, M. F., Armaganijan, D. (2008). Leptin G-2548A promoter polymorphism is associated with increased plasma leptin and BMI in Brazilian women. *Arq Bras Endocrinol Metabol*, 52(4), 611-616.
- Hirsch, J., & Leibel, R. L. (1998). The genetics of obesity. *Hosp Pract (Minneap)*, 33(3), 55-59, 62-55, 69-70 passim.
- Hivert, M. F., Manning, A. K., McAteer, J. B., Dupuis, J., Fox, C. S., Cupples, L. A. (2009). Association of variants in RETN with plasma resistin levels and diabetes-related traits in the Framingham Offspring Study. *Diabetes*, 58(3), 750-756.
- Hjelm Cluff, A., Malmstrom, A., Tingaker, B., David, G., & Ekman-Ordeberg, G. (2005). Normal labor associated with changes in uterine heparan sulfate proteoglycan expression and localization. *Acta Obstet Gynecol Scand*, 84(3), 217-224.
- Holcomb, I. N., Kabakoff, R. C., Chan, B., Baker, T. W., Gurney, A., Henzel, W. (2000). FIZZ1, a novel cysteine-rich secreted protein associated with pulmonary inflammation, defines a new gene family. *EMBO J*, 19(15), 4046-4055.
- Horikoshi, M., Hara, K., Ito, C., Shojima, N., Nagai, R., Ueki, K. (2007). Variations in the HHEX gene are associated with increased risk of type 2 diabetes in the Japanese population. *Diabetologia*, 50(12), 2461-2466.
- Hotamisligil, G. S., Shargill, N. S., & Spiegelman, B. M. (1993). Adipose expression of tumor necrosis factor-alpha: direct role in obesity-linked insulin resistance. *Science*, 259(5091), 87-91.
- Hotta, K., Funahashi, T., Arita, Y., Takahashi, M., Matsuda, M., Okamoto, Y. (2000). Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol*, 20(6), 1595-1599.

- Hotta, K., Nakamura, M., Nakata, Y., Matsuo, T., Kamohara, S., Kotani, K. (2008). INSIG2 gene rs7566605 polymorphism is associated with severe obesity in Japanese. *J Hum Genet*, 53(9), 857-862.
- Hotta, K., Nakata, Y., Matsuo, T., Kamohara, S., Kotani, K., Komatsu, R. (2008). Variations in the FTO gene are associated with severe obesity in the Japanese. *J Hum Genet*, 53(6), 546-553.
- Houseknecht, K. L., & Portocarrero, C. P. (1998). Leptin and its receptors: regulators of whole-body energy homeostasis. *Domest Anim Endocrinol*, 15(6), 457-475.
- Howard, B. V., Knowler, W. C., Vasquez, B., Kennedy, A. L., Pettitt, D. J., & Bennett, P. H. (1984). Plasma and lipoprotein cholesterol and triglyceride in the Pima Indian population. Comparison of diabetics and nondiabetics. *Arteriosclerosis*, 4(5), 462-471.
- Hu, C., Zhang, R., Wang, C., Wang, J., Ma, X., Lu, J. (2009). PPARG, KCNJ11, CDKAL1, CDKN2A-CDKN2B, IDE-KIF11-HHEX, IGF2BP2 and SLC30A8 are associated with type 2 diabetes in a Chinese population. *PLoS One*, 4(10), e7643.
- Hu, E., Liang, P., & Spiegelman, B. M. (1996). AdipoQ is a novel adipose-specific gene dysregulated in obesity. *J Biol Chem*, 271(18), 10697-10703.
- Hu, F. B., Manson, J. E., Stampfer, M. J., Colditz, G., Liu, S., Solomon, C. G. (2001). Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *N Engl J Med*, 345(11), 790-797.
- Hubacek, J. A., Bohuslavova, R., Kuthanova, L., Kubinova, R., Peasey, A., Pikhart, H. (2008). The FTO gene and obesity in a large Eastern European population sample: the HAPIEE study. *Obesity (Silver Spring)*, 16(12), 2764-2766.
- Hubacek, J. A., Kuthanova, L., Bohuslavova, R., Adamkova, V., Lanska, V., Meitinger, T. (2010). INSIG2 promoter variant, obesity markers and lipid parameters - No association in a large Slavonic Caucasian population sample. *Folia Biol (Praha)*, 56(3), 131-134.
- Hug, C., & Lodish, H. F. (2005). The role of the adipocyte hormone adiponectin in cardiovascular disease. *Curr Opin Pharmacol*, 5(2), 129-134.
- Hunt, S. C., Stone, S., Xin, Y., Scherer, C. A., Magness, C. L., Iadonato, S. P. (2008). Association of the FTO gene with BMI. *Obesity (Silver Spring)*, 16(4), 902-904.
- Huszar, D., Lynch, C. A., Fairchild-Huntress, V., Dunmore, J. H., Fang, Q., Berkemeier, L. R. (1997). Targeted disruption of the melanocortin-4 receptor results in obesity in mice. *Cell*, 88(1), 131-141.
- Iacobellis, G., Ribaldo, M. C., Zappaterreno, A., Iannucci, C. V., & Leonetti, F. (2005). Prevalence of uncomplicated obesity in an Italian obese population. *Obes Res*, 13(6), 1116-1122.

- Ingalls, A. M., Dickie, M. M., & Snell, G. D. (1950). Obese, a new mutation in the house mouse. *J Hered*, 41(12), 317-318.
- Insel, P. A. (1996). Seminars in medicine of the Beth Israel Hospital, Boston. Adrenergic receptors--evolving concepts and clinical implications. *N Engl J Med*, 334(9), 580-585.
- IPH. (2008). (Institute for Public Health). *The Third National Health and Morbidity Survey (NHMS III) 2006, General Findings. Ministry of Health Malaysia, Kuala Lumpur*. Kuala Lumpur: Ministry of Health Malaysia.
- Iqbal, N., Seshadri, P., Stern, L., Loh, J., Kundu, S., Jafar, T. (2005). Serum resistin is not associated with obesity or insulin resistance in humans. *Eur Rev Med Pharmacol Sci*, 9(3), 161-165.
- Ishiyama-Shigemoto, S., Yamada, K., Yuan, X., Ichikawa, F., & Nonaka, K. (1999). Association of polymorphisms in the beta2-adrenergic receptor gene with obesity, hypertriglyceridaemia, and diabetes mellitus. *Diabetologia*, 42(1), 98-101.
- Isse, N., Ogawa, Y., Tamura, N., Masuzaki, H., Mori, K., Okazaki, T. (1995). Structural organization and chromosomal assignment of the human obese gene. *J Biol Chem*, 270(46), 27728-27733.
- Jackson, R. S., Creemers, J. W., Farooqi, I. S., Raffin-Sanson, M. L., Varro, A., Dockray, G. J. (2003). Small-intestinal dysfunction accompanies the complex endocrinopathy of human proprotein convertase 1 deficiency. *J Clin Invest*, 112(10), 1550-1560.
- Jackson, R. S., Creemers, J. W., Ohagi, S., Raffin-Sanson, M. L., Sanders, L., Montague, C. T. (1997). Obesity and impaired prohormone processing associated with mutations in the human prohormone convertase 1 gene. *Nat Genet*, 16(3), 303-306.
- Jacobsson, J. A., Danielsson, P., Svensson, V., Klovins, J., Gyllensten, U., Marcus, C. (2008). Major gender difference in association of FTO gene variant among severely obese children with obesity and obesity related phenotypes. *Biochem Biophys Res Commun*, 368(3), 476-482.
- Jacobsson, J. A., Riserus, U., Axelsson, T., Lannfelt, L., Schioth, H. B., & Fredriksson, R. (2009). The common FTO variant rs9939609 is not associated with BMI in a longitudinal study on a cohort of Swedish men born 1920-1924. *BMC Med Genet*, 10, 131.
- Jalba, M. S., Rhoads, G. G., & Demissie, K. (2008). Association of codon 16 and codon 27 beta 2-adrenergic receptor gene polymorphisms with obesity: a meta-analysis. *Obesity (Silver Spring)*, 16(9), 2096-2106.
- James, W. P. (2005). Assessing obesity: are ethnic differences in body mass index and waist classification criteria justified? *Obes Rev*, 6(3), 179-181.



- Janowska, J., Zahorska-Markiewicz, B., & Olszanecka-Glinianowicz, M. (2006). Relationship between serum resistin concentration and proinflammatory cytokines in obese women with impaired and normal glucose tolerance. *Metabolism*, 55(11), 1495-1499.
- Jee, S. H., Sull, J. W., Lee, J. E., Shin, C., Park, J., Kimm, H. (2010). Adiponectin concentrations: a genome-wide association study. *Am J Hum Genet*, 87(4), 545-552.
- Jequier, E. (2002). Leptin signaling, adiposity, and energy balance. *Ann N Y Acad Sci*, 967, 379-388.
- Jequier, E., & Tappy, L. (1999). Regulation of body weight in humans. *Physiol Rev*, 79(2), 451-480.
- Jia, G., Yang, C. G., Yang, S., Jian, X., Yi, C., Zhou, Z. (2008). Oxidative demethylation of 3-methylthymine and 3-methyluracil in single-stranded DNA and RNA by mouse and human FTO. *FEBS Lett*, 582(23-24), 3313-3319.
- Jiang, Y., Wilk, J. B., Borecki, I., Williamson, S., DeStefano, A. L., Xu, G. (2004). Common variants in the 5' region of the leptin gene are associated with body mass index in men from the National Heart, Lung, and Blood Institute Family Heart Study. *Am J Hum Genet*, 75(2), 220-230.
- Jin, S. G., Kadam, S., & Pfeifer, G. P. (2010). Examination of the specificity of DNA methylation profiling techniques towards 5-methylcytosine and 5-hydroxymethylcytosine. *Nucleic Acids Res*, 38(11), e125.
- Johnson, L., van Jaarsveld, C. H., Emmett, P. M., Rogers, I. S., Ness, A. R., Hattersley, A. T. (2009). Dietary energy density affects fat mass in early adolescence and is not modified by FTO variants. *PLoS One*, 4(3), e4594.
- Jones, P. A. (2005). Overview of cancer epigenetics. *Semin Hematol*, 42(3 Suppl 2), S3-8.
- Jorde, L. B. (1995). Linkage disequilibrium as a gene-mapping tool. *Am J Hum Genet*, 56(1), 11-14.
- Julius, S., Valentini, M., & Palatini, P. (2000). Overweight and hypertension : a 2-way street? *Hypertension*, 35(3), 807-813.
- Kao WHL, C. J., Shuldiner AR et al. (1999). Association of obesity and genetic variants of the peroxisome proliferator activated receptor-gamma2 , the beta-2-adrenergic receptor, and the leptin receptor in Africa Americans. The Atherosclerosis Risk in Communities (ARIC) Study. *Diabetes*, 48, A397.
- Kao, W. T., Yen, Y. C., & Lung, F. W. (2008). The effects of beta2 adrenergic receptor gene polymorphism in lipid profiles. *Lipids Health Dis*, 7, 20.
- Kaplan, N. L., Hill, W. G., & Weir, B. S. (1995). Likelihood methods for locating disease genes in nonequilibrium populations. *Am J Hum Genet*, 56(1), 18-32.

- Kaput, J. (2008). Nutrigenomics research for personalized nutrition and medicine. *Curr Opin Biotechnol*, 19(2), 110-120.
- Kaput, J., & Rodriguez, R. L. (2004). Nutritional genomics: the next frontier in the postgenomic era. *Physiol Genomics*, 16(2), 166-177.
- Karasawa, S., Daimon, M., Sasaki, S., Toriyama, S., Oizumi, T., Susa, S. (2010). Association of the common fat mass and obesity associated (FTO) gene polymorphism with obesity in a Japanese population. *Endocr J*, 57(4), 293-301.
- Karasawa S., M. D., Sasu M, Sasaki S, Toriyama S, Oizumi T, Susa S, Kameda W, Wada K, Muramatsu M, Fukao A, Kubota I, Kawata S, Kayama and Kato T, & (2009). Association of the Common Fat Mass and Obesity Associated (FTO) Gene Polymorphism with Obesity in a Japanese Population. *Endocr J*, k09E-305.
- Kawamura, T., Egusa, G., Fujikawa, R., & Okubo, M. (2001). Gln27Glu variant of the beta2-adrenergic receptor gene is not associated with obesity and diabetes in Japanese-Americans. *Metabolism*, 50(4), 443-446.
- Kennedy, G. C. (1953). The role of depot fat in the hypothalamic control of food intake in the rat. *Proc R Soc Lond B Biol Sci*, 140(901), 578-596.
- Khoury MJ, B. T., Cohen BH. (1993). Fundamentals of genetic epidemiology (pp. 383): Oxford University Press.
- Kim, K. H., Lee, K., Moon, Y. S., & Sul, H. S. (2001). A cysteine-rich adipose tissue-specific secretory factor inhibits adipocyte differentiation. *J Biol Chem*, 276(14), 11252-11256.
- Kim, S. H., Kim, D. J., Seo, I. A., Min, Y. K., Lee, M. S., Kim, K. W. (2002). Significance of beta2-adrenergic receptor gene polymorphism in obesity and type 2 diabetes mellitus in Korean subjects. *Metabolism*, 51(7), 833-837.
- Kim, W., Shin, D. J., Harihara, S., & Kim, Y. J. (2000). Y chromosomal DNA variation in east Asian populations and its potential for inferring the peopling of Korea. *J Hum Genet*, 45(2), 76-83.
- Kissebah, A. H., Sonnenberg, G. E., Myklebust, J., Goldstein, M., Broman, K., James, R. G. (2000). Quantitative trait loci on chromosomes 3 and 17 influence phenotypes of the metabolic syndrome. *Proc Natl Acad Sci U S A*, 97(26), 14478-14483.
- Klein, S., Burke, L. E., Bray, G. A., Blair, S., Allison, D. B., Pi-Sunyer, X. (2004). Clinical implications of obesity with specific focus on cardiovascular disease: a statement for professionals from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism: endorsed by the American College of Cardiology Foundation. *Circulation*, 110(18), 2952-2967.

- Kloting, N., Schleinitz, D., Ruschke, K., Berndt, J., Fasshauer, M., Tonjes, A. (2008). Inverse relationship between obesity and FTO gene expression in visceral adipose tissue in humans. *Diabetologia*, 51(4), 641-647.
- Knuutila, S., Aalto, Y., Autio, K., Bjorkqvist, A. M., El-Rifai, W., Hemmer, S. (1999). DNA copy number losses in human neoplasms. *Am J Pathol*, 155(3), 683-694.
- Kobilka, B. K., Dixon, R. A., Frielle, T., Dohlman, H. G., Bolanowski, M. A., Sigal, I. S. (1987). cDNA for the human beta 2-adrenergic receptor: a protein with multiple membrane-spanning domains and encoded by a gene whose chromosomal location is shared with that of the receptor for platelet-derived growth factor. *Proc Natl Acad Sci U S A*, 84(1), 46-50.
- Koerner, A., Kratzsch, J., & Kiess, W. (2005). Adipocytokines: leptin--the classical, resistin--the controversial, adiponectin--the promising, and more to come. *Best Pract Res Clin Endocrinol Metab*, 19(4), 525-546.
- Koh Yuan Ni, Y. S.-Y., Lok Lee Tin and Lim Lay Hong. (2009). Genotyping candidate genes related to obesity in UCSI University student cohort. *Journal for The Advancement of Science and Arts*, 1(1), 29-42.
- Krentz, A. J. (2003). Lipoprotein abnormalities and their consequences for patients with type 2 diabetes. *Diabetes Obes Metab*, 5 Suppl 1, S19-27.
- Kring, S. I., Holst, C., Zimmermann, E., Jess, T., Berentzen, T., Toubro, S. (2008). FTO gene associated fatness in relation to body fat distribution and metabolic traits throughout a broad range of fatness. *PLoS One*, 3(8), e2958.
- Krude, H., Biebermann, H., Schnabel, D., Tansek, M. Z., Theunissen, P., Mullis, P. E. (2003). Obesity due to proopiomelanocortin deficiency: three new cases and treatment trials with thyroid hormone and ACTH4-10. *J Clin Endocrinol Metab*, 88(10), 4633-4640.
- Kruglyak, L., & Nickerson, D. A. (2001). Variation is the spice of life. *Nat Genet*, 27(3), 234-236.
- Kumar, J., Sunkishala, R. R., Karthikeyan, G., & Sengupta, S. (2007). The common genetic variant upstream of INSIG2 gene is not associated with obesity in Indian population. *Clin Genet*, 71(5), 415-418.
- Lafontan, M., & Berlan, M. (1993). Fat cell adrenergic receptors and the control of white and brown fat cell function. *J Lipid Res*, 34(7), 1057-1091.
- Lange, L. A., Norris, J. M., Langefeld, C. D., Nicklas, B. J., Wagenknecht, L. E., Saad, M. F. (2005). Association of adipose tissue deposition and beta-2 adrenergic receptor variants: the IRAS family study. *Int J Obes (Lond)*, 29(5), 449-457.
- Lappalainen, T. J., Tolppanen, A. M., Kolehmainen, M., Schwab, U., Lindstrom, J., Tuomilehto, J. (2009). The common variant in the FTO gene did not modify the effect of lifestyle changes on body weight: the Finnish Diabetes Prevention Study. *Obesity (Silver Spring)*, 17(4), 832-836.

- Larder, R., Cheung, M. K., Tung, Y. C., Yeo, G. S., & Coll, A. P. (2011). Where to go with FTO? *Trends Endocrinol Metab*, 22(2), 53-59.
- Large, V., Hellstrom, L., Reynisdottir, S., Lonnqvist, F., Eriksson, P., Lannfelt, L. (1997). Human beta-2 adrenoceptor gene polymorphisms are highly frequent in obesity and associate with altered adipocyte beta-2 adrenoceptor function. *J Clin Invest*, 100(12), 3005-3013.
- Lasseka WD, S. J. C. G. (2008). Waist-hip ratio and cognitive ability: is gluteofemoral fat a privileged store of neurodevelopmental resources? *Evolution and Human Behavior*, 29, 26-34.
- Lau, C. H., & Muniandy, S. (2011). Novel adiponectin-resistin (AR) and insulin resistance (IRAR) indexes are useful integrated diagnostic biomarkers for insulin resistance, type 2 diabetes and metabolic syndrome: a case control study. *Cardiovasc Diabetol*, 10(1), 8.
- Le Stunff, C., Le Bihan, C., Schork, N. J., & Bougneres, P. (2000). A common promoter variant of the leptin gene is associated with changes in the relationship between serum leptin and fat mass in obese girls. *Diabetes*, 49(12), 2196-2200.
- Lebovitz, H. E. (1999). Type 2 diabetes: an overview. *Clin Chem*, 45(8 Pt 2), 1339-1345.
- Lee, G. H., Proenca, R., Montez, J. M., Carroll, K. M., Darvishzadeh, J. G., Lee, J. I. (1996). Abnormal splicing of the leptin receptor in diabetic mice. *Nature*, 379(6566), 632-635.
- Lee, J. H., Chan, J. L., Yiannakouris, N., Kontogianni, M., Estrada, E., Seip, R. (2003). Circulating resistin levels are not associated with obesity or insulin resistance in humans and are not regulated by fasting or leptin administration: cross-sectional and interventional studies in normal, insulin-resistant, and diabetic subjects. *J Clin Endocrinol Metab*, 88(10), 4848-4856.
- Lee, S. A., Kallianpur, A., Xiang, Y. B., Wen, W., Cai, Q., Liu, D. (2007). Intra-individual variation of plasma adipokine levels and utility of single measurement of these biomarkers in population-based studies. *Cancer Epidemiol Biomarkers Prev*, 16(11), 2464-2470.
- Lee, Y. S. (2009). The role of leptin-melanocortin system and human weight regulation: lessons from experiments of nature. *Ann Acad Med Singapore*, 38(1), 34-11.
- Legry, V., Cottel, D., Ferrieres, J., Arveiler, D., Andrieux, N., Bingham, A. (2009). Effect of an FTO polymorphism on fat mass, obesity, and type 2 diabetes mellitus in the French MONICA Study. *Metabolism*, 58(7), 971-975.
- Lewontin, R. C. (1995). The detection of linkage disequilibrium in molecular sequence data. *Genetics*, 140(1), 377-388.

- Li, H., Kilpelainen, T. O., Liu, C., Zhu, J., Liu, Y., Hu, C. (2011). Association of genetic variation in FTO with risk of obesity and type 2 diabetes with data from 96,551 East and South Asians. *Diabetologia*.
- Li, H., Wu, Y., Loos, R. J., Hu, F. B., Liu, Y., Wang, J. (2008). Variants in the fat mass- and obesity-associated (FTO) gene are not associated with obesity in a Chinese Han population. *Diabetes*, 57(1), 264-268.
- Li, J., Takaishi, K., Cook, W., McCorkle, S. K., & Unger, R. H. (2003). Insig-1 "brakes" lipogenesis in adipocytes and inhibits differentiation of preadipocytes. *Proc Natl Acad Sci U S A*, 100(16), 9476-9481.
- Li, X., Song, F., Jiang, H., Zhang, M., Lin, J., Bao, W. (2010). A genetic variation in the fat mass- and obesity-associated gene is associated with obesity and newly diagnosed type 2 diabetes in a Chinese population. *Diabetes Metab Res Rev*, 26(2), 128-132.
- Liljedahl, U., Fredriksson, M., Dahlgren, A., & Syvanen, A. C. (2004). Detecting imbalanced expression of SNP alleles by minisequencing on microarrays. *BMC Biotechnol*, 4, 24.
- Lillioja, S., Mott, D. M., Spraul, M., Ferraro, R., Foley, J. E., Ravussin, E. (1993). Insulin resistance and insulin secretory dysfunction as precursors of non-insulin-dependent diabetes mellitus. Prospective studies of Pima Indians. *N Engl J Med*, 329(27), 1988-1992.
- Lima, J. J., Feng, H., Duckworth, L., Wang, J., Sylvester, J. E., Kissoon, N. (2007). Association analyses of adrenergic receptor polymorphisms with obesity and metabolic alterations. *Metabolism*, 56(6), 757-765.
- Lin, R. C. Y., Ericsson, J. O., Benjafeld, A. V., & Morris, B. J. (2001). Association of beta2-adrenoceptor Gln27Glu variant with body weight but not hypertension. *Am J Hypertens*, 14(12), 1201-1204.
- Lindsay, R. S., Funahashi, T., Krakoff, J., Matsuzawa, Y., Tanaka, S., Kobes, S. (2003). Genome-wide linkage analysis of serum adiponectin in the Pima Indian population. *Diabetes*, 52(9), 2419-2425.
- Ling, H., Waterworth, D. M., Stirnadel, H. A., Pollin, T. I., Barter, P. J., Kesaniemi, Y. A. (2009). Genome-wide linkage and association analyses to identify genes influencing adiponectin levels: the GEMS Study. *Obesity (Silver Spring)*, 17(4), 737-744.
- Litonjua, A. A., Gong, L., Duan, Q. L., Shin, J., Moore, M. J., Weiss, S. T. (2010). Very important pharmacogene summary ADRB2. *Pharmacogenet Genomics*, 20(1), 64-69.
- Liu, G., Zhu, H., Dong, Y., Podolsky, R. H., Treiber, F. A., & Snieder, H. (2011). Influence of common variants in FTO and near INSIG2 and MC4R on growth curves for adiposity in African- and European-American youth. *Eur J Epidemiol*, 26(6), 463-473.

- Liu, X., Li, Y., Wang, L., Zhao, Q., Lu, X., Huang, J. (2008). The INSIG1 gene, not the INSIG2 gene, associated with coronary heart disease: tagSNPs and haplotype-based association study. The Beijing Atherosclerosis Study. *Thromb Haemost*, 100(5), 886-892.
- Liu, Y., Liu, Z., Song, Y., Zhou, D., Zhang, D., Zhao, T. (2010). Meta-analysis added power to identify variants in FTO associated with type 2 diabetes and obesity in the Asian population. *Obesity (Silver Spring)*, 18(8), 1619-1624.
- Liu, Y. J., Liu, X. G., Wang, L., Dina, C., Yan, H., Liu, J. F. (2008). Genome-wide association scans identified CTNBL1 as a novel gene for obesity. *Hum Mol Genet*, 17(12), 1803-1813.
- Loktionov, A. (2003). Common gene polymorphisms and nutrition: emerging links with pathogenesis of multifactorial chronic diseases (review). *J Nutr Biochem*, 14(8), 426-451.
- Lonnqvist, F., Arner, P., Nordfors, L., & Schalling, M. (1995). Overexpression of the obese (ob) gene in adipose tissue of human obese subjects. *Nat Med*, 1(9), 950-953.
- Loos, R. J. (2009). Recent progress in the genetics of common obesity. *Br J Clin Pharmacol*, 68(6), 811-829.
- Loos, R. J. (2011). The genetic epidemiology of melanocortin 4 receptor variants. *Eur J Pharmacol*, 660(1), 156-164.
- Loos, R. J., Barroso, I., O'Rahilly, S., & Wareham, N. J. (2007). Comment on "A common genetic variant is associated with adult and childhood obesity". *Science*, 315(5809), 187; author reply 187.
- Loos, R. J., & Bouchard, C. (2003). Obesity--is it a genetic disorder? *J Intern Med*, 254(5), 401-425.
- Loos, R. J., & Bouchard, C. (2008). FTO: the first gene contributing to common forms of human obesity. *Obes Rev*, 9(3), 246-250.
- Loos, R. J., Lindgren, C. M., Li, S., Wheeler, E., Zhao, J. H., Prokopenko, I. (2008). Common variants near MC4R are associated with fat mass, weight and risk of obesity. *Nat Genet*, 40(6), 768-775.
- Lopez-Bermejo, A., Petry, C. J., Diaz, M., Sebastiani, G., de Zegher, F., Dunger, D. B. (2008). The association between the FTO gene and fat mass in humans develops by the postnatal age of two weeks. *J Clin Endocrinol Metab*, 93(4), 1501-1505.
- Luo, J. D., Zhang, G. S., & Chen, M. S. (2005). Leptin and cardiovascular diseases. *Timely Top Med Cardiovasc Dis*, 9, E34.
- Lyon, C. J., Law, R. E., & Hsueh, W. A. (2003). Minireview: adiposity, inflammation, and atherogenesis. *Endocrinology*, 144(6), 2195-2200.

- Lyon, H. N., Emilsson, V., Hinney, A., Heid, I. M., Lasky-Su, J., Zhu, X. (2007). The association of a SNP upstream of *INSIG2* with body mass index is reproduced in several but not all cohorts. *PLoS Genet*, 3(4), e61.
- Maccio, A., Madeddu, C., Gramignano, G., Mulas, C., Floris, C., Massa, D. (2010). Correlation of body mass index and leptin with tumor size and stage of disease in hormone-dependent postmenopausal breast cancer: preliminary results and therapeutic implications. *J Mol Med (Berl)*, 88(7), 677-686.
- Maeda, K., Okubo, K., Shimomura, I., Funahashi, T., Matsuzawa, Y., & Matsubara, K. (1996). cDNA cloning and expression of a novel adipose specific collagen-like factor, apM1 (AdiPose Most abundant Gene transcript 1). *Biochem Biophys Res Commun*, 221(2), 286-289.
- Maes, H. H., Neale, M. C., & Eaves, L. J. (1997). Genetic and environmental factors in relative body weight and human adiposity. *Behav Genet*, 27(4), 325-351.
- Maffei, M., Halaas, J., Ravussin, E., Pratley, R. E., Lee, G. H., Zhang, Y. (1995). Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects. *Nat Med*, 1(11), 1155-1161.
- Malathi, M. R. R. a. R. (2007). Possible correlation of leptin with body fat distribution and adiposity: Evaluation of serum leptin in South Indian population. *Reproductive Medicine and Biology*, 6, 117-125.
- Markison S, F. A. (2006). Targeting melanocortin receptors for the treatment of obesity. *Drug Discovery Today: Therapeutic Strategies* 3(4), 569-576.
- Markison, S., & Foster, A. (2006). Targeting melanocortin receptors for the treatment of obesity. *Drug Discovery Today: Therapeutic Strategies* 3(4), 569-576.
- Martinez, J. A. (2000). Body-weight regulation: causes of obesity. *Proc Nutr Soc*, 59(3), 337-345.
- Martinez, J. A., & Fruhbeck, G. (1996). Regulation of energy balance and adiposity: a model with new approaches. *Rev Esp Fisiol*, 52(4), 255-258.
- Marville, A. F., Lange, L. A., Qin, L., Adair, L. S., & Mohlke, K. L. (2008). Association of *FTO* with obesity-related traits in the Cebu Longitudinal Health and Nutrition Survey (CLHNS) Cohort. *Diabetes*, 57(7), 1987-1991.
- Masuo, K., Katsuya, T., Kawaguchi, H., Fu, Y., Rakugi, H., Ogihara, T. (2006). Beta2-adrenoceptor polymorphisms relate to obesity through blunted leptin-mediated sympathetic activation. *Am J Hypertens*, 19(10), 1084-1091.
- Matsubara, M., Maruoka, S., & Katayose, S. (2002). Decreased plasma adiponectin concentrations in women with dyslipidemia. *J Clin Endocrinol Metab*, 87(6), 2764-2769.
- Matsuzawa, Y., Funahashi, T., Kihara, S., & Shimomura, I. (2004). Adiponectin and metabolic syndrome. *Arterioscler Thromb Vasc Biol*, 24(1), 29-33.

- Mattevi, V. S., Zembruski, V. M., & Hutz, M. H. (2004). A resistin gene polymorphism is associated with body mass index in women. *Hum Genet*, *115*(3), 208-212.
- Meilleur, K. G., Doumatey, A., Huang, H., Charles, B., Chen, G., Zhou, J. (2010). Circulating Adiponectin Is Associated with Obesity and Serum Lipids in West Africans. *J Clin Endocrinol Metab*.
- Menzaghi, C., Coco, A., Salvemini, L., Thompson, R., De Cosmo, S., Doria, A. (2006). Heritability of serum resistin and its genetic correlation with insulin resistance-related features in nondiabetic Caucasians. *J Clin Endocrinol Metab*, *91*(7), 2792-2795.
- Meuwissen, T. H., & Goddard, M. E. (2000). Fine mapping of quantitative trait loci using linkage disequilibria with closely linked marker loci. *Genetics*, *155*(1), 421-430.
- Meyre, D., Delplanque, J., Chevre, J. C., Lecoecur, C., Lobbens, S., Gallina, S. (2009). Genome-wide association study for early-onset and morbid adult obesity identifies three new risk loci in European populations. *Nat Genet*, *41*(2), 157-159.
- Montague, C. T., Farooqi, I. S., Whitehead, J. P., Soos, M. A., Rau, H., Wareham, N. J. (1997). Congenital leptin deficiency is associated with severe early-onset obesity in humans. *Nature*, *387*(6636), 903-908.
- Montague, C. T., & O'Rahilly, S. (2000). The perils of portliness: causes and consequences of visceral adiposity. *Diabetes*, *49*(6), 883-888.
- Mori, Y., Kim-Motoyama, H., Ito, Y., Katakura, T., Yasuda, K., Ishiyama-Shigemoto, S. (1999). The Gln27Glu beta2-adrenergic receptor variant is associated with obesity due to subcutaneous fat accumulation in Japanese men. *Biochem Biophys Res Commun*, *258*(1), 138-140.
- Morris, R. W., & Kaplan, N. L. (2002). On the advantage of haplotype analysis in the presence of multiple disease susceptibility alleles. *Genet Epidemiol*, *23*(3), 221-233.
- Moy, F. M., & Bulgiba, A. (2010). The modified NCEP ATP III criteria maybe better than the IDF criteria in diagnosing Metabolic Syndrome among Malays in Kuala Lumpur. *BMC Public Health*, *10*, 678.
- Muller, T. D., Hinney, A., Scherag, A., Nguyen, T. T., Schreiner, F., Schafer, H. (2008). 'Fat mass and obesity associated' gene (FTO): no significant association of variant rs9939609 with weight loss in a lifestyle intervention and lipid metabolism markers in German obese children and adolescents. *BMC Med Genet*, *9*, 85.
- Muoio, D. M., & Lynis Dohm, G. (2002). Peripheral metabolic actions of leptin. *Best Pract Res Clin Endocrinol Metab*, *16*(4), 653-666.



- Mutch, D. M., & Clement, K. (2006). Unraveling the genetics of human obesity. *PLoS Genet*, 2(12), e188.
- Nakano, Y., Tobe, T., Choi-Miura, N. H., Mazda, T., & Tomita, M. (1996). Isolation and characterization of GBP28, a novel gelatin-binding protein purified from human plasma. *J Biochem*, 120(4), 803-812.
- Narkiewicz, K. (2006). Obesity and hypertension--the issue is more complex than we thought. *Nephrol Dial Transplant*, 21(2), 264-267.
- Ng, M. C., Park, K. S., Oh, B., Tam, C. H., Cho, Y. M., Shin, H. D. (2008). Implication of genetic variants near TCF7L2, SLC30A8, HHEX, CDKAL1, CDKN2A/B, IGF2BP2, and FTO in type 2 diabetes and obesity in 6,719 Asians. *Diabetes*, 57(8), 2226-2233.
- Ng, M. C., Tam, C. H., So, W. Y., Ho, J. S., Chan, A. W., Lee, H. M. (2010). Implication of genetic variants near NEGR1, SEC16B, TMEM18, ETV5/DGKG, GNPDA2, LIN7C/BDNF, MTCH2, BCDIN3D/FAIM2, SH2B1, FTO, MC4R, and KCTD15 with obesity and type 2 diabetes in 7705 Chinese. *J Clin Endocrinol Metab*, 95(5), 2418-2425.
- Nishizawa, H., Shimomura, I., Kishida, K., Maeda, N., Kuriyama, H., Nagaretani, H. (2002). Androgens decrease plasma adiponectin, an insulin-sensitizing adipocyte-derived protein. *Diabetes*, 51(9), 2734-2741.
- Norata, G. D., Ongari, M., Garlaschelli, K., Raselli, S., Grigore, L., & Catapano, A. L. (2007). Plasma resistin levels correlate with determinants of the metabolic syndrome. *Eur J Endocrinol*, 156(2), 279-284.
- Nowotny, P., Kwon, J. M., & Goate, A. M. (2001). SNP analysis to dissect human traits. *Curr Opin Neurobiol*, 11(5), 637-641.
- Oberkofler, H., Esterbauer, H., Hell, E., Krempler, F., & Patsch, W. (2000). The Gln27Glu polymorphism in the beta2-adrenergic receptor gene is not associated with morbid obesity in Austrian women. *Int J Obes Relat Metab Disord*, 24(3), 388-390.
- Oki, K., Yamane, K., Kamei, N., Asao, T., Awaya, T., & Kohno, N. (2009). The single nucleotide polymorphism upstream of insulin-induced gene 2 (INSIG2) is associated with the prevalence of hypercholesterolaemia, but not with obesity, in Japanese American women. *Br J Nutr*, 101(3), 322-327.
- Okuda, T., Fujioka, Y., Kamide, K., Kawano, Y., Goto, Y., Yoshimasa, Y. (2002). Verification of 525 coding SNPs in 179 hypertension candidate genes in the Japanese population: identification of 159 SNPs in 93 genes. *J Hum Genet*, 47(8), 387-394.
- Omori, S., Tanaka, Y., Takahashi, A., Hirose, H., Kashiwagi, A., Kaku, K. (2008). Association of CDKAL1, IGF2BP2, CDKN2A/B, HHEX, SLC30A8, and KCNJ11 with susceptibility to type 2 diabetes in a Japanese population. *Diabetes*, 57(3), 791-795.

- Orkunoglu-Suer, F. E., Gordish-Dressman, H., Clarkson, P. M., Thompson, P. D., Angelopoulos, T. J., Gordon, P. M. (2008). INSIG2 gene polymorphism is associated with increased subcutaneous fat in women and poor response to resistance training in men. *BMC Med Genet*, 9, 117.
- Otto, T. C., & Lane, M. D. (2005). Adipose development: from stem cell to adipocyte. *Crit Rev Biochem Mol Biol*, 40(4), 229-242.
- Ouchi, N., Kihara, S., Arita, Y., Maeda, K., Kuriyama, H., Okamoto, Y. (1999). Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin. *Circulation*, 100(25), 2473-2476.
- Ozcan, L., Ergin, A. S., Lu, A., Chung, J., Sarkar, S., Nie, D. (2009). Endoplasmic reticulum stress plays a central role in development of leptin resistance. *Cell Metab*, 9(1), 35-51.
- Ozer, A., & Bruick, R. K. (2007). Non-heme dioxygenases: cellular sensors and regulators jelly rolled into one? *Nat Chem Biol*, 3(3), 144-153.
- Paracchini, V., Pedotti, P., & Taioli, E. (2005). Genetics of leptin and obesity: a HuGE review. *Am J Epidemiol*, 162(2), 101-114.
- Park, H. S., Shin, E. S., & Lee, J. E. (2008). Genotypes and haplotypes of beta2-adrenergic receptor and parameters of the metabolic syndrome in Korean adolescents. *Metabolism*, 57(8), 1064-1070.
- Park, J. H., Gail, M. H., Weinberg, C. R., Carroll, R. J., Chung, C. C., Wang, Z. (2011). Distribution of allele frequencies and effect sizes and their interrelationships for common genetic susceptibility variants. *Proc Natl Acad Sci U S A*, 108(44), 18026-18031.
- Peeters, A., Beckers, S., Verrijken, A., Roevens, P., Peeters, P., Van Gaal, L. (2008). Variants in the FTO gene are associated with common obesity in the Belgian population. *Mol Genet Metab*, 93(4), 481-484.
- Peng, S., Zhu, Y., Xu, F., Ren, X., Li, X., & Lai, M. (2011). FTO gene polymorphisms and obesity risk: a meta-analysis. *BMC Med*, 9, 71.
- Percik, R., & Stumvoll, M. (2009). Obesity and cancer. *Exp Clin Endocrinol Diabetes*, 117(10), 563-566.
- Perry, G. H., Dominy, N. J., Claw, K. G., Lee, A. S., Fiegler, H., Redon, R. (2007). Diet and the evolution of human amylase gene copy number variation. *Nat Genet*, 39(10), 1256-1260.
- Perusse, L., & Bouchard, C. (2000). Gene-diet interactions in obesity. *Am J Clin Nutr*, 72(5 Suppl), 1285S-1290S.
- Piestrzeniewicz, K., Luczak, K., Komorowski, J., Maciejewski, M., Jankiewicz Wika, J., & Goch, J. H. (2008). Resistin increases with obesity and atherosclerotic risk factors in patients with myocardial infarction. *Metabolism*, 57(4), 488-493.

- Pischon, T., Nothlings, U., & Boeing, H. (2008). Obesity and cancer. *Proc Nutr Soc*, 67(2), 128-145.
- Pizzuti, A., Argiolas, A., Di Paola, R., Baratta, R., Rauseo, A., Bozzali, M. (2002). An ATG repeat in the 3'-untranslated region of the human resistin gene is associated with a decreased risk of insulin resistance. *J Clin Endocrinol Metab*, 87(9), 4403-4406.
- Plomin, R., Owen, M. J., & McGuffin, P. (1994). The genetic basis of complex human behaviors. *Science*, 264(5166), 1733-1739.
- Poh BK, S. M., Tahir A, Siti Haslinda MD, Siti Norazlin N, Norimah AK., & Wan Manan WM, M. K., Zalilah MS, Azmi MY & Fatimah S. (2010). Physical Activity Pattern and Energy Expenditure of Malaysian Adults: Findings from the Malaysian Adult Nutrition Survey (MANS). *Mal J Nutr*, 16(1), 13-37.
- Poirier, P., Giles, T. D., Bray, G. A., Hong, Y., Stern, J. S., Pi-Sunyer, F. X. (2006). Obesity and cardiovascular disease: pathophysiology, evaluation, and effect of weight loss: an update of the 1997 American Heart Association Scientific Statement on Obesity and Heart Disease from the Obesity Committee of the Council on Nutrition, Physical Activity, and Metabolism. *Circulation*, 113(6), 898-918.
- Portoles, O., Sorli, J. V., Frances, F., Coltell, O., Gonzalez, J. I., Saiz, C. (2006). Effect of genetic variation in the leptin gene promoter and the leptin receptor gene on obesity risk in a population-based case-control study in Spain. *Eur J Epidemiol*, 21(8), 605-612.
- Postma, D. S., Bleeker, E. R., Amelung, P. J., Holroyd, K. J., Xu, J., Panhuysen, C. I. (1995). Genetic susceptibility to asthma--bronchial hyperresponsiveness coinherited with a major gene for atopy. *N Engl J Med*, 333(14), 894-900.
- Price, R. A., Li, W. D., & Zhao, H. (2008). FTO gene SNPs associated with extreme obesity in cases, controls and extremely discordant sister pairs. *BMC Med Genet*, 9, 4.
- Prieto-Hontoria, P. L., Perez-Matute, P., Fernandez-Galilea, M., Bustos, M., Martinez, J. A., & Moreno-Aliaga, M. J. (2011). Role of obesity-associated dysfunctional adipose tissue in cancer: a molecular nutrition approach. *Biochim Biophys Acta*, 1807(6), 664-678.
- Pritchard, J. K. (2001). Are rare variants responsible for susceptibility to complex diseases? *Am J Hum Genet*, 69(1), 124-137.
- Pritchard, J. K., & Przeworski, M. (2001). Linkage disequilibrium in humans: models and data. *Am J Hum Genet*, 69(1), 1-14.
- Przeworski, M., & Wall, J. D. (2001). Why is there so little intragenic linkage disequilibrium in humans? *Genet Res*, 77(2), 143-151.

- Radhakrishnan, A., Ikeda, Y., Kwon, H. J., Brown, M. S., & Goldstein, J. L. (2007). Sterol-regulated transport of SREBPs from endoplasmic reticulum to Golgi: oxysterols block transport by binding to Insig. *Proc Natl Acad Sci U S A*, *104*(16), 6511-6518.
- Ramachandran, A., & Snehalatha, C. (2010). Rising burden of obesity in Asia. *J Obes*, *2010*.
- Rampersaud, E., Mitchell, B. D., Pollin, T. I., Fu, M., Shen, H., O'Connell, J. R. (2008). Physical activity and the association of common FTO gene variants with body mass index and obesity. *Arch Intern Med*, *168*(16), 1791-1797.
- Rankinen, T., Zuberi, A., Chagnon, Y. C., Weisnagel, S. J., Argyropoulos, G., Walts, B. (2006). The human obesity gene map: the 2005 update. *Obesity (Silver Spring)*, *14*(4), 529-644.
- Reich, D. E., Cargill, M., Bolk, S., Ireland, J., Sabeti, P. C., Richter, D. J. (2001). Linkage disequilibrium in the human genome. *Nature*, *411*(6834), 199-204.
- Reilly, M. P., Lehrke, M., Wolfe, M. L., Rohatgi, A., Lazar, M. A., & Rader, D. J. (2005). Resistin is an inflammatory marker of atherosclerosis in humans. *Circulation*, *111*(7), 932-939.
- Reizes, O., Benoit, S. C., & Clegg, D. J. (2008). The role of syndecans in the regulation of body weight and synaptic plasticity. *Int J Biochem Cell Biol*, *40*(1), 28-45.
- Reizes, O., Lincecum, J., Wang, Z., Goldberger, O., Huang, L., Kaksonen, M. (2001). Transgenic expression of syndecan-1 uncovers a physiological control of feeding behavior by syndecan-3. *Cell*, *106*(1), 105-116.
- Reynisdottir, S., Langin, D., Carlstrom, K., Holm, C., Rossner, S., & Arner, P. (1995). Effects of weight reduction on the regulation of lipolysis in adipocytes of women with upper-body obesity. *Clin Sci (Lond)*, *89*(4), 421-429.
- Reynisdottir, S., Wahrenberg, H., Carlstrom, K., Rossner, S., & Arner, P. (1994). Catecholamine resistance in fat cells of women with upper-body obesity due to decreased expression of beta 2-adrenoceptors. *Diabetologia*, *37*(4), 428-435.
- Rimm, E. B., Stampfer, M. J., Giovannucci, E., Ascherio, A., Spiegelman, D., Colditz, G. A. (1995). Body size and fat distribution as predictors of coronary heart disease among middle-aged and older US men. *Am J Epidemiol*, *141*(12), 1117-1127.
- Rong LUO, X.-p. L., Yan ZHAO. (2007). Serum resistin and adiponectin concentrations in patients with overweight and obesity. *Journal of Medical Colleges of PLA*, *22*(3), 160-164.
- Rosenbaum, M., Leibel, R. L., & Hirsch, J. (1997). Obesity. *N Engl J Med*, *337*(6), 396-407.

- Rosenbaum, M., Nicolson, M., Hirsch, J., Heymsfield, S. B., Gallagher, D., Chu, F. (1996). Effects of gender, body composition, and menopause on plasma concentrations of leptin. *J Clin Endocrinol Metab*, 81(9), 3424-3427.
- Rosmond, R., Chagnon, M., Bouchard, C., & Bjorntorp, P. (2001). A missense mutation in the human melanocortin-4 receptor gene in relation to abdominal obesity and salivary cortisol. *Diabetologia*, 44(10), 1335-1338.
- Ross, R. (1999). Atherosclerosis--an inflammatory disease. *N Engl J Med*, 340(2), 115-126.
- Roskopf, D., Bornhorst, A., Rimmbach, C., Schwahn, C., Kayser, A., Kruger, A. (2007). Comment on "A common genetic variant is associated with adult and childhood obesity". *Science*, 315(5809), 187; author reply 187.
- Rutanen, J., Pihlajamaki, J., Karhapaa, P., Vauhkonen, I., Kuusisto, J., Moilanen Mykkanen, L. (2004). The Val103Ile polymorphism of melanocortin-4 receptor regulates energy expenditure and weight gain. *Obes Res*, 12(7), 1060-1066.
- Salisbury, B. A., Pungliya, M., Choi, J. Y., Jiang, R., Sun, X. J., & Stephens, J. C. (2003). SNP and haplotype variation in the human genome. *Mutat Res*, 526(1-2), 53-61.
- Samad, F., & Loskutoff, D. J. (1996). Tissue distribution and regulation of plasminogen activator inhibitor-1 in obese mice. *Mol Med*, 2(5), 568-582.
- Samad, F., Yamamoto, K., Pandey, M., & Loskutoff, D. J. (1997). Elevated expression of transforming growth factor-beta in adipose tissue from obese mice. *Mol Med*, 3(1), 37-48.
- Sanghera, D. K., Ortega, L., Han, S., Singh, J., Ralhan, S. K., Wander, G. S. (2008). Impact of nine common type 2 diabetes risk polymorphisms in Asian Indian Sikhs: PPARG2 (Pro12Ala), IGF2BP2, TCF7L2 and FTO variants confer a significant risk. *BMC Med Genet*, 9, 59.
- Santiago Rodriguez , T. R. G. a. L. N. (2009). Hardy-Weinberg Equilibrium Testing of Biological Ascertainment for Mendelian Randomization Studies. *American Journal of Epidemiology*.
- Saris, W. H., Blair, S. N., van Baak, M. A., Eaton, S. B., Davies, P. S., Di Pietro, L. (2003). How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st Stock Conference and consensus statement. *Obes Rev*, 4(2), 101-114.
- Schaffler, A., Buchler, C., Muller-Ladner, U., Herfarth, H., Ehling, A., Paul, G. (2004). Identification of variables influencing resistin serum levels in patients with type 1 and type 2 diabetes mellitus. *Horm Metab Res*, 36(10), 702-707.
- Scherer, P. E., Williams, S., Fogliano, M., Baldini, G., & Lodish, H. F. (1995). A novel serum protein similar to C1q, produced exclusively in adipocytes. *J Biol Chem*, 270(45), 26746-26749.

- Schrauwen, P., van Marken Lichtenbelt, W. D., Saris, W. H., & Westerterp, K. R. (1997). Changes in fat oxidation in response to a high-fat diet. *Am J Clin Nutr*, 66(2), 276-282.
- Schuring, A. N., Lutz, F., Tuttelmann, F., Gromoll, J., Kiesel, L., & Gotte, M. (2009). Role of syndecan-3 polymorphisms in obesity and female hyperandrogenism. *J Mol Med*, 87(12), 1241-1250.
- Schutz, Y. (1995). Abnormalities of fuel utilization as predisposing to the development of obesity in humans. *Obes Res*, 3 Suppl 2, 173S-178S.
- Scott, J. (1996). New chapter for the fat controller. *Nature*, 379(6561), 113-114.
- Scuteri, A., Sanna, S., Chen, W. M., Uda, M., Albai, G., Strait, J. (2007). Genome-wide association scan shows genetic variants in the FTO gene are associated with obesity-related traits. *PLoS Genet*, 3(7), e115.
- Sentinelli, F., Romeo, S., Arca, M., Filippi, E., Leonetti, F., Banchieri, M. (2002). Human resistin gene, obesity, and type 2 diabetes: mutation analysis and population study. *Diabetes*, 51(3), 860-862.
- Serre, D., & Hudson, T. J. (2006). Resources for genetic variation studies. *Annu Rev Genomics Hum Genet*, 7, 443-457.
- Sever, N., Lee, P. C., Song, B. L., Rawson, R. B., & Debose-Boyd, R. A. (2004). Isolation of mutant cells lacking Insig-1 through selection with SR-12813, an agent that stimulates degradation of 3-hydroxy-3-methylglutaryl-coenzyme A reductase. *J Biol Chem*, 279(41), 43136-43147.
- Sever, N., Yang, T., Brown, M. S., Goldstein, J. L., & DeBose-Boyd, R. A. (2003). Accelerated degradation of HMG CoA reductase mediated by binding of insig-1 to its sterol-sensing domain. *Mol Cell*, 11(1), 25-33.
- Sewter, C., Berger, D., Considine, R. V., Medina, G., Rochford, J., Ciaraldi, T. (2002). Human obesity and type 2 diabetes are associated with alterations in SREBP1 isoform expression that are reproduced ex vivo by tumor necrosis factor-alpha. *Diabetes*, 51(4), 1035-1041.
- Sha, B. Y., Yang, T. L., Zhao, L. J., Chen, X. D., Guo, Y., Chen, Y. (2009). Genome-wide association study suggested copy number variation may be associated with body mass index in the Chinese population. *J Hum Genet*, 54(4), 199-202.
- Shi, Y. Y., & He, L. (2005). SHEsis, a powerful software platform for analyses of linkage disequilibrium, haplotype construction, and genetic association at polymorphism loci. *Cell Res*, 15(2), 97-98.
- Siegel, E. M., Ulrich, C. M., Poole, E. M., Holmes, R. S., Jacobsen, P. B., & Shibata, D. (2010). The effects of obesity and obesity-related conditions on colorectal cancer prognosis. *Cancer Control*, 17(1), 52-57.

- Smith, A. J., Cooper, J. A., Li, L. K., & Humphries, S. E. (2007). INSIG2 gene polymorphism is not associated with obesity in Caucasian, Afro-Caribbean and Indian subjects. *Int J Obes (Lond)*, *31*(11), 1753-1755.
- Sonestedt, E., Roos, C., Gullberg, B., Ericson, U., Wirfalt, E., & Orho-Melander, M. (2009). Fat and carbohydrate intake modify the association between genetic variation in the FTO genotype and obesity. *Am J Clin Nutr*, *90*(5), 1418-1425.
- Song, Y., You, N. C., Hsu, Y. H., Howard, B. V., Langer, R. D., Manson, J. E. (2008). FTO polymorphisms are associated with obesity but not diabetes risk in postmenopausal women. *Obesity (Silver Spring)*, *16*(11), 2472-2480.
- Speakman, J. R., Rance, K. A., & Johnstone, A. M. (2008). Polymorphisms of the FTO gene are associated with variation in energy intake, but not energy expenditure. *Obesity (Silver Spring)*, *16*(8), 1961-1965.
- Speliotes, E. K., Willer, C. J., Berndt, S. I., Monda, K. L., Thorleifsson, G., Jackson, A. U. (2010). Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet*, *42*(11), 937-948.
- Spiegelman, B. M., & Flier, J. S. (1996). Adipogenesis and obesity: rounding out the big picture. *Cell*, *87*(3), 377-389.
- Stefansson, H., Helgason, A., Thorleifsson, G., Steinthorsdottir, V., Masson, G., Barnard, J. (2005). A common inversion under selection in Europeans. *Nat Genet*, *37*(2), 129-137.
- Stephens, J. C., Schneider, J. A., Tanguay, D. A., Choi, J., Acharya, T., Stanley, S. E. (2001). Haplotype variation and linkage disequilibrium in 313 human genes. *Science*, *293*(5529), 489-493.
- Steppan, C. M., Bailey, S. T., Bhat, S., Brown, E. J., Banerjee, R. R., Wright, C. M. (2001). The hormone resistin links obesity to diabetes. *Nature*, *409*(6818), 307-312.
- Steppan, C. M., Brown, E. J., Wright, C. M., Bhat, S., Banerjee, R. R., Dai, C. Y. (2001). A family of tissue-specific resistin-like molecules. *Proc Natl Acad Sci U S A*, *98*(2), 502-506.
- Steppan, C. M., & Lazar, M. A. (2002). Resistin and obesity-associated insulin resistance. *Trends Endocrinol Metab*, *13*(1), 18-23.
- Stratigopoulos, G., Padilla, S. L., LeDuc, C. A., Watson, E., Hattersley, A. T., McCarthy, M. I. (2008). Regulation of Fto/Ftm gene expression in mice and humans. *Am J Physiol Regul Integr Comp Physiol*, *294*(4), R1185-1196.
- Strosberg, A. D., & Issad, T. (1999). The involvement of leptin in humans revealed by mutations in leptin and leptin receptor genes. *Trends Pharmacol Sci*, *20*(6), 227-230.

- Stumvoll, M., Tschritter, O., Fritsche, A., Staiger, H., Renn, W., Weisser, M. (2002). Association of the T-G polymorphism in adiponectin (exon 2) with obesity and insulin sensitivity: interaction with family history of type 2 diabetes. *Diabetes*, *51*(1), 37-41.
- Stunkard, A. J., Foch, T. T., & Hrubec, Z. (1986). A twin study of human obesity. *JAMA*, *256*(1), 51-54.
- Stunkard, A. J., Harris, J. R., Pedersen, N. L., & McClearn, G. E. (1990). The body-mass index of twins who have been reared apart. *N Engl J Med*, *322*(21), 1483-1487.
- Stunkard, A. J., & Sorensen, T. I. (1993). Obesity and socioeconomic status--a complex relation. *N Engl J Med*, *329*(14), 1036-1037.
- Stunkard, A. J., Sorensen, T. I., Hanis, C., Teasdale, T. W., Chakraborty, R., Schull, W. J. (1986). An adoption study of human obesity. *N Engl J Med*, *314*(4), 193-198.
- Stutzmann, F., Cauchi, S., Durand, E., Calvacanti-Proenca, C., Pigeyre, M., Hartikainen, A. L. (2009). Common genetic variation near MC4R is associated with eating behaviour patterns in European populations. *Int J Obes (Lond)*, *33*(3), 373-378.
- Stutzmann, F., Vatin, V., Cauchi, S., Morandi, A., Jouret, B., Landt, O. (2007). Non-synonymous polymorphisms in melanocortin-4 receptor protect against obesity: the two facets of a Janus obesity gene. *Hum Mol Genet*, *16*(15), 1837-1844.
- Suzana, S., Kee, C., Jamaludin, A., Noor Safiza, M., Khor, G., Jamaiyah, H. (2012). The third national health and morbidity survey: prevalence of obesity, and abdominal obesity among the malaysian elderly population. *Asia Pac J Public Health*, *24*(2), 318-329.
- Syvanen, A. C., Landegren, U., Isaksson, A., Gyllensten, U., & Brookes, A. (1999). First International SNP Meeting at Skokloster, Sweden, August 1998. Enthusiasm mixed with scepticism about single-nucleotide polymorphism markers for dissecting complex disorders. *Eur J Hum Genet*, *7*(1), 98-101.
- Tabara, Y., Osawa, H., Guo, H., Kawamoto, R., Onuma, H., Shimizu, I. (2009). Prognostic significance of FTO genotype in the development of obesity in Japanese: the J-SHIPP study. *Int J Obes (Lond)*, *33*(11), 1243-1248.
- Takeuchi, F., Serizawa, M., Yamamoto, K., Fujisawa, T., Nakashima, E., Ohnaka, K. (2009). Confirmation of multiple risk Loci and genetic impacts by a genome-wide association study of type 2 diabetes in the Japanese population. *Diabetes*, *58*(7), 1690-1699.
- Tan, J. T., Dorajoo, R., Seielstad, M., Sim, X. L., Ong, R. T., Chia, K. S. (2008). FTO variants are associated with obesity in the Chinese and Malay populations in Singapore. *Diabetes*, *57*(10), 2851-2857.



- Tanofsky-Kraff, M., Han, J. C., Anandalingam, K., Shomaker, L. B., Columbo, K. M., Wolkoff, L. E. (2009). The FTO gene rs9939609 obesity-risk allele and loss of control over eating. *Am J Clin Nutr*, 90(6), 1483-1488.
- Templeton, A. R., Boerwinkle, E., & Sing, C. F. (1987). A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping. I. Basic theory and an analysis of alcohol dehydrogenase activity in *Drosophila*. *Genetics*, 117(2), 343-351.
- Teo, Y. Y., Sim, X., Ong, R. T., Tan, A. K., Chen, J., Tantoso, E. (2009). Singapore Genome Variation Project: a haplotype map of three Southeast Asian populations. *Genome Res*, 19(11), 2154-2162.
- Teo, Y. Y., Small, K. S., Fry, A. E., Wu, Y., Kwiatkowski, D. P., & Clark, T. G. (2009). Power consequences of linkage disequilibrium variation between populations. *Genet Epidemiol*, 33(2), 128-135.
- Terwilliger, J. D., & Weiss, K. M. (1998). Linkage disequilibrium mapping of complex disease: fantasy or reality? *Curr Opin Biotechnol*, 9(6), 578-594.
- Teucher, B., Rohrmann, S., & Kaaks, R. (2010). Obesity: focus on all-cause mortality and cancer. *Maturitas*, 65(2), 112-116.
- Thelle, D. S., Shaper, A. G., Whitehead, T. P., Bullock, D. G., Ashby, D., & Patel, I. (1983). Blood lipids in middle-aged British men. *Br Heart J*, 49(3), 205-213.
- Thorleifsson, G., Walters, G. B., Gudbjartsson, D. F., Steinthorsdottir, V., Sulem, P., Helgadóttir, A. (2009). Genome-wide association yields new sequence variants at seven loci that associate with measures of obesity. *Nat Genet*, 41(1), 18-24.
- Tishkoff, S. A., Reed, F. A., Ranciaro, A., Voight, B. F., Babbitt, C. C., Silverman, J. S. (2007). Convergent adaptation of human lactase persistence in Africa and Europe. *Nat Genet*, 39(1), 31-40.
- Tiwari, A. K., Zai, C. C., Meltzer, H. Y., Lieberman, J. A., Muller, D. J., & Kennedy, J. L. (2010). Association study of polymorphisms in insulin induced gene 2 (INSIG2) with antipsychotic-induced weight gain in European and African-American schizophrenia patients. *Hum Psychopharmacol*, 25(3), 253-259.
- Tonjes, A., Zeggini, E., Kovacs, P., Bottcher, Y., Schleinitz, D., Dietrich, K. (2010). Association of FTO variants with BMI and fat mass in the self-contained population of Sorbs in Germany. *Eur J Hum Genet*, 18(1), 104-110.
- Tousoulis, D., Antoniadis, C., & Stefanadis, C. (2007). Assessing inflammatory status in cardiovascular disease. *Heart*, 93(8), 1001-1007.
- Trewick, S. C., Henshaw, T. F., Hausinger, R. P., Lindahl, T., & Sedgwick, B. (2002). Oxidative demethylation by *Escherichia coli* AlkB directly reverts DNA base damage. *Nature*, 419(6903), 174-178.

- Tung, Y. C., Ayuso, E., Shan, X., Bosch, F., O'Rahilly, S., Coll, A. P. (2010). Hypothalamic-specific manipulation of Fto, the ortholog of the human obesity gene FTO, affects food intake in rats. *PLoS One*, 5(1), e8771.
- Ukkola, O., Kunnari, A., & Kesaniemi, Y. A. (2008). Genetic variants at the resistin locus are associated with the plasma resistin concentration and cardiovascular risk factors. *Regul Pept*, 149(1-3), 56-59.
- Ukkola, O., Rankinen, T., Weisnagel, S. J., Sun, G., Perusse, L., Chagnon, Y. C. (2000). Interactions among the alpha2-, beta2-, and beta3-adrenergic receptor genes and obesity-related phenotypes in the Quebec Family Study. *Metabolism*, 49(8), 1063-1070.
- Vaisse, C., Clement, K., Durand, E., Hercberg, S., Guy-Grand, B., & Froguel, P. (2000). Melanocortin-4 receptor mutations are a frequent and heterogeneous cause of morbid obesity. *J Clin Invest*, 106(2), 253-262.
- van Kruijsdijk, R. C., van der Wall, E., & Visseren, F. L. (2009). Obesity and cancer: the role of dysfunctional adipose tissue. *Cancer Epidemiol Biomarkers Prev*, 18(10), 2569-2578.
- Vasseur, F., Meyre, D., & Froguel, P. (2006). Adiponectin, type 2 diabetes and the metabolic syndrome: lessons from human genetic studies. *Expert Rev Mol Med*, 8(27), 1-12.
- Vendrell, J., Broch, M., Vilarrasa, N., Molina, A., Gomez, J. M., Gutierrez, C. (2004). Resistin, adiponectin, ghrelin, leptin, and proinflammatory cytokines: relationships in obesity. *Obes Res*, 12(6), 962-971.
- Villalobos-Comparan, M., Teresa Flores-Dorantes, M., Teresa Villarreal-Molina, M., Rodriguez-Cruz, M., Garcia-Ulloa, A. C., Robles, L. (2008). The FTO gene is associated with adulthood obesity in the Mexican population. *Obesity (Silver Spring)*, 16(10), 2296-2301.
- Vimalaswaran, K. S., Li, S., Zhao, J. H., Luan, J., Bingham, S. A., Khaw, K. T. (2009). Physical activity attenuates the body mass index-increasing influence of genetic variation in the FTO gene. *Am J Clin Nutr*, 90(2), 425-428.
- Vinik, A. I. (2005). The metabolic basis of atherogenic dyslipidemia. *Clin Cornerstone*, 7(2-3), 27-35.
- Vojarova de Courten, B., Degawa-Yamauchi, M., Considine, R. V., & Tataranni, P. A. (2004). High serum resistin is associated with an increase in adiposity but not a worsening of insulin resistance in Pima Indians. *Diabetes*, 53(5), 1279-1284.
- Wahlen, K., Sjolín, E., & Hoffstedt, J. (2008). The common rs9939609 gene variant of the fat mass- and obesity-associated gene FTO is related to fat cell lipolysis. *J Lipid Res*, 49(3), 607-611.

- Wang, D., Ma, J., Zhang, S., Hinney, A., Hebebrand, J., Wang, Y. (2010). Association of the MC4R V103I polymorphism with obesity: a Chinese case-control study and meta-analysis in 55,195 individuals. *Obesity (Silver Spring)*, 18(3), 573-579.
- Wang, H., Chu, W. S., Hemphill, C., & Elbein, S. C. (2002). Human resistin gene: molecular scanning and evaluation of association with insulin sensitivity and type 2 diabetes in Caucasians. *J Clin Endocrinol Metab*, 87(6), 2520-2524.
- Wang, H. J., Zhang, H., Zhang, S. W., Pan, Y. P., & Ma, J. (2008). Association of the common genetic variant upstream of INSIG2 gene with obesity related phenotypes in Chinese children and adolescents. *Biomed Environ Sci*, 21(6), 528-536.
- Wang, K., Li, W. D., Zhang, C. K., Wang, Z., Glessner, J. T., Grant, S. F. (2011). A genome-wide association study on obesity and obesity-related traits. *PLoS One*, 6(4), e18939.
- Wang, T. N., Huang, M. C., Chang, W. T., Ko, A. M., Tsai, E. M., Liu, C. S. (2006). G-2548A polymorphism of the leptin gene is correlated with extreme obesity in Taiwanese aborigines. *Obesity (Silver Spring)*, 14(2), 183-187.
- Wannamethee, S. G., & Shaper, A. G. (1999). Weight change and duration of overweight and obesity in the incidence of type 2 diabetes. *Diabetes Care*, 22(8), 1266-1272.
- Wardle, J., Carnell, S., Haworth, C. M., Farooqi, I. S., O'Rahilly, S., & Plomin, R. (2008). Obesity associated genetic variation in FTO is associated with diminished satiety. *J Clin Endocrinol Metab*, 93(9), 3640-3643.
- Wardle, J., Carnell, S., Haworth, C. M., & Plomin, R. (2008). Evidence for a strong genetic influence on childhood adiposity despite the force of the obesogenic environment. *Am J Clin Nutr*, 87(2), 398-404.
- Wardle, J., Llewellyn, C., Sanderson, S., & Plomin, R. (2009). The FTO gene and measured food intake in children. *Int J Obes (Lond)*, 33(1), 42-45.
- Wauters, M., & Van Gaal, L. (1999). Gender differences in leptin levels and physiology: a role for leptin in human reproduction. *J Gend Specif Med*, 2(5), 46-51.
- Way, J. M., Gorgun, C. Z., Tong, Q., Uysal, K. T., Brown, K. K., Harrington, W. W. (2001). Adipose tissue resistin expression is severely suppressed in obesity and stimulated by peroxisome proliferator-activated receptor gamma agonists. *J Biol Chem*, 276(28), 25651-25653.
- Weisberg, S. P., McCann, D., Desai, M., Rosenbaum, M., Leibel, R. L., & Ferrante, A. W., Jr. (2003). Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest*, 112(12), 1796-1808.

- Westerterp, K. R., & Goran, M. I. (1997). Relationship between physical activity related energy expenditure and body composition: a gender difference. *Int J Obes Relat Metab Disord*, 21(3), 184-188.
- Weyer, C., Bogardus, C., Mott, D. M., & Pratley, R. E. (1999). The natural history of insulin secretory dysfunction and insulin resistance in the pathogenesis of type 2 diabetes mellitus. *J Clin Invest*, 104(6), 787-794.
- Weyer, C., Funahashi, T., Tanaka, S., Hotta, K., Matsuzawa, Y., Pratley, R. E. (2001). Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J Clin Endocrinol Metab*, 86(5), 1930-1935.
- Whitlock, G., Lewington, S., Sherliker, P., Clarke, R., Emberson, J., Halsey, J. (2009). Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*, 373(9669), 1083-1096.
- WHO. (2012a). 10 facts on obesity. Retrieved 20 September 2011 <http://www.who.int/features/factfiles/obesity/facts/en/index.html>
- WHO. (2012b). Global Strategy on Diet, Physical Activity and Health. Retrieved 15 February 2012 <http://www.who.int/dietphysicalactivity/pa/en/index.html>
- Wiecek, A., Kokot, F., Chudek, J., & Adamczak, M. (2002). The adipose tissue--a novel endocrine organ of interest to the nephrologist. *Nephrol Dial Transplant*, 17(2), 191-195.
- Wiedmann, S., Neureuther, K., Stark, K., Reinhard, W., Kallmunzer, B., Baessler, A. (2009). Lack of association between a common polymorphism near the INSIG2 gene and BMI, myocardial infarction, and cardiovascular risk factors. *Obesity (Silver Spring)*, 17(7), 1390-1395.
- Willer, C. J., Speliotes, E. K., Loos, R. J., Li, S., Lindgren, C. M., Heid, I. M. (2009). Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. *Nat Genet*, 41(1), 25-34.
- Wing, M. R., Ziegler, J., Langefeld, C. D., Ng, M. C., Haffner, S. M., Norris, J. M. (2009). Analysis of FTO gene variants with measures of obesity and glucose homeostasis in the IRAS Family Study. *Hum Genet*, 125(5-6), 615-626.
- Wu, Q., Saunders, R. A., Szkudlarek-Mikho, M., Serna Ide, L., & Chin, K. V. (2010). The obesity-associated Fto gene is a transcriptional coactivator. *Biochem Biophys Res Commun*, 401(3), 390-395.
- Xi, B., & Mi, J. (2009). FTO polymorphisms are associated with obesity but not with diabetes in East Asian populations: a meta-analysis. *Biomed Environ Sci*, 22(6), 449-457.
- Xi, B., Shen, Y., Zhang, M., Liu, X., Zhao, X., Wu, L. (2010). The common rs9939609 variant of the fat mass and obesity-associated gene is associated with obesity risk in children and adolescents of Beijing, China. *BMC Med Genet*, 11, 107.

- Xiang, Z., Litherland, S. A., Sorensen, N. B., Proneth, B., Wood, M. S., Shaw, A. M. (2006). Pharmacological characterization of 40 human melanocortin-4 receptor polymorphisms with the endogenous proopiomelanocortin-derived agonists and the agouti-related protein (AGRP) antagonist. *Biochemistry*, 45(23), 7277-7288.
- Xita, N., Georgiou, I., Chatzikyriakidou, A., Vounatsou, M., Papassotiriou, G. P., Papassotiriou, I. (2005). Effect of adiponectin gene polymorphisms on circulating adiponectin and insulin resistance indexes in women with polycystic ovary syndrome. *Clin Chem*, 51(2), 416-423.
- Xu, H., Barnes, G. T., Yang, Q., Tan, G., Yang, D., Chou, C. J. (2003). Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest*, 112(12), 1821-1830.
- Yabe, D., Brown, M. S., & Goldstein, J. L. (2002). Insig-2, a second endoplasmic reticulum protein that binds SCAP and blocks export of sterol regulatory element-binding proteins. *Proc Natl Acad Sci U S A*, 99(20), 12753-12758.
- Yajnik, C. S., Janipalli, C. S., Bhaskar, S., Kulkarni, S. R., Freathy, R. M., Prakash, S. (2009). FTO gene variants are strongly associated with type 2 diabetes in South Asian Indians. *Diabetologia*, 52(2), 247-252.
- Yamamoto, Y., Hirose, H., Saito, I., Nishikai, K., & Saruta, T. (2004). Adiponectin, an adipocyte-derived protein, predicts future insulin resistance: two-year follow-up study in Japanese population. *J Clin Endocrinol Metab*, 89(1), 87-90.
- Yanagiya, T., Tanabe, A., Iida, A., Saito, S., Sekine, A., Takahashi, A. (2007). Association of single-nucleotide polymorphisms in MTMR9 gene with obesity. *Hum Mol Genet*, 16(24), 3017-3026.
- Yang, J., Li, M., Wu, C. Y., Wang, H., Xu, Q. S., & Deng, J. Y. (2003). [Reduced resistin levels in patients with type 2 diabetes mellitus]. *Zhonghua Yi Xue Za Zhi*, 83(17), 1471-1474.
- Yang, L., Wu, Y., Li, H., Yu, Z., Li, X., Liu, Y. (2008). Potential association of INSIG2 rs7566605 polymorphism with body weight in a Chinese subpopulation. *Eur J Hum Genet*, 16(6), 759-761.
- Yang, W. S., & Chuang, L. M. (2006). Human genetics of adiponectin in the metabolic syndrome. *J Mol Med (Berl)*, 84(2), 112-121.
- Yannakoulia, M., Yiannakouris, N., Bluher, S., Matalas, A. L., Klimis-Zacas, D., & Mantzoros, C. S. (2003). Body fat mass and macronutrient intake in relation to circulating soluble leptin receptor, free leptin index, adiponectin, and resistin concentrations in healthy humans. *J Clin Endocrinol Metab*, 88(4), 1730-1736.
- Yeo, G. S., Connie Hung, C. C., Rochford, J., Keogh, J., Gray, J., Sivaramakrishnan, S. (2004). A de novo mutation affecting human TrkB associated with severe obesity and developmental delay. *Nat Neurosci*, 7(11), 1187-1189.

- Yiannakouris, N., Melistas, L., Yannakoulia, M., Mungal, K., & Mantzoros, C. S. (2003). The-2548G/A polymorphism in the human leptin gene promoter region is associated with plasma free leptin levels; interaction with adiposity and gender in healthy subjects. *Hormones (Athens)*, 2(4), 229-236.
- Yoshihiro Miyamoto, H. M., Yoshihiro Kokubo, Itaru Yamanaka, Hitonobu Tomoike, Akira Okayama, Yasunao Yoshimasa, Takayuki Morisaki (2009). Resistin gene variations are associated with the metabolic syndrome in Japanese men. *Obesity Research & Clinical Practice*, 3(2), 65-74.
- Young, E. H., Wareham, N. J., Farooqi, S., Hinney, A., Hebebrand, J., Scherag, A. (2007). The V103I polymorphism of the MC4R gene and obesity: population based studies and meta-analysis of 29 563 individuals. *Int J Obes (Lond)*, 31(9), 1437-1441.
- Yudkin, J. S., Stehouwer, C. D., Emeis, J. J., & Coppack, S. W. (1999). C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arterioscler Thromb Vasc Biol*, 19(4), 972-978.
- Zabena, C., Gonzalez-Sanchez, J. L., Martinez-Larrad, M. T., Torres-Garcia, A., Alvarez-Fernandez-Represa, J., Corbaton-Anchuelo, A. (2009). The FTO obesity gene. Genotyping and gene expression analysis in morbidly obese patients. *Obes Surg*, 19(1), 87-95.
- Zhang, B., Graziano, M. P., Doebber, T. W., Leibowitz, M. D., White-Carrington, S., Szalkowski, D. M. (1996). Down-regulation of the expression of the obese gene by an antidiabetic thiazolidinedione in Zucker diabetic fatty rats and db/db mice. *J Biol Chem*, 271(16), 9455-9459.
- Zhang, G., Karns, R., Narancic, N. S., Sun, G., Cheng, H., Missoni, S. (2010). Common SNPs in FTO gene are associated with obesity related anthropometric traits in an island population from the eastern Adriatic coast of Croatia. *PLoS One*, 5(4), e10375.
- Zhang, J., Lin, R., Wang, F., Lu, M., Lin, R. Y., Wang, S. Z. (2008). A common polymorphism is associated with body mass index in Uyghur population. *Diabetes Res Clin Pract*, 81(2), e11-13.
- Zhang, K., Calabrese, P., Nordborg, M., & Sun, F. (2002). Haplotype block structure and its applications to association studies: power and study designs. *Am J Hum Genet*, 71(6), 1386-1394.
- Zhang, K., & Kaufman, R. J. (2004). Signaling the unfolded protein response from the endoplasmic reticulum. *J Biol Chem*, 279(25), 25935-25938.
- Zhang, Y., Proenca, R., Maffei, M., Barone, M., Leopold, L., & Friedman, J. M. (1994). Positional cloning of the mouse obese gene and its human homologue. *Nature*, 372(6505), 425-432.

- Zhao, H., Pfeiffer, R., & Gail, M. H. (2003). Haplotype analysis in population genetics and association studies. *Pharmacogenomics*, 4(2), 171-178.
- Zhao, K., Aranzana, M. J., Kim, S., Lister, C., Shindo, C., Tang, C. (2007). An Arabidopsis example of association mapping in structured samples. *PLoS Genet*, 3(1), e4.