

REFERENCES

- Banks, W.A. (2009). Characteristics of compounds that cross the blood-brain barrier. BMC Neurology, 9, S3.
- Bishayee, A., Rao, D.V., Bouchet, L.G., Bolch, W.E., Howell, R.W. (2000). Protection by DMSO against cell death caused by intracellularly localized iodine-125, iodine-131 and polonium-210. Radiation Research, 153, 416-427.
- Capsoni, S., Giannotta, S., Cattaneo, A. (2002). Nerve growth factor and galantamine ameliorate early signs of neurodegeneration in anti-nerve growth factor mice. Proceedings of the National Academy of Sciences, 99, 12432–12437.
- Chang, S.T., & Miles, P.G. (2004). Mushroom: Cultivation, nutritional values, medicinal effect, and environmental impact. CRC Press. pp.385-387.
- Chatellier, G., & Lacomblez, L. (1990). Tacrine (tetrahydroaminoacridine; THA) and lecithin in senile dementia of the Alzheimer type: a multicentre trial. British Medical Journal, 300, 495-499.
- Chen, G., Luo, Y.C., Ji, B.P., Li, B., Guo, Y., Li, Y., Su, W., & Xiao, Z.L. (2008). Effect of Polysaccharide from *Auricularia auricula* on Blood Lipid Metabolism and Lipoprotein Lipase Activity of ICR Mice Fed a Cholesterol-Enriched Diet. Journal of Food Science, 73, 103-108.
- Cheung, P.C.K. (1996). The hypocholesterolemic effect of two edible mushrooms-*Auricularia auricula* (tree-ear) and *Tremella fuciformis* (white jelly-leaf) in hypercholesterolemic rats. Nutrition Research, 16, 1721-1725.
- Cheung, W.M.W., Hui, W.S., Chu, P.W.K., Chiu, S.W., & Ip, N.Y. (2000). Ganoderma extract activates MAP kinases and induces neuronal differentiation of rat pheochromocytoma PC12 cells. FEBS Letter, 486, 291-296.
- Darios, F., & Davletov, B. (2006). Omega-3 and omega-6 fatty acids stimulate cell membrane expansion by acting on syntaxin 3. Nature, 440, 813–817.
- Fan, Y.M., Xu, M.Y., Wang, L.Y., Zhang, Y., Zhang, L., Yang, H., Wang, P., & Cui, P. (1989). The effect of edible black tree fungus (*Auricularia auricula*) on experimental atherosclerosis in rabbits. Chinese Medical Journal, 102, 100-105.
- Farlow, M., Gracon, S.I., Hershey, L.A., Lewis, K.W., Sadowsky, C.H., & Dolan-Ureno, J. (1992). A controlled trial of tacrine in Alzheimer's disease. The Journal of the American Medical Association, 268, 2523–2529.
- Fischer, W., Wictorin, K., Bjorklund, A., Williams, L.R., Varon, S., & Gage, F.H. (1987). Amelioration of cholinergicneuron atrophy and spatila memory impairment in aged rats by nerve growth factor. Nature, 329, 65-68.

Francia, C., Rapior, S., Courtecuisse, R., & Siroux, Y. (1999). Current research findings on the effects of selected mushrooms on cardiovascular diseases. International Journal of Medicinal Mushrooms, 1, 169-172.

Fukaya, T., Gondaira, T., Kashiyae, Y., Kotani, S., Ishikura, Y., Fujikawa, S., Kiso, Y., & Sakakibara, M. (2007). Arachidonic acid preserves hippocampal neuron membrane fluidity in senescent rats. Neurobiology of Aging, 28, 1179–1186.

Goate, A. M., Chartier-Harlin, M.C., Mullan, M., Brown, J., Crawford,F., *et al.*, (1991). Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimers's disease. Nature, 349, 704-706.

Greene, L.A. (1977). A quantitative bioassay for nerve growth factor (NGF) activity employing a clonal pheochromocytoma cell line. Brain Research, 133, 350-353.

Greene, L.A., & Tischler, A.S. (1982). PC12 pheochromocytoma cultures in neurobiological research. Advances in Cellular Neurobiology, 3, 373-414.

Halliwell, B., & Gutteridge, J.C. (1999). Free radicals in biology and medicine (3rd ed.). London: Oxford University Press.

Halpern, G.M. (2007). Healing Mushrooms. Garden City (NY): Square One Publishers.

Hamprecht, B., Glaser, T., Reiser, G., Bayer, E., & Propst, F. (1985). Culture and characteristics of hormone-responsive neuroblastoma x glioma hybrid cells. Methods in enzymology, 109, 316-341.

Harrison, P.K., Connor, M., & Kelly, E. (1997). Chronic ethanol promotes the neuronal differentiation of NG108-15 cells independently of toxin-sensitive G-proteins. Environmental Toxicology and Pharmacology, 3, 307-319.

Hawksworth, D.L. (2001). The magnitude of fungal diversity: the 1.5 million species estimate revisited . Mycological Research, 12, 1422-1432.

Hobbs, C. (1995). Medicinal mushrooms: An exploration of tradition, healing and culture. Botanica Press, Santa Cruz, CA. pp.252.

Ismail, A., Marjan, Z.M., & Foong, C.W. (2004). Total antioxidant activity and phenolic content in selected vegetables. Food Chemistry, 87, 581-586.

Jong, S.C., & Denovick, R. (1989). Antitumor and antiviral substances from fungi. Advances in Applied Microbiology, 34, 183-262.

Kabir, Y., & Kimura, S. (1989). Dietary mushrooms reduce blood pressure in spontaneously hypertensive rats (SHR). Journal of Nutritional Science and Vitaminology, 35, 91-94.

Kabir, Y., Kimura, S., & Tamura, T. (1988). Dietary effect of Ganoderma lucidum mushroom on blood pressure and lipid levels in spontaneously hypertensive rats (SHR). Journal of Nutritional Science and Vitaminology, 34, 433-438.

Kawagishi, H., Ando, M., & Mizuno, T. (1990). Hericenone A and B as cytotoxic principles from the mushroom *Hericium erinaceum*. Tetrahedron Letter, 31, 373-376.

Kawagishi, H., Ando, M., Sakamoto, H., Yoshida, S., Ojima, F., Ishiguro, Y., Ukai, N., & Furukawa, S. (1991). Hericenones C, D, and E, stimulators of nerve growth factor (NGF)- synthesis, from mushroom *Hericium erinaceum*. Tetrahedron Letter, 32, 4561-4564.

Kawagishi, H., Ando, M., Sakamoto, H., Yoshida, S., Ojima, F., Ishiguro, Y., Ukai, N., & Furukawa, S. (1991). Hericenones C, D, and E, stimulators of nerve growth factor (NGF)- synthesis, from mushroom *Hericium erinaceum*. Tetrahedron Letter, 32, 4561-4564.

Kawagishi, H., Ando, M., Shinba, K., Sakamoto, H., Yoshida, S., Ojima, F., Ishiguro, Y., Ukai, N., & Furukawa, S. (1993). Chromans, hericenones F, G and H from the mushroom *Hericium erinaceum*. Phytochemistry, 32, 175-178.

Kawagishi, H., Furukawa, Y., Furukawa, S., Zhuang, C., & Yunoki, R. (2002). The inducer of the synthesis of nerve growth factor from Lion's Mane (*Hericium erinaceus*). Explore, 11, 4-51.

Kawagishi, H., Shimada, A., Hosokawa, S., Mori, H., Sakamoto, H., Ishiguro, Y., Sakemi, S., Bordner, J., Kojima, N., & Furukawa, S. (1996a). Erinacines E, F, and G, stimulators of nerve growth factor (NGF)-synthesis, from the mycelia of *Hericium erinaceum*. Tetrahedron Letters, 37, 7399-7402.

Kawagishi, H., Shimada, A., Shirai, R., Okamoto, K., Ojima, F., Sakamoto, H., Ishiguro, Y., & Furukawa, S. (1994). Erinacines A, B and C, strong stimulators of nerve growth factor (NGF)-synthesis, from the mycelia of *Hericium erinaceum*. Tetrahedron Letters, 35, 1569-1572.

Kawagishi, H., Shimada, A., Shizuki, K., Mori, H., Okamoto, K., Sakamoto, H., & Furukawa, S. (1996b). Erinacine D, a stimulator of NGF-synthesis, from the mycelia of *Hericium erinaceum*. Heterocyclic Communications, 2, 51-54.

Kawagishi, H., Shirai, R., Sakamoto, H., Yoshida, S., Ojima, F., & Ishiguro, Y. (1992). Erinapyrones A and B from the cultured mycelia of *Hericium erinaceum*. Chemistry Letters, 21, 2475-2476.

Keyzers, R.A., & Davies- Coleman, M.T. (2005). Anti-inflammatory metabolites from marine sponges. Chemical Society Reviews, 34, 355-365.

Kiho, T., Sobue, S., & Ukai, S. (1994). Structural features and hypoglycemic activities of two polysaccharides ffrom a hot-water extract of *Agrocybe cylindracea*. Carbohydrate Reseach, 251, 81-87.

Kiho, T., Yamane, A., Hui, J., Usui, S., & Ukai, S. (1996). Polysaccharides in fungi, XXXVI Hypoglycemic activity of a polysaccharide (CS-F30) from the cultural mycelium of *Cordyceps sinensis* and its effect on glucose metabolism in mouse liver. Biological and Pharmaceutical Bulletin, 19, 294-296.

Kim, D.M., Pyun, C.W., Ko, H.G., & Park, W.M. (2000). Isolation of antimicrobial substances from *Hericeum erinaceum*. *Mycobiology*, 28, 33-38.

Kim, J.H., Ha, H.C., Lee, M.S., Kang, J.I., Kim, H.S., Lee, S.Y., Pyun, K.H., & Shim, I. (2007). Effect of *Tremella fuciformis* on the neurite outgrowth of PC12h cells and the improvement of memory in rats. *Biological and Pharmaceutical Bulletin*, 30, 708-714.

Kirmizigul, S., Boke, N., Sumbul, H., Gokturk, R.S., & Arda, N. (2007). Essential fatty acid components and antioxidant activities of eight *Cephalaria* species from southwestern Anatolia. *Pure and Applied Chemistry*, 79, 2297-2304.

Kromer, L.F. (1987). Nerve growth factor treatment after brain injury prevents neuronal death. *Science*, 235, 214-216.

Kubo, K., Aoki, H., & Nanba, H. (1994). Anti-diabetic activity present in the fruit body of *Grifola frondosa* (Maitake). *Biological and Pharmaceutical Bulletin*, 17, 1106-1110.

Lee, L.K., Yun, B.S., Kim, J.P., Ryoo, I.J., Kim, Y.H., & Yoo, I.D. (2003). Neuroprotective activity of p-terphenyl leucomentins from the mushroom *Paxillus panuoides*. *Bioscience, Biotechnology and Biochemistry*, 67, 1813-1816.

Li, H., Park, S., Moon, B.K., Yoo, Y.B., Lee, Y.W., & Lee, C. (2012). Targeted phenolic analysis in *Hericium erinaceum* and its antioxidant activities. *Food Science Biotechnology*, 21, 881-888.

Littlefield, J.W. (1964). Selection of hybrids from matings of fibroblasts *in vitro* and their presumed recombinants. *Science*, 145, 709-710.

Lu, H., Zou, W.X., Meng, J.C., Hu, J., & Tan, R.X. (2000). New bioactive metabolites produced by *Colletotrichum sp.*, an endophytic fungus in *Artemesia annua*. *Plant Science*, 151, 67-73.

Malingre, M.M., Schellens, J.H., van Tellingen, O., Ouwehand, M., Bardelmeijer, H.A., Rosing, H., Koopman, F.J., Schot, M.E., Ten Bokkel Huinink, W.W., & Beijnen, J.H. (2001). The co-solvent Cremophor EL limits absorption of orally administered paclitaxel in cancer patients. *British Journal of Cancer*, 85, 1472-1477.

Manakunakorn, P., Tohda, M., Murakami, Y., Matsumoto, K., Watanabe, H., & Vajragupta, O. (2003). Cytoprotective and cytotoxic effects of curcumin: dual action on H₂O₂- induced oxidative cell damage in NG108-15 cells. *Biological and Pharmaceutical Bulletin*, 26, 725-728.

Mandana, B., Russly, A.R., Farah, S.T., Noranizam, M.A., Zaidul, I.S., & Ali, G. (2012). Antioxidant activity of winter melon (*Benincasa Hispida*) seeds using conventional soxhlet extraction technique. *International Food Research Journal*, 19, 229-234.

Mau, J.L., Chao, G.R., & Wu, K.T. (2001). Antioxidant properties of methanolic extracts from several ear mushrooms. *Journal of Agricultural and Food Chemistry*, 49, 5461-5467.

Mau, J.L., Lin, H.C., & Song, S.F. (2002). Antioxidant activity of several specialty mushrooms. Food Research International, 35, 519-526.

Mizuno, T. (1999). Bioactive substances in *Hericium erinaceus* (Bull.: Fr.) Pers. (Yamabushitake), and its medicinal utilization. International Journal of Medicinal Mushroom, 1, 105-119.

Miyazawa, M., Matsuda, N., Tamura, N. (2008). Characteristic flavor of volatile oil from dried fruiting bodies of *Hericium erinaceus* (Bull.: Fr.) Pers. Journal of Essential Oil Research, 20, 420-423.

Mori, K., Obara, Y., Hirota, M., Azumi, Y., Kinugasa, S., Inatomi, S., & Nakahata, N. (2008). Nerve growth factor-inducing activity of *Hericium erinaceus* in 1321N1 human astrocytoma cells. Biological and Pharmaceutical Bulletin, 31, 1727-1732.

Mullan, M., Houlden, H., Crawford, F., Kennedy, A., Rogues, P., Rossor, M. (1993). Age of onset in familial early onset Alzheimer's disease correlates with genetic aetiology. American Journal of Medical Genetics, 48, 129-130.

Nagai, K., Chiba, A., Nishini, T., Kubota, T., & Kawagishi. H. (2006). Dilinoleoyl-phosphatidylethanoamine from *Hericium erinaceum* protects against ER stress-dependent Neuro2a cell death via protein kinase C pathway. Journal of Nutritional Biochemistry, 17, 525-530.

Obara, Y., Nakahata, N., Kita, T., Takaya, Y., Kobayashi, H., Hosoi, S., Kiuchi, F., Ohta, T., Oshima, Y., & Ohizuma, Y. (1999). Stimulation of neurotrophic factor secretion from 1321N1 human astrocytoma cells by novel diterpenoids, scabronines A and G. European Journal of Pharmacology, 370, 79-84.

Ohta, T., Kita, T., Kobayashi, N., Obara, Y., Nakahata, N., Ohizumi, Y., Takaya, Y., & Oshima, Y. (1998). Scabronine A, a novel diterpenoid having a potent inductive activity of nerve growth factor synthesis isolated from the mushroom *Sarcodon scabrosus*. Tetrahedron Letters, 39, 6229-6232.

Okamoto, K., Shimada A., Shirai, R., Sakamoto, H., Yoshida, S., Ojima, F., Ishiguro, Y., & Sakai, T., & Kawagishi, H. (1993). Antimicrobial chlorinated orcinol derivaties from mycelia of *Hericium erinaceum*. Phytochemistry, 34, 1445-1446.

Oser, B.L., Oser, M. (1956). Nutritional studies on rats on diets containing high levels of partial ester emulsifiers. I. General plan and procedures; growth and food utilization. Journal of Nutrition, 60, 367-390.

O'Sullivan, S.M., Woods, J.A., & O'Brien, N.M. (2004). Use of Tween 40 and Tween 80 to deliver a mixture of phytochemicals to human colonic adenocarcinoma cell (CaCo-2) monolayers. British Journal of Nutrition, 91, 757-764.

Ricceri, L., Alleva, E., Chiarotti, F., & Calamandrei, G. (1996). Nerve growth factor affects passive avoidance learning and retention in developing mice. Brain Reseach Bulletin, 39, 219-226.

Rizzo, W. B., Watkins, P. A., Phillips, M. W., Cranin, D., Campbell, B., Avigan, J. (1986). Adrenoleukodystrophy: oleic acid lowers fibroblast saturated C22-26 fatty acids. *Neurology*, 36, 257-361.

Rizzo, W.B., Philips, M.W., Dammann, A.L., Leshner, R.T., Jennings, S.S., Avigan, J., & Proud, V.K. (1987). Adrenoleukodystrophy: Dietary oleic acid lowers hexacosanoate levels. *Annals of Neurology*, 21, 232-239.

Sarafianos, S.G., Marchand, B., Das, K., Himmel, D.M., Parniak, M.A., Hughes, S.H., Arnold, E. (2009). Structure and function of HIV-1 reverse transcriptase: Molecular mechanisms of polymerization and inhibition. *Journal of Molecular Biology*, 385, 693-713.

Schorderet, M. (1995). Alzheimer's disease: fundamental and therapeutic aspects. *Experientia*, 51, 99-105.

Shang, X.D., Wang, G.Y., Pan, W., Sun, R., Cai Y.H., Tan, Q., & Zhao, G.P. (2012). Advances in research on low molecular weight bioactive components of *Hericium erinaceus*. *Acta Edulis Fungi*, 19, 85-91.

Shimbo, M., Kawagishi, H., & Yokogoshi, H. (2005). Erinacine A increases catecholamine and nerve growth factor content in the central nervous system of rats. *Nutrition Research*, 25, 617-623.

Silva,D. (2004). Cellular and physiological effects of *Ganoderma lucidum* (Reishi). *Mini- Reviews in Medicinal Chemistry*, 4, 873-879.

Sim, K.S., Sri Nurestri, A.M., & Norhanom, A. W. (2010). Phenolic content and antioxidant activity of *Pereskia grandifolia Haw.* (Cactaceae) extracts. *Pharmacognosy Magazine*, 6, 248-254.

Smalheiser N. R. (1991). Role of laminin in stimulating rapid onset neurites in NG108-15 cells: relative contribution of attachment and motility responses. *Developmental Brain Research*, 62, 81-89.

Smalheiser N. R., & Schwartz N. B. (1987). Kinetic analysis of 'rapid onset' neurite formation in NG108-15 cells reveals a dual role for substratum-bound laminin. *Developmental Brain Research*, 34, 111-121.

Smit, M., Leng, J., & Klemke, R.L. (2003). Assay for neurite outgrowth quantification. *Biotechniques*, 35, 254-256.

Son, C.G., Shin, J.W., Cho, J.H., Cho, C.H., Yun, C.H., Chung, W., & Han, S.H. (2006). Macrophage activation abd nitric oxide production by water soluble components of *Hericium erinaceum*. *International Immunopharmacology*, 6, 1363-1369.

Swanson-Flatt, S.K., Day, C., Flatt, P.R., Gould, B.J., & Bailey, C.J. (1989). Glycaemic effects of traditional european plant treatments for diabetes. Studies in normal and streptozotocin diabetic mice. *Diabetes Research*, 10, 69-73.

Tabart, J., Kevers, C., Pincemail, J., Defraigne, J.O., & Dommes, J. (2006). Antioxidant capacity of black current varies with organ, season and cultivar. Journal of Agricultural and Food Chemistry, 54, 6271-6276.

Takaku, T., Kimura, Y., Okuda, H. (2001). Isolation of an antitumor compound from *Agaricus blazei Murill* and its mechanism of action. The Journal of Medicine, 131, 1409-1413.

Uterman, G. (1994). The apolipoprotein E connection. Current Biology, 4, 362-365.

Wang, H.X., & Ng, T.B. (2004). A new laccase from dried fruiting bodies of the monkey head mushroom *Hericium erinaceum*. Biochemical and Biophysical Research Communications, 322, 17-21.

Wang, J.C., Hu, S.H., Lee, W.L., & Tsai, L.Y. (2001). Antimutagenicity of extracts of *Hericium erinaceus*. Kaohsiung Journal of Medical Sciences, 17, 230-238.

Wang, J.C., Hu, S.H., Wang, J.T., Chen, K.S., & Chia, Y.C. (2005). Hypoglycemic effect of extract of *Hericium erinaceus*. Journal of Science of Food and Agriculture, 85, 641-646.

Wang, Z.J., Liang, C.L., Li, G.M., Yu, C.Y., Yin, M. (2006). Neuroprotective effects of arachidonic acid against oxidative stress on rat hippocampal slices. Chemico-biological Interactions, 163, 207–217.

Wasser, S.P., & Weis, A.L. (1999). Medicinal properties of substances occurring in higher Basidiomycetes mushrooms: Current perspective (review). International Journal of Medicinal Mushroom, 1, 31-62.

Weiss, J., & Liao, W. (2000). Addition of sugars influences colour of oil-in-water emulsions. Journal of Agricultural and Food Chemistry, 48, 5053-5060.

Wong, K. H., Murali Naidu, David, P., Mahmood A. A., Noorlidah Abdullah, Kuppusamy, U.R. & Vikineswary, S. (2011). Peripheral nerve regeneration following crush injury to rat peroneal nerve by aqueous extract of medicinal mushroom *Hericium erinaceus* (Bull.: Fr) Pers. (Aphyllophoromycetideae). Evidenced-Based Complementary and Alternative Medicine, 2010, 1-10.

Wong, K. H., Vikineswary S., Murali Naidu, David, P., Noorlidah Abdullah, Mahmood, A. A., & Kuppusamy, U.R. (2009). Functional recovery enhancement following nerve crush injury to rodent peroneal nerve by medicinal mushroom *Hericium erinaceus*. International Journal of Medicinal Mushrooms, 11, 225-236.

Wong, K.H., Vikineswary, S., Noorlidah Abdullah, Kuppusamy, U.R., & Murali Naidu (2009). Effects of cultivation techniques and processing on antimicrobial and antioxidant activities of *Hericium erinaceus* (Bull.: Fr.) Pers. Extracts. Food Technology and Biotechnology, 47, 47-56.

Wong, K.H., Vikineswary S., Noorlidah Abdullah, Murali Naidu, & Roger Keynes (2007). Activity of aqueous extracts of lion's mane mushroom *Hericium erinaceus* (Bull.: Fr.) Pers. (Aphyllophoromycetideae) on the neural cell line NG108-15. International Journal of Medicinal Mushroom, 9, 57-65.

Yang, B.K., Park, J.B., & Song, C.H. (2003). Hypolipidemic effect of an exobiopolymer produced from a submerged mycelia culture of *Hericium erinaceus*. Bioscience, Biotechnology and Biochemistry, 67, 1292-1298.

Yang, Q.Y., & Jong, S.C. (1989). Medicinal mushrooms in China. Mushroom Science, 9, 631-643.

Zhang, A.Q., Sun, P.L., Zhang, J.S., Tang, C.H., Fan, J.M., Shi, X.M., & Pan, Y.J. (2007). Structural investigation of a novel fucoglucogalactan isolated from the fruiting bodies of the fungus *Hericium erinaceus*. Food Chemistry, 104, 451-456.

Zhang, H.J., Yao, M., Morrison, R.A., & Chong, S. (2003). Commonly used surfactant, Tween 80, improves absorption of P-glycoprotein substrate, digoxin, in rats. Archives of Pharmacal Research, 26, 768-772.