

## REFERENCES

- Abdus, S.A., Imran, M., & Ashfaq, M. (2009). Estimation of genetic variability and correlation for grain yield components in rice (*Oryza sativa* L.). *American-Eurasian J. Agric. & Environ. Sci.*, 6(5), 585-590.
- Ahmed, S. A., Borua, I., Sarkar, C. R., & Thakur, A. C. (1996). Volatile component (2-acetyl-1-pyrroline) in scented rice. *Proceedings of the Seminar on Problems and Prospects of Agricultural Research and Development in North-east India, Assam Agricultural University, Jorhat, India, 27-28 November, 1995.* (pp 55-57).
- Ahn, S.N., Bollisch, C.N., & Tanksley, S.D. (1992). RFLP tagging of a gene for aroma in rice. *Theor. Appl. Genet.*, 84, 825–8.
- Azeez, M.A., & Shafi, M. (1966). Quality in rice. *Technical Bulletin No. 13. Department of Agriculture, West Pakistan, Government of West Pakistan*, pp.50.
- Bao, J. S., Cai, Y. Z., & Corke, H. (2001). Prediction of rice starch quality parameters by near-infra red reflectance spectroscopy. *J. of Food Sci.*, 66, 036-939.
- Bemer, D.K., & Hoff, B.J. (1986). Inheritance of scent in American long grain rice. *Crop Science*, 26, 876-878.
- Blakeney, A. B. (1992). Developing rice varieties with different textures and tastes. *Chemistry in Australia*, 59, 475-6.
- Bollich, C. N. (1989). Release of new rice cultivar Jasmine 85 in USA. *Intl. Rice Res. Newsl.*, 14, 12.
- Bounphanousay, C., Jaisil, P., Sanitchon, J., Fitzgerald, M., & Sackville Hamilton, N. R., (2008). Chemical and molecular characterization of fragrance in black glutinous rice from Lao PDR. *Asian J. of Plant Sci.*, 7(1), 1-7.
- Bradbury, L. M. T., Fitzgerald, T. L., Henry, R. J., Jin, Q., & Waters, D. L. E. (2005a). The gene for fragrance in rice. *Plant Biotechnology Journal*, 3, 363-370.
- Bradbury, L. M. T., Henry, R. J., Jin, Q., Reinke, R. F., & Waters, D. L. E. (2005b). A perfect marker for fragrance genotyping in rice. *Mol. Breed.*, 16, 279-283.
- Bradbury, L. M. T., Gillies, S.A., Brushett, D.J., Waters, D. L. E., Henry, R.J. (2008). Inactivation of an aminoaldehyde dehydrogenase is responsible for fragrance in rice. *Plant Mol. Biol.*, 68, 439–49.
- Bryant, R.J., & McClung, A.M. (2011). Volatile profile of aromatic and non-aromatic rice cultivars using SPME/GC-MS. *Food Chemistry*, 124 (2), 501-513.
- Bullard, R. W., & Holguin, G. (1977). Volatile components of unprocessed rice (*Oryza sativa*). *J. Agric. Food Chem.*, 25, 99.
- Buttery, R. G., & Ling, L. C. (1982). Cooked rice aroma and 2-acetyl-1-pyrroline in rice. *Chem. Ind (London)*, 24, 958-959.

- Buttery, R. G., & Ling, L. C. (1995). Volatile flavor components of corn tortillars and related products. *J. Agric. Food Chem.*, *43*, 1878-1882.
- Buttery, R. G., Juliano, B. O., & Ling, L.C. (1983b). Identification of rice aroma compound 2-acetyl-1-pyrroline in pandan leaves. *Chem. Ind. (London)*, *23*, 478-479.
- Buttery, R. G., Ling, L. C., Juliano, B. O. & Tumbaugh, J. (1983a). Cooked rice aroma and 2-acetyl-1-pyrroline. *J. Agric Food Chem.*, *31*, 823-826.
- Buttery, R. G., Turnbaugh, J. G. & Ling, L.C. (1988). Contributions of volatiles to rice aroma. *J. Agric. Food Chem.*, *36*, 1006-1009.
- Buttery, R. G., Ling, L. C., & Mon, T. R. (1986). Quantitative analysis of 2-acetyl-1-pyrroline in rice. *J. Agric. Food Chem.*, *34*, 112-114.
- Buu, B.C., & Truong, D.X. (1988). Path analysis of rice grain yield under saline conditions. *Inter. Rice Res. Newsletter*, *13*, 20-21.
- Carrapiso, A. I., Jurado, A., Timon, M. L. & Garcia, C. (2002). Odor-active compounds of Iberian hams with different aroma characteristics. *J. Agric. Food Chem.*, *50*, 6453-6458.
- Champagne, E. T. (2008). Rice aroma and flavor: a literature review. *Cereal Chem.*, *85*, 445-54.
- Chen, S., Wu, J., Yang, Y., Shi, W., & Xu, M. (2006). The *fgr* gene responsible for rice fragrance was restricted within 69 kb. *Plant Science*, *171*, 505-514.
- Chen, S., Yang, Y., Shi, W., Ji, Q., He, F., Zhang, Z., et al. (2008). *Badh2*, encoding betaine aldehyde dehydrogenase, inhibits the biosynthesis of 2-acetyl-1-pyrroline, a major component in rice fragrance. *Plant Cell*, *20*, 1850-61.
- Cordeiro, G. M., Christopher, M. J., Henry, R. J., & Reinke, R. F. (2002). Identification of microsatellite markers for fragrance in rice by analysis of the rice genome sequence. *Mol. Breed.*, *9*, 245-250.
- Dela Cruz, N., & Khush, G. S. (2000). Rice Grain Quality Evaluation Procedure. In: R. K. Singh, U. S. Singh & G. S. Khush (Eds.), *Aromatic rices* (pp. 15-28). New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.
- Dhulappanavar, C.V. (1976). Inheritance of scent in rice. *Euphytica*, *25*, 659-662.
- Dhulappanavar, C. V., & Mensinkai, S.W. (1969). Inheritance of scent in rice. *Karnataka Univ. J.*, *14*, 125-129.
- Efferson, J.N. (1985). Rice quality in world markets. In: *Rice Grain Quality and Marketing*, International Rice Research Insititute, Los Banos, Philippines, pp. 1-13.
- Faruq, G., Mohamad, O., Hadjim, K., & Craig, A. M. (2004). Inheritance of kernel elongation in rice. *Int J Agri Biol.* *1560 8530 /2004 /06*, *5*, 813-815.

- Faruq, G., Mohamad, O., Hadzim, K. & Craig, M. A. (2003). Optimization of aging time and temperature of four Malaysian rice cultivars. *Pakistan J. Nutr.*, 2, 125–31.
- Faruq, G., NorZulaani, K., Jennifer, A. H., Subha, B., Zulqarnain, M., Nazia, A. M., & Mohammad, O. (2010). Evaluation of kernel elongation ratio and aroma association in global popular aromatic rice cultivars in tropical environment. *Afr J Agric Res.*, 5 (12), 1515-1522.
- Faruq, G., Yeap, H. Y., Masitah, A., Afnierna, N., Nazia, A. M., Norzulaani, K., Mohamad, O. (2011). Analysis of aroma and yield components of aromatic rice in Malaysian tropical environment. *Australian J. of Crop Science*, 5 (11), 1318-1325.
- Fushimi, T., Itani, T., Kohyama, N., & Sekiya, K. (1995). Variation of 2-acetyl-1-pyrroline concentration of the aromatic rice (cv. Hier) cultivated at Kubokawa-area in Kochi prefecture (abst). In: R. Ishii, & T. Horie (Eds.), *Crop Research in Asia, 1996: Achievements and Perspectives: Proc. 2nd Asia Crop Sci. Conf. at Fukui, Japan, August 21-23, 1995* (pp. 738-739).
- George, S. P., Dijee Bastian, Radhakrishnan, N. V., & Aipe, K. C. (2005). Short communication: Evaluation of aromatic rice varieties in Wayanad, Kerala. *Journal of Tropical Agriculture*, 43(1-2), 67-69.
- Ghose, R.L.M., & Butany, W.T. (1952). Studies on the inheritance of some characters in rice (*Oryza sativa* L.). *Indian J. Genet. Plant Breed.*, 12, 26-30.
- Ghosh, A. K., & Govindswamy, S. (1972). Inheritance of starch iodine blue value and alkali digestion value in rice and their genetic association. *Riso*, 21, 123-132.
- Girish, T.N., Gireesha, T.M., Vaishali, M.G., Hnamareddy B.G., & Hittalmani, S. (2006). Response of a new IR50/Moroberekan recombinant inbred population of rice (*Oryza sativa* L.) from an indicaxjaponica cross for growth and yield traits under aerobic conditions. *J. Euphytica*, 152(2), 149-161.
- Goff, S. A., Ricke, D., Lan, T. H., Presting, G., Wang, R., Dunn, M., et al. (2002). A draft sequence of the rice genome (*Oryza sativa* L. ssp japonica). *Science*, 296, 92-100.
- Grosch W., & Schieberle, P. (1997). Flavor of cereal products-A review. *Cereal. Chem.*, 74, 91-97.
- Habib, S.H., Bashir, M.K., Khalequzzaman, M., Ahmed, M.S., & Rashid, E.S.M.H. (2005). Genetic analysis and morphological selection criteria for traditional Broin Bangladesh rice germplasms. *J. Biol. Sci.*, 5(3), 315-318.
- Heda, G. D., & Reddy, G. M. (1984). Genetic analysis of cooking qualities of rice. *J.Cytol Gent.*, 19, 38-42.
- Heda, G.D., & Reddy, G. M. (1986). Studies on the inheritance of amylose content and Gelatinization temperature in rice. *Genet. Agric.*, 40, 1-8.

- Herderich, M., Costello, P. J., Grbin, P. R., & Henschke, P. A. (1995). Occurrence of 2-acetyl-1-pyrroline in mousy wines. *Nat. Product Lett.*, 7, 129-132.
- Huang, N., McCouch, S. R., Mew, T., & Parco, A. (1994). Development of an RFLP Map from a Doubled Haploid Population in Rice. *Rice Genet. Newsl.*, 11, 134-137.
- Huang, T. C., Teng, C. S., Chang, J. L., Chuang, H.S., Ho, C.T., Wu, M.L. (2008). Biosynthetic mechanism of 2-acetyl-1-pyrroline and its relationship with  $\Delta$ 1-pyrroline-5-carboxylic acid and methylglyoxal in aromatic rice (*Oryza sativa* L.) callus. *J. Agric. Food Chem.*, 56, 7399-404.
- Hossain, M.F., Bhuiya, M.S.U., & Ahmed, M. (2005). Morphological and Agronomic Attributes of some local and modern aromatic rice varieties of Bangladesh. *Asian J. of Plant Sci.*, 4(6), 664-666.
- IBPGR-IRRI rice advisory committee. (1980). Descriptors for rice (*Oryza sativa*). International Rice Research Institute, Manila, Philippines.
- Islam, M. A. (1983). Effect of heterozygosity on grain and nutritional quality of rice (*Oryza sativa* L.). Master 's thesis, Univ. Philippines, Los Banos.
- Itani, T., & Fushimi, T. (1996). Influence of pre-and post-harvest conditions on 2-acetyl-1-pyrroline concentration in aromatic rice (abst.). In: R. Ishii, & T. Horie (Eds.), *Crop Research in Asia, 1996: Achievements and Perspectives: Proc. 2nd Asia Crop Sci. Conf. at Fukui, Japan, August 21-23, 1995* (pp. 728-729).
- Jennings, P. R., Coffman, W. R., & Kauffman, H. E. (1979). Rice improvements. International Rice Research Institute, Los Banos, Philippines., pp.186.
- Jin, Q., Waters, D., Cordeiro, G. M., Henry, R. J., & Reinke, R. F. (2003). A single nucleotide polymorphism (SNP) marker lined to the fragrance gene in rice (*Oryza sativa* L.). *Plant Sci.*, 165, 359-64.
- Jodon, N. E. & Sonnier, E. A. (1973). Registration of Della rice. *Crop Science*, 13, 773.
- Juliano, B.O. (1971). A simplified assay for milled-rice amylose. *Cereal Sci. Today* , 16, 334-340,360.
- Juliano, B. O. (1972). Physicochemical properties of starch and protein in relation to grain quality and nutrition value of rice. In: *Rice Breeding*. IRRI, Manila, Philippines, pp.389-404.
- Juliano, B. O., & Duff, D. (1991). Rice grain quality as an emerging priority in National rice Breeding programmes. In: *Rice grain marketing and quality issues*. IRRI, Los Banos, Laguna, pp.55-64.
- Juliano, B. O. & Perez, C. M. (1984). Result of a collaborative test on the measurement of grain elongation of milled rice during cooking. *J. Cereal Sci.*, 2, 281-292.
- Juliano, B.O., & Villareal, C. P. (1993). Grain quality evaluation of world rice. International Rice Research Institute, Manila, Philippines, pp.205.

- Kadam, B.S., & Patankar, V.K. (1983). Inheritance of aroma in rice. *Chron. Bot.*, 4, 32.
- Kaosa-ard, M., & Juliano, B.O. (1991). Assessing rice quality characteristics and prices in selected international markets. In: Rice Grains Marketing and Quality Issues. Selected papers from the International Rice Research Conference. International Rice Research Institute, Los Banos, Laguna, Philippines, pp 25-35.
- Khush, G.S. (1997). Origin, dispersal, cultivation and variation of rice. *Plant Mol. Biol.*, 35, 25-34.
- Khush G.S. (2000). Taxonomy and origin of rice. In: R. K. Singh, U. S. Singh & G. S. Khush (Eds.), *Aromatic rices* (pp. 5-14). New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.
- Khush, G. S., Kumar, I., & Varmani, S. S. (1988). Grain quality of hybrid rice. In: Hybrid rice. IRRI, Manila, Philippines, pp. 201-215.
- Khush, G.S., Paule, C.M., & Delacruz, N.M. (1979). Rice grain quality evaluation and improvement of IRRI. In *Proc.on Workshop of Chemicals Aspects of Rice Grain Quality*, Los Banos, Philippines, pp.21–31.
- Kibria, K., Islam, M. M. & Begum, S. N. (2008). Screening of aromatic rice lines by phenotypic and molecular markers. *Bangladesh J. Bot.*, 37(2), 141-147.
- Kim, C. Y. (1999). A study on the Growth Characteristics and Analysis of Chemical Compounds of Grains as Affected by Cultivation Method in Coloured and Aromatic Rice Varieties. Ph.D. Thesis submitted to Chungnam National University, Taejeon, Korea. pp.112.
- Kirstin W, Wootton M. Flavour qualities of new Australian fragrant rice cultivars; 2004. RIRDC Publication No 04/ 160 RIRDC Project No UNS-12A <http://www.rirdc.gov.au/reports/RIC/04-160sum.html>.
- Kumar, I., & Khush, G. S. (1986). Gene dosage effect of amylose content in rice endosperm. *Japanese J. Genet.*, 61, 559-568.
- Kumazawa, K., & Masuda, H. (2002). Identification of potent odorants in different green tea varieties using flavor dilution technique. *J. Agric. Food Chem.*, 50, 5660-5663.
- Laksanalamai, V., & Ilangantileke, S. (1994). Comparison of aroma compound (2-acetyl-1-pyrroline) in leaves from pandan (*Pandanus amaryllifolius*) and Thai fragrant rice (Khao Dawk Mali-105). *Cereal Chem.*, 70, 381-384.
- Lin, C. F., Hsieh, R. C. Y., & Hoff, B. J. (1990). Identification and quantification of the popcorn-like aroma in Louisiana aromatic Della rice (*Oryza sativa*). *J. Food Sci.*, 35, 1466-1467.
- Liu, Q., Li, Q., Cai, X., Wang, H., Tang, S., Yu, H., Wang, Z., & Gu, M. (2006). Molecular marker-assisted selection for improved cooking and eating quality of two elite parents of hybrid rice. *Crop Sci.*, 46, 2354–2360.

- Lorieux, M., Petrov, M., Huang, N., Guiderdoni, E., & Ghesquiere, A. (1996). Aroma in rice: genetic analysis of quantitative trait. *Theor. Appl. Genet.*, *93*, 1145-1151.
- Maga, J. A. (1984). 'Rice product volatiles': a review. *J. Agric. Food Chem.*, *32*, 964-970.
- Mahadevappa, M., Shankaregowda, B. T. & Reddy, T. G. (1977). Scope for scented rice improvement in Karnataka. *Curr. Res.*, *6*, 1-3.
- Malaysian Agricultural Research & Development Institute, MARDI. (2007). Commercial production of high quality rice MRQ 74. <http://www.mardi.my>.
- Mann, R. A. (1987). Basmati rice: a wonder of Pakistan's agriculture IRRC Newslett. XXXVI (1), 23-28.
- Maningat, C. C., & Juliano, B. O. (1978). Alkali digestibility pattern, apparent solubility and gel consistency of milled rice. *Starch/Staerke*, *30*, 125-127.
- McKenzie, K. S., & Rutger, J. N. (1983). Genetic analysis of amylose content, alkali spreading score and grain dimensions in rice. *Crop Science*, *23*, 306-313.
- Mittal, U. K., Preet, K., Singh, D., Shukla, K. K., & Saini, R. G. (1995). Variability for aroma in some land races and cultivars of scented rice. *Crop Improv.*, *22*, 109-112.
- Mohamad, O., Mohd. Nazir, B., Alias, I., Azlan, S., Abdul Rahim, H., Abdullah, M. Z., et al. (2006, May). Development of improved rice varieties through the use of induced mutations in Malaysia. *Plant Mutation Report*, *1*(1), 26-27. Retrieved November 20, 2011, from <http://www-naweb.iaea.org/nafa/pbg/public/pmr-01-01.pdf>
- Mohamad, O., Hadzim, K., Azlan, S., Abdullah, M. Z., Amiran, N., Nur Samahah, M. Z., et al. (2008). Molecular screening for high quality rice. Retrieved November 20, 2011, from <http://www.fao.org/ag/agp/agpc/doc/services/pbn/pbn195.htm#a134>
- Myint Yi, Nwea, K. T., Vanavichit, A., Chai-arree, W., & Toojinda, T. (2009). Marker assisted backcross breeding to improve cooking quality traits in Myanmar rice cultivar Manawthukha. *Field Crops Research*, *113*, 178-186.
- Muraue, B. (1997). The advance of rice taste evaluation. Japanese Farming News Press, 7-164.
- Murrey, S. S. (2004). Searching for the origins of African rice domestication. *Antiquity*, *78*(300). Retrieved October 3, 2007, from <http://antiquity.ac.uk/ProjGall/murray/index.html>.
- Nagaraju, M., Choudhary, D., & Rao, M.J.B. (1975). A simple technique to identify scent in rice and inheritance pattern of scent. *Curr. Sci.*, *44*, 59.
- Nguyen, T. L., & Bui, C. B. (2008). Development of PCR-based markers for aroma (*fgr*) gene in rice (*Oryza sativa L.*). *Omonrice*, *16*, 16-23.

- Nijssen, L. M., Vischer, C. A., Maarse, H., Willensens, L. C., & Boelens, M. H. (1996). Volatile compounds in food. Qualitative and quantitative data. In: Rice (No.123). Central Institute for Nutrition and Food Research: Zeist, The Netherlands.
- Niu, X., Tang, W., Huang, W., Ren, G., Wang, Q., Luo, D., et al. (2008). RNAi-directed down regulation of OsBADH2 results in aroma (2-acetyl-1-pyrroline) production in rice (*Oryza sativa* L.) *BMC Plant Biol.*, 8, 1-10.
- Ookawa, T., & Ishihara, K. (1993). Varietal differences in physical characteristics of the culm related to lodging resistance in paddy rice. *Tpn. J. Crop Sci.*, 61, 419-425.
- Panwar, D. V. S., Gupta, K. R., Battan, K. R., & Singh, A. (1991). HKR 228, a semi dwarf aromatic rice strain for Havana, India. *Intl. Rice Res. Newsl.*, 16, 16-17.
- Paule, C. M., & Powers, J. J. (1989). Sensory and chemical examination aromatic and nonaromatic rices. *J. Food Sci.*, 54, 343-346.
- Perez, C. M. & Juliono, B. O. (1981). Texture changes and storage of rice. *J. Texture Stud.*, 12, 321-33.
- Pinson, S.R.M. (1994). Inheritance of aroma in six cultivars. *Crop Sci.*, 34, 1151-1157.
- Rasheed, M.S., Sadaqat, H.A., & Babar, M. (2002). Correlation and path co-efficient analysis for yield and its components in rice. *Asian J. Pl. Sci.*, 1(3), 241-244.
- Reddy, V. D., & Reddy, G. M. (1987). Genetic and biochemical basis of scent in rice (*Oryza sativa*). *Theor. Appl. Genet.*, 73, 699-700.
- Reddy, P. R. & Sathyanarayanaiah, K. (1980). Inheritance of aroma in rice. *Indian J. Genet. Plant Breed.*, 40, 327-329.
- Reinke, R.F., Welsch, L.A., Reece, J.E., Lewin, L.G., Blakeney, A.B. (1991). Procedure for quality selection of aromatic rice varieties. *Int Rice Res Newsl*, 16, 10-1.
- Rister, M. E., Goodwin, H. L., Brabson, R. R., Stamel, J. W., & Webb, B. D. (1992). The U.S. Aromatic rice market-results of an Asian-American household test panel. Texas Agric. Exp. Stn. Texas A & M Univ., College Station, TX.
- Saini, S. S., & Gagneja, M. R. (1973). PAU 29-295, a superfine scented rice variety. *Oryza* 10, 92, 541-551.
- Saini, S. S., & Kumar, Ish. (1978). Advances in breeding high yielding scented rice variety. *Oryza* 15, 216-218.
- Sakthivel, K., Sundaram, R. M., Shobha Rani, N., Balachandran, S. M., & Neeraja, C. N. (2009). Genetic and molecular basis of fragrance in rice. *Biotechnology Advances*, 27, 468-473.
- Sarhadi, W. A., Hien, N. L., Zanjani, M., Yosofzai, W., Yoshihashi, T., & Hirata, Y. (2008). Comparative analyses for aroma and agronomic traits of native rice cultivars from Central Asia. *J. Crop Sci. Biotech.*, 11 (1), 17-22.

- Sanjiva, R. B., Vasudeva Murthy, A. R., & Subrahmanya, R. S. (1952). The amylose and the amylopectin contents of rice and their influence on the cooking quality of the cereal. *Proc. Indian Acad. Sci.*, 36B, (2), 70-80.
- Schieberle, P., (1995). Quantitation of important roast-smelling odorants in popcorn by stable-isotope dilution assays and model studies on flavor formation during popping. *J. Agric. Food Chem.*, 43, 2442-2448.
- Schieberle, P., & Grosch, W. (1985). Identification of volatile flavour compounds of wheat bread crust-comparison with rye bread crust. *Z. Lebensm-Unters. Forsch.*, 180, 474-478.
- Sekhar, B. P. S., & Reddy, G. M. (1982). Amino acid profiles in some scented rice varieties. *Theor. Appl. Genet.*, 62, 35-37.
- Seguchi, M., Hayashi, M., Suzuki, Y., Sano, Y., & Hirano, Y.H. (2003). Role of amylose in the maintenance of the configuration of rice starch granules. *Starch/Staerke*, 55, 524-528.
- Shobha Rani, N., Pandey, M. K., Prasad, G. S. V., & Sudharshan, I. (2006). Historical significance, grain quality features and precision breeding for improvement of export quality basmati varieties in India. *Indian J. Crop Science*, 1 (1-2), 29-41.
- Shukla, R.S., & Chandel, P.S. (2004). Cytogenetics evolution and plant breeding. (3<sup>rd</sup> ed.). New Delhi: S. Chand & Company Ltd.
- Singh, B. D. (2008). Plant Breeding: Principles and Methods. Hybridization: Techniques and consequences. (7<sup>th</sup> ed.). New Delhi: Kalyani Publishers.
- Singh, M. V., Tripathi, H. N., & Tripathi, H. P. (1997). Effect of nitrogen and planting date on yield and quality of scented rice (*Oryza sativa*). *Indian J. Agron.*, 42, 602-606.
- Singh, R. K., Gautam, P. L., Sanjeev Saxena, & Singh, S. (2000a). Scented rice germplasm: conservation, evaluation and utilization. In: R. K. Singh, U. S. Singh & G. S. Khush (Eds.), *Aromatic rices* (pp. 107-134). New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.
- Singh, R. K., Khush, U. S., Singh, A. K., & Sanjay Singh. (2000b). Breeding aromatic rice for high yield, improved aroma and grain quality. In: R. K. Singh, U. S. Singh & G. S. Khush (Eds.), *Aromatic rices* (pp. 71-106). New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.
- Singh, U. S., Rashmi Rohilla, Amita Singh, Tewari, S. N., & Singh, H. M. (2000c). Plant protection in aromatic rice. In: R. K. Singh, U. S. Singh & G. S. Khush (Eds.), *Aromatic rices* (pp. 217-256). New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.
- Singh, R.K., Singh, U. S., & Khush, G. S. (2000d). Prologue. In: R. K. Singh, U. S. Singh & G. S. Khush (Eds.), *Aromatic rices* (pp. 1-4). New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.



- Sood, B. C., & Siddiq, E. A. (1978). A rapid technique for scent determination in rice. *Indian J. Genet. Plant Breed.*, 38, 268-271.
- Sood, G. B., & Siddiq, E. A. (1979). Geographical distribution of kernel elongation gene(s) in rice. *Indian J. Genet. Plant Breed.*, 40, 439-442.
- Sood, G. B., Siddiq, E. A. & Zaman, U. F. (1983). Genetic analysis of kernel elongation in rice. *Indian J. Genet. Plant Breed.*, 43, 40-43.
- Subudhi, P. K., Sasaki, T., & Khush, G. S. (2006). Rice. Genome Mapping and Molecular Breeding in Plants. In: C. Kole (Ed.), Cereals and millets. pp.1-8.
- Sun, S. X., Gao, F. Y., Lu, X. J., Wu, X. J., Wang, X. D., Ren, G. J., & Luo, H. (2008). Genetic analysis and gene fine mapping of aroma in rice (*Oryza sativa* L. Cyperales, Poaceae). *Genetics and Molecular Biology*, 31(2), 532-538.
- Suvamalatha, G., Narayan, M. S., Ravishankar, G. A. & Venkataraman, L. V. (1994). Flavour production in plant cell cultures of Basmati rice. *J. Sci. Food Agric.*, 66, 439-442.
- Tanchotikul, U., & Hsieh, T. C. Y. (1991). An improved method for quantification of 2-acetyl-1-pyrroline a popcorn-like aroma, in aromatic rice by high-resolution gas chromatography/mass spectrometry/selected ion monitoring. *J. Agri. Food Chem.*, 39, 944-947.
- Tomar, J. B., & Nanda, J. S. (1985). Genetics and association studies of kernel shape in rice. *Indian J. Genet.*, 45 (2), 278-283.
- Timmer, C. P. (2010). The changing role of rice in Asia's food security. ADB Sustainable Development Working Paper Series No. 15
- Toojinda, T., Tragoonrung, S., Vanavichit, A., Siangliw, J.L., Pa-In, N., Jantaboon, J., Siangliw, M., & Fukai, S. (2005). Molecular breeding for rainfed lowland rice in the Mekong region. *Plant Prod. Sci.*, 8, 330-333.
- Tripathi, R.S., & Rao, M.J. (1979). Inheritance and linkage relationship of scent in rice. *Euphytica*, 28, 319-323.
- Tsugita, T. (1985-1986). Aroma of cooked rice. *Food Rev. Intl.*, 1, 497-520.
- Tsuzuki, E., Kurata, T., & Kato, H. (1980). *Agric. Biol. Chem.*, 44, 835.
- Tsuzuki, E., Tanaka, K., & Shida, S. (1981). *Bull. Fac. Agric. Miyazaki Univ.*, 28, 31-37.
- Varzirzanjani et al., 2011, Grain Quality Laboratory lab at International Rice Research Institute (IRRI).
- Vanavichit, A., Yoshihashi, T., Wanchana, S., Areekit, S., Saengraku, D., Kamolsukyonyong, W., et al. (2005). Cloning of Os2AP, the aromatic gene controlling the biosynthetic switch of 2-acetyl-1-pyrroline and gamma

aminobutyric acid (GABA) in rice. 5th International Rice Genetics Symposium. Philippines, IRRI, November 19–23, pp. 44.

- Varaporn, L., & Sarath, I. (1993). Comparison of Aroma Compound (2-acetyl-1-pyrroline) in Leaves from Pandan (*Pandanus amaryllifolius*) and Thai Fragrant Rice (Khao Dawk Mali-105). *Cereal Chem.*, 70(4), 381-384.
- Villaral, R. M., Resurreccion, P. A. Suzuki, B. L., & Juliono, O. B. (1976). Changes in physioco-chemical properties of rice during storage. *Starch*, 28, 88-94.
- Weber, D. J., Rohilla, R., & Singh, U. S. (2000). Chemistry and biochemistry of aroma in scented rice. In: R. K. Singh, U. S. Singh & G. S. Khush (Eds.), *Aromatic rices* (pp. 29-46). New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd.
- Widjaja, R., Craske, J. D., & Wootton, M. (1996). Comparative studies on volatile components of non-fragrant and fragrant rices. *J. Sci Food Agric.*, 70, 151-161.
- Wongpornchai, S., Sriseadka, T., & Choovisase, S. (2003). Identification and quantitation of the rice aroma compound, 2-acetyl-1-pyrroline, in bread flowers (*Vallaris glabra* Ktze). *J. Agric. Food Chem.*, 51, 457-462.
- Yajima, I., Yanai, T., & Nakamura, M. (1978). Volatile flavour components of cooked rice. *Agric Biol. Chem.*, 42, 1229-1233.
- Yajima, I., Yanai, T., Nakamura, M., Sakakibara, H., & Habu, T. (1979). Volatile flavor components of cooked rice kaorimai (scented rice, *O. sativa japonica*), *J. Agric. Biol. Chem.*, 43, 2425-2429.
- Yoshihashi, T., Nguyen, T. T. H., & Kabaki, N. (2002). Quality evaluation of Khao Dawk Mali 105, an aromatic rice variety of Northeast Thailand. JIRCAS Working Report No. 30
- Zahid, M.A., Akhtar, M., Sabir, M., Manzoor, Z. & Awan, T. (2006). Correlation and path analysis studies of yield and economic traits in Basmati rice (*Oryza sativa* L.), *Asian J. Pl. Sci.*, 5(4), 643-645.
- Zehentbauer, G., & Grosch, W. (1998). Crust aroma of Baguettes-1. Key odorants of baguettes prepared in two different ways. *J. Cereal Sci.*, 28, 81-92.
- Zehentbauer, G., & Reineccius, G. A., (2002). Determination of key aroma components of Cheddar cheese using dynamic headspace dilution assay. *Flavor Fragrance J.*, 17, 300-305.
- Zhang, W. & Jiarong, Y. (1998). A preliminary study of ancient excavated rice from Yuchanyan site, Dao country, Human province, People's Republic of China. *Acta Agronomica Sinica*, 24 (4), 416-420.
- Zhang, Q. (2007). Strategies for developing green super rice. *Proc. Natl. Acad. Sci. U.S.A.*, 104, 16402–16409.
- Zhang, Q., & Yu, S. (2000). Molecular marker-based gene tagging and its impact on

rice improvement. In: J.S. Nanda, (Ed.). Rice breeding and genetics research priorities and challenges (pp. 241-270). New York: Science Publishers, Inc.

Zhang, X. M., Chun, H. S., Jian, G. W., Horiuchi, H., Tomita, K., Shui, Y. F., Gen, L. B., & Sheng, H.Y. (2003). Analysis of variations in the amylose content of grains located at different positions in the rice panicle and the effect of milling. *Starch/Staerke*, 55(6), 258-264.

Zhou, P.H., Tan, Y.F., He, Y.Q., Xu, C.G., & Zhang, Q. (2003). Simultaneous improvement for four quality traits of Zhenshan 97, an elite parent of hybrid rice, by molecular marker-assisted selection. *Theor. Appl. Genet.*, 106, 326–331.