

TABLE OF CONTENTS

| | Page |
|---|-------------|
| TITLE PAGE | i |
| ORIGINAL LITERARY WORK DECLARATION | ii |
| ABSTRACT | iii |
| ABSTRAK | v |
| ACKNOWLEDGEMENTS | vii |
| TABLE OF CONTENTS | viii |
| LIST OF FIGURES | xiii |
| LIST OF TABLES | xv |
| ABBREVIATIONS | xvii |

CHAPTER ONE: INTRODUCTION AND LITERATURE REVIEW

| | | |
|-----|--|---|
| 1.1 | Introduction | 1 |
| | 1.1.1 Objective of this study | 3 |
| 1.2 | Dengue virus (DENVs) | 5 |
| | 1.2.1 Geographic and Seasonal Distribution | 7 |
| | 1.2.2 Taxonomy and Classification | 9 |

| | |
|---|----|
| 1.2.3 Morphology of dengue virus | 10 |
| 1.2.4 Strains of dengue virus | 12 |
| 1.2.4.1 Structural proteins | 14 |
| 1.2.4.2 Non-structural proteins | 15 |
| 1.2.5 Immune Response | 17 |
| 1.2.6 Prevention and Control of dengue | 19 |
| 1.2.7 Dengue Virus Challenges | 20 |
| 1.3 Plant under study | 21 |
| 1.3.1 <i>Quercus Infectoria</i> | 22 |
| 1.3.1.1 Distribution and Description | 22 |
| 1.3.1.2 History | 25 |
| 1.3.1.3 Chemical Composition | 25 |
| 1.3.1.4 Action and Medicinal Uses | 26 |
| 1.4 Natural products evolution | 27 |
| 1.4.1 Ellagic Acid | 29 |
| 1.4.2 Gallic Acid | 30 |
| CHAPTER TWO: MATERIALS AND METHODS | |
| 2.1 Materials | 31 |
| 2.2 Chemical and Biological Agents | 31 |

| | |
|---|----|
| 2.3 Escherichia coli | 32 |
| 2.4 Media for Bacterial Cell Growth | 32 |
| 2.5 Antibiotic and Solution for Bacterial Culture | 33 |
| 2.6 Reagents and Buffers for SDS-PAGE | 34 |
| 2.7 Buffers and Reagents for western Blot | 37 |
| 2.8 Prestained SDS-PAGE Standards Protein Marker | 38 |
| 2.9 Bradford Reagent for Protein Estimation (Bio-Rad) | 38 |
| 2.10 Buffers for Protein Purification | 38 |
| 2.11 Buffer for Protease Assay | 39 |
| 2.12 Instruments | 40 |
| 2.13 PHYTOCHEMICAL ANALYSIS WORK | 40 |
| 2.13.1 Crude Plant Extract | 41 |
| 2.13.2 Fractionation by Column Chromatography | 41 |
| 2.13.3 Thin-Layer Chromatography (TLC) | 42 |
| 2.13.4 High-performance liquid chromatography (HPLC) | 42 |
| 2.13.5 Nuclear Magnetic Resonance (NMR) | 43 |
| 2.14 Biological work | 43 |
| 2.14.1 Transformation of E.coli | 43 |
| 2.14.2 Expression and Purification of Recombinant Protein | 44 |
| 2.14.3 Protein Estimation | 45 |
| 2.14.3.1 Bovine serum albumin (BSA) standard curve | 45 |

| | |
|--|----|
| 2.14.3.2 Protein Estimation Using Bradford Reagent | 45 |
| 2.15 7-Amino-4-methylcoumarin (AMC) Standard plot for protein assay | 46 |
| 2.16 One Dimension Sodium Dodecyl Sulphate – Polyacrylamide Gel Electrophoresis (SDS PAGE) | 46 |
| 2.17 Western blot | 48 |
| 2.18 Preparation of Fluorogenic Peptide Substrate | 49 |
| 2.19 Expression and Purification of Dengue-2 complex | 49 |
| 2.20 Determination of Optimum Activity of the Den2 NS2B/NS3 Protease and Substrate | 50 |
| 2.21 Determination of Solvent Effect on Enzyme Activity (Methanol and DMSO) | 51 |
| 2.22 Inhibition Assays Using Fluorogenic Peptide | 51 |
| 2.23 Inhibition Assay by Using Tecan Infinite M200 Pro fluorescence spectrophotometer | 51 |
| 2.24 Determination of NS2B/NS3 Protease Biochemical and Kinetic Properties | 52 |
| 2.24.1 Analysis of Kinetics Parameter, K_m and V_{max} | 52 |
| 2.24.2 Determination of the Inhibition Constant, K_i | 53 |
| 2.25 Calculation Methods | 53 |
| 2.25.1 Calculation of R_f Values | 53 |
| 2.25.2 Calculation of Inhibition Activity | 54 |
| 2.26 Cytotoxicity assay | 55 |

CHAPTER THREE: RESULTS

| | |
|--|----|
| 3.1 Expression and purification of DEN-2 complex | 56 |
| 3.2 Western Blot | 59 |
| 3.3 Determination of Enzyme Concentration | 60 |
| 3.4 Determination of 7-amino-4-methylcoumarin (AMC) Released | 62 |
| 3.5 Determination of Optimum Enzyme Concentration for Assay | 64 |
| 3.6 Determination of Optimum Substrate Concentration for Assay | 66 |
| 3.7 Biochemical and Kinetic Parameters of the DEN-2 Protease Complex | 66 |
| 3.8 Effect of solvents on DEN-2 virus protease complex | 68 |
| 3.9 Initial screening of <i>Quercus Infectoria</i> aqueous extract and partition fractions against DEN-2 NS2B/NS3 Protease complex | 70 |
| 3.10 <i>Quercus Infectoria</i> | 73 |
| 3.10.1 Fractionation and purification of <i>Quercus Infectoria</i> | 73 |
| 3.10.2 <i>Quercus Infectoria</i> fractions activity towards NS2B/NS3 Protease | 75 |
| 3.10.3 High-Performance Liquid Chromatography of Active Fractions | 75 |
| 3.10.4 Nuclear Magnetic Resonance (NMR) of Purified compounds | 83 |
| 3.10.5 Inhibition Assay of Purified compounds of <i>Quercus Infectoria</i> aqueous extract on DEN-2 NS2B/NS3 Protease Complexes | 89 |
| 3.11 Kinetic Analysis to Determine Type of Inhibition | 91 |
| 3.12 Purified compounds of <i>Quercus Infectoria</i> | 91 |

| | |
|---|-----|
| 3.13 Cytotoxicity assay | 96 |
| CHAPTER FOUR: DISCUSSION | |
| 4.1 Discussion | 100 |
| CHAPTER FIVE: | |
| 5.1 Conclusion | 112 |
| REFERENCES | 113 |
| LIST OF FIGURES | |
| Figure 1.1 Mosquito Vectors of Dengue | 1 |
| Figure 1.2 The global distribution of the predominant vector for dengue and regions endemic with dengue activity (Adapted from World Health Organization, 2011) | 8 |
| Figure 1.3 Diagram of a Flavivirus Virion | 11 |
| Figure: 1.4 Diagram of the Flaviviral Genome | 13 |
| Figure 1.5 <i>Quercus infectoria</i> (Manjakani) | 24 |
| Figure 1.6 Structure of Ellagic Acid | 29 |
| Figure 1.7 Structure of Gallic Acid | 30 |
| Figure 3.1 Elution profile of protein purification of DEN-2 protease precursor fraction on Nickle-column | 57 |

| | |
|--|----|
| Figure 3.2 SDS-PAGE of NS2B/NS3 before and after purification | 58 |
| Figure 3.3 Western blot of NS2B/NS3 elution after SDS-PAGE electrophoresis apparatus | 59 |
| Figure 3.4 Bovine serum Albumin (BSA) standard curve was used to determination of concentration of the protein | 61 |
| Figure 3.5 7-amino-4-methylcounarin (AMC) standard curve | 63 |
| Figure 3.6 Graph of enzyme optimum by GraphPad Prism 5.0 software | 65 |
| Figure 3.7 Graph of substrate optimum of NS2B/NS3 by Graphpad 5.0 software | 67 |
| Figure 3.8 Chart of Inhibition assay with <i>Quercus Infectoria</i> and 5 fractions from column chromatography towards NS2B/NS3 | 72 |
| Figure 3.9 Thin layer chromatography of different fractions isolated by column chromatography | 74 |
| Figure 3.10 The chromatogram of fraction 1 | 76 |
| Figure 3.11 The chromatogram of fraction 2 | 77 |
| Figure 3.12 The chromatogram of fraction 3 | 78 |
| Figure 3.13 The chromatogram of fraction 4 | 79 |
| Figure 3.14 The chromatogram of fraction 5 | 80 |
| Figure 3.15 Chromatogram of the active compound 1 (Ellagic acid) | 81 |
| Figure 3.16 Chromatogram of the active compound 2 (Gallic acid) | 82 |
| Figure 3.17 ¹ H NMR Spectrum of the active compound 1 of aqueous extract of <i>Quercus Infectoria</i> after silica gel chromatography | 84 |

| | |
|---|----|
| Figure 3.18 ^1H NMR Spectrum of the active compound 2 of aqueous extract of <i>Quercus Infectoria</i> after silica gel chromatography | 85 |
| Figure 3.19 ^{13}C NMR Spectrum of the active compound 1 of aqueous extract of <i>Quercus Infectoria</i> after silica gel chromatography | 86 |
| Figure 3.20 ^{13}C NMR Spectrum of the active compound 2 of aqueous extract of <i>Quercus Infectoria</i> after silica gel chromatography | 87 |
| Figure 3.21 Chart of inhibition assay purified compounds on DEN 2 NS2B/NS3 protease complex | 90 |
| Figure 3.22 Kinetic assay plot of ellagic acid from <i>Quercus Infectoria</i> | 92 |
| Figure 3.23 Kinetic assay plot of gallic acid from <i>Quercus Infectoria</i> | 94 |
| Figure 3.24 The cytotoxicity effect of <i>Quercus Infectoria</i> crude extract on MK2 cells. | 97 |
| Figure 3.25 The cytotoxicity effect of purified ellagic acid on MK2 cells | 98 |
| Figure 3.26 The cytotoxicity effect of purified gallic acid on MK2 cells | 99 |

LIST OF TABLES

| | |
|--|----|
| Table 2.1 Preparation of 12% (w/v) polyacrylamide gel and 5% (w/v) stacking gel | 47 |
| Table 3.1 The optimum enzyme concentration V_{max} and K_m by GraphPad Prism 5.0 software | 65 |
| Table 3.2 Kinetic properties of NS2B/NS3 protease complex parameter by GraphPad Prism 5.0 software | 67 |

| | |
|--|----|
| Table 3.3 Determination of inhibitory effect on the active enzyme and substrate against DMSO as solvent at different concentrations range from 0% to 50% | 69 |
| Table 3.4 Determination of inhibitory effect on the active enzyme and substrate against methanol as solvent at different concentrations range from 0% to 50% | 69 |
| Table 3.5 Inhibition assay with <i>Quercus Infectoria</i> and 5 fractions from column chromatography | 71 |
| Table 3.6 Table showing the retention time of the active fraction 1 | 76 |
| Table 3.7 The table showed the retention time of the active fraction 2 | 77 |
| Table 3.8 The table showed the retention time of the active fraction 3 | 78 |
| Table 3.9 The table showed the retention time of the active fraction 4 | 79 |
| Table 3.10 The table showed the retention time of the active fraction 5 | 80 |
| Table 3.11 ¹³ C and ¹ H NMR spectral data of the Ellagic acid and Gallic acid | 88 |
| Table 3.12 Result of inhibition assay of purified plant extracts on DEN-2 NS2B/NS3 protease complex | 89 |
| Table 3.13 The values of K_i , α , V_{max} and K_m by GraphPad Prism 5.0 software | 93 |
| Table 3.14 The values of K_i , α , V_{max} and K_m by GraphPad Prism 5.0 software | 95 |