

REFERENCES

- Adom, K. K. & Liu, R. H. (2002). Antioxidant activity of grains. *Journal of Agricultural and Food Chemistry*, 50 (21), 6182-6187.
- Ali, B.H., Mousa, H.M., & El-Mougy, S. (2003). The effect of a water extract and anthocyanins of *Hibiscus sabdariffa* L. on paracetamol-induced hepatotoxicity in rats. *Phytotherapy Research*, 17(1), 56-59.
- Anderson, T.J., Meredith, I.T., Yeung, A.C., Frei, B., Selwyn, A.P., & Ganz, P. (1995). The effect of cholesterol-lowering and antioxidant therapy on the endothelium-dependent coronary vasomotion. *New England Journal of Medicine*, 332, 488-493.
- Anderson, K. M., Wilson, P. W., Odell, P. M., & Kannel, W. B. (1991). An updated coronary risk profile:
A statement for health professionals. *Circulation*, 83, 356-362.
- Antonio, M.G.Jr. (2002). High-Density lipoprotein cholesterol and triglycerides as therapeutic target for preventing and treating coronary artery disease. *American Heart Journal*, 144(6), 33-42.
- Badreldin H. Ali, Naser Al Wabel, & Gerald Blunden. (2005). Phytochemical, Pharmacological and Toxicological Aspects of *Hibiscus sabdariffa* L.: A Review. *Phytotherapy Research*, 19, 369-375.
- Beltrán-Debón, R., Alonso-Villaverde, C., Aragonès, G., Rodríguez-Medina, I., Rull, A., Micol, V. (2010). The aqueous extract of *Hibiscus sabdariffa* calices modulates the production of monocyte chemoattractant protein-1 in humans. *Phytomedicine*, 17, 186-191.

- Bloomfield, R.H., Davenport, J., Babikian, V., Brass, L.M., Collins, D., Wexler, L. (2001). VA-HIT Study Group. Reduction in stroke with gemfibrozil in men with coronary heart disease and low HDL cholesterol: the Veterans Affairs HDL Intervention Trial (VA-HIT). *Circulation*, *103*, 2828-2833.
- Brown, M.S., & Goldstein, J.L. (1980). Multivalent feedback regulation of HMG Co-A reductase, a control mechanism coordinating isoprenoid synthesis and cell growth. *Journal of Lipid Research*, *21*, 505-517.
- Chait, A., Brazg, R.L., Tribble, D.L., & Krauss, R.M. (1993). Susceptibility of small, dense, low-density lipoproteins to oxidative modification in subjects with the atherogenic phenotype, pattern-B. *American Journal of Medicine*, *94*, 350-356.
- Chang, Y.C., Huang, H.P., Hsu, J.D., Yang, S.F., & Wang, C.J. (2005). Hibiscus anthocyanins rich extract-induced apoptotic cell death in human promyelocytic leukemia cells. *Toxicology and Applied Pharmacology*, *205*, 201- 212.
- Chen, C.C., Chou, F.P., Ho, Y.C., Lin, W.L., Wang, C.P., Hao, E.S. (2004). Inhibitory effects of *Hibiscus sabdariffa* L., extract on low-density lipoprotein oxidation and anti-hyperlipidemia in fructose-fed and cholesterol-fed rats. *Journal of the Science of Food and Agriculture*, *84*, 1989- 1996.
- Chen, C.C., Hsu, J.D., Wang, S.F., Chiang, H.C., Yang, M.Y., Kao, E.S., Ho, Y.C., Wang, C.J., 2003. Hibiscus sabdariffa extract inhibits the development of atherosclerosis in cholesterol-fed rabbits. *Journal of Agricultural and Food Chemistry*, *51*, 5472-5477.
- Chewonarin, T., Kinouchi, T., Kataoka, K., Arimochi, H., Kuwahara, T., Vinitketkumnun, U. (1999). Effects of roselle (*Hibiscus sabdariffa* Linn.), a Thai medicinal plant, on the mutagenicity of various known mutagens in *Salmonella typhimurium* and on formation of aberrant crypt foci induced by the colon carcinogens azoxymethane and 2-amino-1-methyl-6-phenylimidazo(4,5b)pyridine in F344 rats. *Food Chemistry and Toxicology*, *37*, 591-601.

- Chiang, J.Y. (2003). Bile acid regulation of hepatic physiology: III. Bile acids and nuclear receptors. *American Journal of Physiology - Gastrointestinal and Liver Physiology*, 284, 349-356.
- Christian, K.R., Nair, M.G., & Jackson, J.C. (2006). Antioxidant and cyclooxygenase inhibitory activity of sorrel (*Hibiscus sabdariffa*). *Journal of Food Composition and Analysis*, 19, 778–783.
- Chrisselis, M.C. (2000). Hypocholesterolemic and hypolipidemic activities of some novel morphine derivatives with antioxidant activity. *Journal of Medicinal Chemistry*, 43, 609-612.
- Crawford, R.S., Kirk, E.A., Rosenfeld, M.E., LeBoeuf, R.C., & Chait, A. (1998). Dietary antioxidants inhibit development of fatty streak lesions in the LDL receptor-deficient mouse. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 18, 1506-1513.
- Dafallah, A.A., & Al-Mustafa, Z. (1996). Investigation of the anti-inflammatory activity of *Acacia nilotica* and *Hibiscus sabdariffa*. *American Journal of Chinese Medicine*, 24, 263-269.
- Diaz, M.N., Frei, B., Vita, J.A., & Keaney, J.F. (1997). Antioxidants and atherosclerosis heart disease. *New England Journal of Medicine*, 37, 408-416.
- Dinis, T.C., Madeira, V.M., & Almeida, L.M. (1994). Action of phenolic derivatives (acetoaminophen, salicylate and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxyl radical scavengers. *Archives Biochemistry and Biophysics*, 315, 161-169.
- Droge, W. (2002). Free radicals in the physiological control of cell function. *Physiological Reviews*, 82, 47-95.

- Eklund, P.C., Langvik, O.K., Warna, J.P., Salmi, T.O., Willfor, S.M., & Sjöholm, R.E. (2005). Chemical studies on antioxidant mechanisms and free radical scavenging properties of lignans. *Organic and Biomolecular Chemistry*, 21, 3336–3347.
- El-Saadany, S.S., Sitohy, M.Z., & Labib, S.M. (1991). Biochemical dynamic and hypocholesterolemic action of *Hibiscus sabdariffa* L. (Karkada). *Die Nahrung*, 35, 567-576.
- El-Shayeb, N.M., & Mabrook, S.S. (1984). Utilization of some edible and medicinal plants to inhibit aflatoxin formation. *Nutrition reports international*, 29, 273-282.
- Farnworth, N.R., & Bunyapraphatsara, N. (1992). Thai Medicinal Plants. (pp. 163-166). Bangkok: Prachachon Press.
- Fernández-Arroyo, S., Rodríguez-Medina, I.C., Beltrán-Debón, R., Joven, F.P.J., Micol, V., & Segura-Carretero, A. (2011). Quantification of the polyphenolic fraction and in vitro antioxidant and in vivo anti-hyperlipemic activities of *Hibiscus sabdariffa* aqueous extract. *Food Research International*, 44, 1490-1495.
- Gabay, C., & Kushner, I. (1999). Acute-phase proteins and other systemic response to inflammation. *New England Journal of Medicine*, 340, 448-454.
- Gangrade, H., Mishra, S.H., & Kaushal, R. (1979). Antimicrobial activity of the oil and unsaponifiable matter of red roselle. *Indian Drugs*, 16, 147-148.
- Goodwin, B., Jones, S. A., Price, R. R., Watson, M. A., McKee, D. D., Moore, L.B. (2000). A regulatory cascade of the nuclear receptors FXR, SHP-1, and LRH-1 represses bile acid biosynthesis. *Molecular Cell*, 6, 517-526.
- Graaf, J., Hendriks, J.C.M., Demacker, P.N.M., & Stalenhoef, A.F.H. (1991). Enhanced susceptibility to in vitro oxidation of the dense low density lipoprotein subfraction in healthy subjects. *Arteriosclerosis Thrombosis and Vascular Biology*, 11, 298-306.

- Guerin, J.C., & Reveillere, H.P. (1984). Antifungal activity of plant extracts used in therapy. 1. Study of 41 plant extracts against 9 fungi species. *Annales Pharmaceutiques Francaises*, 42, 553-559.
- Halliwell, B. (1996). Antioxidants in human health and disease. *Annual Review of Nutrition*, 16, 33-50.
- Halliwell, B., & Gutteridge, J.M.C. (1984). Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochemistry Journal*, 219, 1-14.
- Haslinger, B., Kleemann, R., Toet, K.H., & Kooistra, T. (2003). Simvastatin suppresses tissue factor expression and increases fibrinolytic activity in tumor necrosis factor-alpha-activated human peritoneal mesothelial cells. *Kidney International*, 63, 2065-2074.
- Herrera-Arellano, A., Flores-Romero, S., Chavez-Soto, M. A., & Tortoriello, J. (2004). Effectiveness and tolerability of a standardized extract from *Hibiscus sabdariffa* in patients with mild to moderate hypertension: A controlled and randomized clinical trial. *Phytomedicine*, 11, 375-382.
- Hirunpanich, V., Utaipat, A., Morales, N.P, Bunyaphatsara, N., Sato, H., Herunsalee, A., et al. (2005) Antioxidant effects of aqueous extracts from dried calyx of *Hibiscus sabdariffa* Linn. (Roselle) in vitro using rat low-density lipoprotein (LDL). *Biological & Pharmaceutical Bulletin*, 28, 481-484.
- Hirunpanich, V., Utaipat, A., Morales, N.P., Bunyaphatsara, N., Sato, H., Herunsalee, A., et. al. (2006). Hypocholesterolemic and antioxidant effects of aqueous extracts from the dried calyx of *Hibiscus sabdariffa* L. in hypercholesterolemic rats. *Journal of Ethnopharmacology*, 103, 252-260.
- Hirunpanich, V., Utaipat, A., Morales, N.P., Bunyaphatsara, N., Sato, H., Herunsalee, A. (2005). Antioxidant effects of aqueous extracts from dried calyx of *Hibiscus sabdariffa* linn. (Roselle) in vitro using rat low-density lipoprotein (LDL). *Biological and Pharmaceutical Bulletin*, 28, 481-484.

- Humphries, S.E., Lane, A., Green, F.R., Cooper, J., Miller, G.J. (1994). Factor VII coagulant activity and antigen concentrations in healthy men are determined by interaction between factor VII genotype and plasma triglyceride concentration. *Arteriosclerosis and Thrombosis*, 14, 193-198.
- Ishige, K., Schubert, D., & Sagara, Y. (2001). Flavonoids protect neuronal cells from oxidant stress by three distinct mechanisms. *Free Radical Biology and Medicine*, 30, 433-446.
- James, M.S. (1994). *Roselle — Hibiscus sabdariffa* L. Institute of Food and Agricultural Sciences (IFAS).
- Jialal, I., Vega, G.L., & Grundy, S.M. (1990). Physiologic levels of ascorbate inhibit the oxidative modification of low density lipoprotein. *Atherosclerosis*, 82, 185-191.
- Juliani, H.R., Welch, C.R., Wu, Q., Diouf, B., Malainy, D., & Simon, J.E. (2009). Chemistry and quality of hibiscus (*Hibiscus sabdariffa*) for developing the natural-product industry in Senegal. *Journal of Food Science*, 74 (2), 113–121.
- Khafaga, E.S.R., & Koch, H. (1980). Stage of maturity and quality of roselle (*Hibiscus sabdariffa* L. var. *sabdariffa*): Organic acids. *Angewandte Botanik*, 54, 287–293.
- Kim, H.K., & Verpoorte, R. (2004). Sample preparation for plant metabolomics. *Phytochemical analysis*, 21(1), 4-13.
- Kirdpon, S., Nakorn, S.N., & Kirdpon, W. (1994). Changes in urinary chemical composition in healthy volunteers after consuming roselle (*Hibiscus sabdariffa* Linn.) juice. *Journal of Medical Association of Thailand*, 77, 314–321.
- Kleemann, R., & Kooistra, T. (2005). HMG-CoA Reductase Inhibitors: Effects on Chronic Subacute Inflammation and Onset of Atherosclerosis Induced by Dietary Cholesterol. *Current Drug Targets - Cardiovascular & Haematological Disorders*, 5, 441-453.

- Koivisto, U.M., Hamalainen, L., Taskinen, M.R., Kettunen, K., & Kontula, K. (1993). Prevalence of familial hypercholesterolemia among young North Karelian patients with coronary heart disease: a study based on diagnosis by polymerase chain reaction. *Journal of Lipid Research*, *34*, 269-277.
- Lazze, M.C., Pizzala, R., Savio, M., Stivala, L., Prosperi, E., & Bianchi, L. (2003). Anthocyanins protect against DNA damage induced by tert-butyl-hydroperoxide in rat smooth muscle and hepatoma cells. *Mutation Research*, *535*(1), 103-115.
- Lee, H.L. & Halliwell, B. (2001). Antioxidant and prooxidant abilities of foods and beverages. *Methods in Enzymology*, *335*, 181-190.
- Lee, J., Koo, N., & Min, D.B. (2004). Reactive oxygen species, aging and antioxidative nutraceuticals. *Comparative Reviews of Food Science*, *3*, 21-33.
- Lee, M.J., Chou, F.P., Tseng, T.H., Hsieh, M.H., Lin, M.C., & Wang, C.J. (2002). Hibiscus protocatechuic acid or esculetin can inhibit oxidative LDL induced by either copper ion or nitric oxide donor. *Journal Agriculture and Food Chemistry*, *50*, 2130- 2136.
- Liyana-Pathirana, C., & Shahidi, F. (2004). Antioxidant activity of wheat extracts as effected by *in vitro* digestion. *Biofactors*, *21*(1-4), 325-328.
- Lin, H.H., Huang, H.P., Huang, C.C., Chen, J.H., & Wang, C.J. (2005). Hibiscus polyphenol-rich extract induces apoptosis in human gastric carcinoma cells via p53 phosphorylation and p38 MAPK/FasL cascade pathway. *Molecular Carcinogenesis*, *43*, 86-99.
- Lin, T.L., Lin, H.H., Chen, C.C., Lin, M.C., Chou, M.C., Wang, C.J. (2007). *Hibiscus sabdariffa* extract reduces serum cholesterol in men and women. *Nutrition Research*, *27*, 140–145.
- Liu, C.L., Wang, J.M., Chu, C.Y., Cheng, M.T., & Tseng, T.H. (2002). In vivo protective effect of protocatechuic acid on tert-butyl hydroperoxide-induced rat hepatotoxicity. *Chemical Toxicology*, *40*, 635-641.

- Liu, J. Y., Chen, C. C., Wang, W. H., Hsu, J. D., Yang, M. Y., & Wang, C. J. (2006). The protective effects of *Hibiscus sabdariffa* extract on CCl₄ -induced liver fibrosis in rats. *Food and Chemical Toxicology*, *44*, 336-343.
- Maganha, E.G., Halmenschlager, R.D.C., Rosa, R.M., Henriques, J.A.P., Ramos, A.L.L.D.P & Saffi, J. (2010). Pharmacological evidences for the extracts and secondary metabolites from plants of the genus Hibiscus. *Food Chemistry*, *118*, 1-10.
- Maxfield, F.R., Wustner, D. (2002). Intracellular cholesterol transport. *Journal of Clinical Investigation*, *110*, 891-898.
- McKenna, N.J., & O'Malley, B.W. (2002). Minireview: nuclear receptor coactivators - an update. *Endocrinology*, *143*, 2461-2465.
- McLaughlin, J.L., Rogers, L.L & Anderson, J.E. (1998). The use of biological assays to evaluate botanicals. *Drug Information Journal*, *32*, 513-524.
- Mierzejewski, A. (2004). *Bring Your Triglycerides Down Naturally: A Drug-Free Solution to High Blood Lipids*. Full of health inc. ISBN 0-9736799-0-5
- Miller, H.E., Rigelhof, F., Marquart, L., Prakash, A., & Kanter, M. (2000). Antioxidant content of whole grain breakfast cereals, fruits and vegetables, *Cereal Foods World*, *45*(2), 59-63.
- Miller, M., Seidler, A., Moalemi, A., & Pearson, T.A. (1998). Normal triglyceride concentrations and coronary artery disease events: the Baltimore Coronary Observational Long-Term Study. *Journal of the American College of Cardiology*, *31*, 1252– 1257.
- Milletti, M., Dore, F., & Palmieri, S. (1959). Paper chromatography of extracts of *Hibiscus sabdariffa*: Identification of the coloring matter. *Annali Di Chimica* (Rome), *49*, 655–662.

- Muller, B.M., & Franz, G. (1992). Chemical structure and biological activity of polysaccharides from *Hibiscus sabdariffa*. *Planta Medica*, 58, 60–67.
- National Cholesterol Education Program (NCEP)(2001). Executive Summary of The Third Report of The NCEP Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *Journal of the American Medical Association*, 285, 2486-2497.
- Navab, M., Berliner, J.A., Subbanagounder, G., Hama, S., Lusis, A.J., Castellani, L.W. (2001). HDL and the inflammatory response induced by LDL-derived oxidized phospholipids. *Arteriosclerosis Thrombosis and Vascular Biology*, 21, 481-488.
- Odigie, I.P., Ettarh, R.R., & Adigun, S.A. (2003). Chronic administration of aqueous extract of *Hibiscus sabdariffa* attenuates hypertension and reverses cardiac hypertrophy in 2K-1 C hypertensive rats. *Journal of Ethnopharmacology*, 86, 181-185.
- Onyenekwe, P.C., Ajani, E.O., Ameh, D.A., Gamaniel, K.S. (1999). Antihypertensive effect of roselle (*Hibiscus sabdariffa*) calyx infusion inspontaneously hypertensive rats and a comparison of its toxicity with that in Wistar rats. *Cell Biochemistry and Function*, 17,199-206.
- Orisakwe, O.E., Husaini, D.C., & Afonne, O.J. (2004). Testicular effects of sub-chronic administration of *Hibiscus sabdariffa* calyx aqueous extract in rats. *Reproductive Toxicology*, 18, 295-298.
- Osman, M., Golam, F., Saberi, S., Majid, N.A., Nagoor, N.A., Zulqarnain, M. (2011). Morpho-agronomic analysis of three roselle (*Hibiscus sabdariffa* L.) mutants in tropical Malaysia. *Australian Journal of Crop Sciences*, 5(10), 1150-1156.
- Oyaizu, M. (1986). Studies on products of browning reaction prepared from glucosamine. *Japanese Journal of Nutrition*, 44, 307-314.

- Prenesti, E., Berto, S., Daniele, P.G., & Toso, S. (2007). Antioxidant power quantification of decoction and cold infusions of *Hibiscus sabdariffa* flowers. *Food Chemistry*, *100*, 433-438.
- Prior, R.L., Wu, X., & Schaich, K. (2005) Standardized Methods for the Determination of Antioxidant Capacity and Phenolics in Foods and Dietary Supplements. *Journal of Agricultural and Food Chemistry*, *53*, 4290-4302.
- Pullinger, C.R., Eng, C., Salen, G., Shefer, S., Batta, A.K., Erickson, S.K. (2002). Human cholesterol 7alpha-hydroxylase (CYP7A1) deficiency has a hypercholesterolemic phenotype. *Journal of Clinical Investigation*, *110*, 109-117.
- Queiroz, O. (1974). Circadian-rhythms and metabolic patterns. *Annual Review of Plant Physiology and Plant Molecular Biology*, *25*, 115-134.
- Reddy, S.T., Wadleigh, D.J., Grijalva, V., Ng, C., Hama, S., Gangopadhyay, A., et al. (2001). Human paraoxonase-3 is an HDL-associated enzyme with biological activity similar to paraoxonase-1 protein but is not regulated by oxidized lipids. *Arteriosclerosis Thrombosis and Vascular Biology*, *21*, 542-547.
- Ridker, P.M. (2003). Rosuvastatin in the primary prevention of cardiovascular disease among patients with low levels of low-density lipoprotein cholesterol and elevated high-sensitivity C-reactive protein: rationale and design of the JUPITER trial. *Circulation*, *108*, 2292-2297.
- Rieser, M.J., Gu, Z.M., Fang, X.P., Zeng, L., Wood, K. V., & McLaughlin, J. L. (1996). Five novel mono-tetrahydrofuran ring acetogenins from the seeds of *Annona muricata*. *Journal of Natural Products*, *59*, 100-108.
- Rimm, E.B., & Stamfer, M.J. (2000). Antioxidants for vascular disease. *Medical Clinics of North America*, *84*, 239-249.

- Russell, D.W. (2009). Fifty years of advances in bile acid synthesis and metabolism. *Journal of Lipid Research*, 50, 120-125.
- Salah, A.M., Gathumbi, J., & Vierling, W. (2002). Inhibition of intestinal motility by methanolic extracts of *Hibiscus sabdariffa* L. (Malvaceae) in rats. *Phytotherapy Research*, 16, 283–285.
- Seshadri, T.R., & Thakur, R.S. (1961). Constitution of gossytrin, a new glycoside from the flower petals of *Hibiscus sabdariffa*. *Journal of Indian Chemistry Society*, 38, 649-651.
- Shahidi, F. (2000). Antioxidants in food and food antioxidants. *Nahrung*, 44, 158-163.
- Shibata, M., & Furukawa, M. (1969). Reexamination of the structure of the so-called hiverscin. *Botanical Magazine Tokyo*, 82, 341–347.
- Singaraja, R.R., Van Eck, M., Bissada, N., Zimetti, F., Collins, H.L., Hildebrand, R.B., et al. (2006). Both hepatic and extrahepatic ABCA1 have discrete and essential functions in the maintenance of plasma high-density lipoprotein cholesterol levels in vivo. *Circulation*, 114, 1301-1309.
- Stampfer, M.J., Krauss, R.M., Ma, J., Blanche, P.J., Holl, L.G., Sacks, F.M. (1996). A prospective study of triglyceride concentration, low-density lipoprotein particle diameter, and risk of myocardial infarction. *Journal of the American Medical Association*, 276, 882-888.
- Subramanian, S.S., & Nair A.G.R. (1972). Flavonoids of four Malvaceous plants. *Phytochemistry*, 11, 1518–1519.
- Sullivan, D.R., Babiy, A.V., & Gebiciki, J.M. (1990). Vitamin E content and low density lipoprotein oxidizability induced by free radicals. *Atherosclerosis*, 81, 175-182.

- Takeda, N., & Yasui, Y. (1985). Identification of mutagenic substances in roselle color, elderberry color and safflower yellow. *Agricultural and Biological Chemistry*, *49*, 1851–1852.
- Tangirala, R.K., Casanada, F., Miller, E., Witztum, J.L., Steinberg, D., & Palinski, W. (1995). Effect of the antioxidant N,NV-diphenyl 1,4-phenylenedi- amine (DPPD) on atherosclerosis in apoE-deficient mice. *Arteriosclerosis Thrombosis and Vascular Biology*, *15*, 1625- 1630.
- Tanne, D., Koren-Morgan, N., Graff, E., & Goldbourt, U. (2001). Blood lipids and first-ever stroke/transient ischemic attack in the bezafibrate infarction prevention registry: high triglycerides consistute an independent risk factor. *Circulation*, *104*, 2892-2897.
- Teo, S., Stirling, D., Thomas, S., Hoberman, A., Kiorpes, A., & Khetani, V. (2002). A 90-day oral gavage toxicity study of d-methylphenidate and d,l-methylphenidate in Sprague-Dawley rats. *Toxicology*, *179*, 183-196.
- Timmins, J. M., Lee, J. Y., Boudyguina, E., Kluckman, K. D., Brunham, L. R., Mulya, A., et al. (2005). Targeted inactivation of hepatic Abca1 causes profound hypoalphalipoproteinemia and kidney hypercatabolism of apoA-I. *Journal of Clinical Investigation*, *115*, 1333-1342.
- Tjoelker, L.W., Eberhardt, C., Unger, J., Trong, H.L., Zimmerman, G.A., McIntyre, T.M., et al. (1995). Plasma platelet-activating factor acetylhydrolase is a secreted phospholipase A2 with acatalytic triad. *Journal of Biological Chemistry*, *270*, 25481-25487.
- Tseng, T.H., Kao, E.S., Chu, H.Y., Chou, F.P., Linwu, H.W., & Wang, C.J. (1997). Protective effects of dried flower extracts of *Hibiscus sabdariffa* L. against oxidative stress in rat primary hepatocytes. *Food and Chemical Toxicology*, *35*, 1159-1164.

- Tseng, T.H., Wang, C.J., Kao, E.S., & Chu, H.Y. (1996). Hibiscus protocatechuic acid protects against oxidative damage induced by tert-butylhydroperoxide in rat primary hepatocytes. *Chemico-Biological Interactions*, 101(2), 137–148.
- Tsuda, T., Horio, F., & Osawa, T. (2000). The role of anthocyanins as an antioxidant under oxidative stress in rats. *Biofactors*, 13, 122–133.
- Urbanczyk-Wochniak, E., Baxter, C., Kolbe, A., Kopka, J., Sweetlove, L.J., & Fernie, A.R. (2005). Profiling of diurnal patterns of metabolite and transcript abundance in potato (*Solanum tuberosum*) leaves. *Planta*, 221, 891-903.
- Vaisar, T., Pennathur, S., Green, P. S., Gharib, S. A., Hoofnagle, A. N., Cheung, M. C., et al. (2007). Shotgun proteomics implicates protease inhibition and complement activation in the antiinflammatory properties of HDL. *Journal of Clinical Investigation*, 117, 746-756.
- Van Lenten, B.J., Wagner, A.C., Nayak, D.P., Hama, S., Navab, M., & Fogelman, A.M. (2001). High-density lipoprotein lost its anti-inflammatory properties during acute influenza A infection. *Circulation*, 103, 2283-2288.
- Vilasinee, H., Anocha, U., Noppawan, P.M., Nuntavan, B., Hitoshi, S., Angkana, H. (2005). Antioxidant effects of aqueous extracts from dried calyx of *Hibiscus sabdariffa*. L (Roselle) *in vitro* using rat low density lipoprotein (LDL). *Biological and Pharmaceutical Bulletin*, 28(3), 481-484.
- Wang, C.J., Wang, J.M., Lin, W.L., Chu, C.Y., Chou, F.P., & Tseng, T.H. (2000). Protective effect of Hibiscus anthocyanins against tert-butyl hydroperoxide-induced hepatic toxicity in rats. *Food Chemistry and Toxicology*, 38(5), 411–416.
- Wang, J., Gu, Q., Li, K., & Zhang, C. (2007). CETP and oxidized LDL levels increase in dyslipidemic subjects. *Clinical Biochemistry*, 40, 995–999.

- Welson, L.T. (2006). *Triglycerides and Cholesterol Research*. Nove Science Publisher, inc. ISBN 1-60021-109-7.
- Wong, P.K., Yusof, S., Ghazali, H.M. & Che Man Y.B. (2002). Physicochemical characteristics of roselle (*Hibiscus sabdariffa* L.). *Nutrition & Food Science*, 32, 68–73.
- Xia, Y., Mirzai, B., Chackalamannil, S., Czamiecki, M., Wang, S., Clemmons, A. (1996). Substituted 1,3,5-triazines as cholesteryl ester transfer protein inhibitors. *Bioorganic & Medicinal Chemistry Letters*, 6(7), 919-922.
- Yamaguchi, F., Saito, M., Ariga, T., Yoshimura, Y., & Nakazawa, H. (2000). Free radical scavenging activity and antiulcer activity of garcinol from *Garcinia indica* fruit rind. *Journal of Agricultural and Food Chemistry*, 48, 2320-2325.
- Yamamoto, R., & Oshima, Y. (1932). Red coloring matter of *Hibiscus sabdariffa* L. (A new glucoside, hibiscin). *Scientific Papers of Institute of Physical and Chemical Research (Tokyo)*, 19, 134–141.
- Yusuf, S., Reddy, S., Ounpuu, S., & Anand, S. (2001). Global burden of cardio-vascular diseases part I: general considerations, the epidemiologic transition, risk factors, and impact of urbanization. *Circulation*, 104, 2746-2753.