

## CHAPTER 7: REFERENCES

- Abol-Munafi, A.B., Ambak, M.B., Ismail, P., MinhTam, B. (2006) Genetic Variation of Snakehead Fish (*Channa striata*) Populations Using Random Amplified Polymorphic DNA. *Biotechnology*, **5**: 104-110.
- Abol-Munafi, A.B., Ambak M.B., Ismail P., MinhTam, B. (2007) Molecular Data from the Cytochrome *b* for the Phylogeny of Channidae (*Channa* sp.) in Malaysia. *Biotechnology*, **6**: 22-27.
- Allard, M.W., Miyamoto M.M., Jareche, L., Krau F., and Tennant, M.R. (1992) DNA systematic and evolution of the artiodactyls family Bovidae. In: *Proceedings of the National Academy of Sciences USA*, **89**: 3972-3976.
- Allen, P.J., Amos, W., Pomeroy, P.P. and Twiss, S.D. (1995) Microsatellite variation in grey seals (*Halichoerus grypus*) shows evidence of genetic differentiation between two British breeding colonies. *Molecular Ecology*, **4**: 653-662.
- Avise, J.C. (1994) *Molecular Markers, Natural History and Evolution*. Chapman & Hall, New York.
- Baie, S.H., and Sheikh, K.A. (2000) The wound healing properties of *Channa striatus* cetrimide cream; tensile strength measurement. *Journal of Ethnopharmacology*, **71**: 93-100.

Balloux, F. and Lugon-Moulin, N. (2001) The estimation of population differentiation with microsatellite markers. *Molecular Ecology*, **11**: 155-165.

Bekkevold, D., Hansen, M.M. and Loeschcke, V. (2002) Male reproductive competition in spawning aggregations of cod (*Gadus morhua*, L.). *Molecular Ecology*, **11**: 91-102.

Billington, N. (2003) Mitochondrial DNA. In: E.M. Hallerman (ed.) *Population genetics: principles and applications for fisheries scientists*. Bethesda, Maryland: American Fisheries Society pp. 59-100.

Bucklin, A., Frost, B.W., and Kocher, T.D. (1995) Molecular systematic of six *Calanus* and three *Metridia* species (Calanoida: Copepoda). *Marine Biology*, **121**: 655-664.

Callen, D.F., Thompson, A.D., Shen, Y., Phillips, H., Richards, R.I., Mulley, J.C., and Sutherland, G.R. (1993) Incidence and origin of ‘null’ alleles in the (AC) $n$  microsatellite markers. *The American Journal of Human Genetics*, **52**: 922–927.

Canapa, A., Marota, I., Rollo, F. and Olmo, E., (1996) Phylogenetic analysis of veneridae (Bivalvia): Comparison of molecular and palaeontological data. *Journal of Molecular Evolution*, **43**: 517-522.

Carvalho, G.R. and Hauser, L. (1998) Advances in the molecular analysis of fish population structure. *The Italian journal of zoology (Modena)*, **65**: 21-33.

- Chambers, G.K. and MacAvoy, E.S. (2000) Microsatellites: consensus and controversy. *Comparative Biochemistry Physiology Part B Biochemistry and Molecular Biology*, **126**: 455-476.
- Çiftci, Y. and Okumus, I. (2002) Fish Population Genetics and Application of Molecular Markers to Fisheries and Aquaculture – I: Basic Principles of Fish Population Genetics. *Turkish Journal of Fisheries and Aquatic Sciences*, **2**: 145-155.
- Craig, J., Fowler, S., Burgoyne, L.A., Scott, A.C., and Harding, H.W.J. (1988) Repetitive deoxyribonucleic acid (DNA) and human genome variation: a concise review relevant to forensic biology. *Journal of Forensic Science*, **33**: 1111-1126.
- Crooijmans, R.P.M.A., Bierbooms, V.A.F., Komen, J., van der Poel, J.J., and Groenen, M.A.M. (1997) Microsatellite markers in common carp (*Cyprinus carpio* L.). *Animal Genetics*, **28**: 129–134.
- Cunningham, C.W., Blackstone, N.W. and Buss, L.W. (1992) Evolution of king crabs from hermit crab ancestors. *Nature*, **355**: 359-342.
- de Bruyn, M., Wilson, J.A. and Mather, P.B. (2004) Huxley's line demarcates extensive genetic divergence between eastern and western forms of the giant freshwater prawn, *Macrobrachium rosenbergii*. *Molecular Phylogenetics and Evolution*, **30**: 251-257.

Estroup, A., Solignac, M., Harry, M., and Cornut, J.M. (1993) Characterization of (GT)<sub>n</sub> and (CT)<sub>n</sub> microsatellites in two insect species: *Apis mellifera* and *Bombus terrestris*. *Nucleic Acids Research*, **21**: 1427–1431.

Evanno, G., Regnaut, S., Goudet, J. (2005) Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, **14**: 2611–2620.

Excoffier, L., Smouse, P.E., and Quattro, J.M. (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics*, **131**: 479-491.

Falush, D., Stephens, M., and Pritchard, J. K. (2003) Inference of population structure: Extensions to linked loci and correlated allele frequencies. *Genetics*, **164**: 1567-1587.

Falush, D., Stephens, M., and Pritchard, J. K. (2007) Inference of population structure using multilocus genotype data: dominant markers and null alleles. *Molecular Ecology Notes*, **7**: 574-578.

FAO. (1981) Conservation of the Genetic Resources of Fish: Problems and Recommendations. FAO Fisheries Technical Paper No. 217. Rome: Food and Agriculture Organisation of the United Nations.

Felsenstein, J. (2004) PHYLIP (Phylogeny Inference Package) version 3.6. *Distributed by the author. Department of Genome Sciences, University of Washington, Seattle.*

Ferguson, A., Taggart, J.B., Prodohl, P.A., McMeel, O., Thompson, C., Stone, C., McGinnity, P. and Hynes, R.A. (1995) The application of molecular markers to the study and conservation of fish populations, with special reference to *Salmo*. *Journal of Fish Biology*, **47**: 103-126.

Geller, J.B., Carlton, J.T. and Powers, D.A. (1993) Interspecific and intrapopulation variation in mitochondrial ribosomal DNA sequences of *Mytilus* sp (Bivalvia: Mollusca). *Molecular Marine Biology and Biotechnology*, **2**: 44-20.

Glaubitz, J.C. (2004) Convert: A user-friendly program to reformat diploid genotypic data for commonly used population genetic software packages. *Molecular Ecology Notes*, **4**: 309-310.

Goldstein, D.B., Linares, A.R., Cavalli-Sforza, L.L. and Feldman, M.W. (1995) An evaluation of genetic distances for use with microsatellite loci. *Genetics*, **139**: 463-471.

Hansen, M.M. (2002) Estimating the long-term effects of stocking domesticated trout into wild brown trout (*Salmo trutta*) populations: An approach using microsatellite DNA analysis of historical and contemporary samples. *Molecular Ecology*, **11**: 1003-1015.

Hansen, M.M. (2003) Application of molecular markers in population and conservation genetics, with special emphasis on fishes. DSc thesis, Faculty of Natural Sciences, University of Aarhus.

Hansen, M.M., Mensberg, K.L.D., and Berg, S. (1999) Postglacial recolonisation patterns and genetic relationships among whitefish (*Coregonus sp.*) populations in Denmark, inferred from mitochondrial DNA and microsatellite markers. *Molecular Ecology*, **8**: 239-252.

Hansen, M.M., Ruzzante, D.E., Nielsen, E.E., and Mensberg, K.L.D. (2000) Microsatellite and mitochondrial DNA polymorphism reveals lifehistory dependent interbreeding between hatchery and wild brown trout (*Salmo trutta* L.). *Molecular Ecology*, **9**: 583-594.

Hansen, M.M., Ruzzante, D.E., Nielsen, E.E. and Mensberg, K.-L.D. (2001) Brown trout (*Salmo trutta*) stocking impact assessment using microsatellite DNA markers. *Ecological Applications*, **11**: 148-160.

Hansen, M.M., Ruzzante, D.E., Nielsen, E.E., Bekkevold, D., and Mensberg, K.-L.D. (2002) Long-term effective population sizes, temporal stability of genetic composition and potential for local adaptation in anadromous brown trout (*Salmo trutta*) populations. *Molecular Ecology*, **11**: 2523– 2535.

Hartl, D.L. and Clark, A.G. (1989) *Principles of population genetics*. 2nd ed. Sinauer Associates, Sunderland, MA.

Hedrick, P.W. (2005) *Genetics of populations*. Jones and Bartlett Publishers, New York.

Hoh, B.P. (2005) Isolation, Characterization and Genetic Linkage Mapping of Microsatellite Markers in *Mystus nemurus*. Phd thesis, Universiti Putra Malaysia.

Inger, R.F. and Kong, C.P. (1962) The fresh-water fishes of North Borneo: Fieldiana: *Zoology*, **45**: 1-268.

Jackson, T.R., Ferguson, M.M., Danzmann, R.G., Fishback, A.G., Ihssen, P.E., Crease, T.J. (1998) Identification of two QTL influencing upper temperature tolerance in three rainbow trout (*Oncorhynchus mykiss*) half-sib families. *Heredity*, **80**: 143-151.

Jarne, P. and Lagoda, P.J.L. (1996) Microsatellites, form molecules to populations and back. *Trends in Ecology and Evolution*, **11**: 424–429.

Kaeuffer, R., Coltman, D.W., Chapuis, J.L., Pontier, D., Reale, D. (2007) Unexpected heterozygosity in an island mouflon population founded by a single pair of individuals. In: *Proceedings of the Royal Society B: Biological Sciences*, **274**: 527-533.

Kumar, S.V. (2003) Isolation, characterization and application of DNA microsatellite markers in mungbean (*Vigna radiata* L.Wilczek) and other selected legumes. PhD thesis, Universiti Putra Malaysia.

Lee, P.G., and Ng, P.K.L. (1991) The snakehead fishes of the Indo-Malayan Region.

*Nature Malaysiana*, **16**: 113-129.

Magoulas, A. (1998) Application of molecular markers to aquaculture and broodstock management with special emphasis on microsatellite DNA. *Cahiers Options Méditerranées*, **34**: 153- 168.

Mat Jais, A.M., McCulloch, R., and Croft, K.D. (1994) Fatty acid and amino acid composition in Haruan as potential role for wound healing. *General Pharmacology*, **25**: 947-950.

Mat Jais, A.M., Dambisya, Y.M., and Lee, T.L. (1997) Anti-nociceptive activity of *Channa striatus* (Haruan) extracts in mice. *Journal of Ethnopharmacology*, **57**: 125-130.

Mat Jais, A.M., Matori, M.F. Kittakoop, P., and Suwanborirux, K. (1998) Fatty acid compositions in mucus and roe of Haruan, *Channa striatus*, for wound healing. *General Pharmacology*, **30**: 561-563.

Mat Jais, A.M., Zakaria, Z.A., Luo, A. and Song, Y.X. (2008) Antifungal activity of *Channa striatus* (Haruan) crude extracts. *International Journal of Tropical Medicine*, **3**: 43-48.

May, B. and Krueger, C.C. (1990) Use of allozyme data for population analysis. In D.H. Whitmore (ed.) *Electrophoretic and Isoelectric Focusing Techniques in Fisheries Management*. CRC Pres, Boston. pp. 157-171.

McConnell, S.K., O'Reilly, P., Hamilton, L., Wright, J.N. and Bentzen, P. (1995) Polymorphic microsatellite loci from Atlantic salmon (*Salmo salar*) – genetic differentiation of North American and European populations. *Canadian Journal of Fisheries and Aquatic Sciences*, **52**: 1863–1872.

Menon, A.G.K. (1999) Check list - fresh water fishes of India. *Rec. Zool. Surv. India, Misc. Publ., Occas. Pap.*, **175**: 366.

Milinkovitch, M.C., Orti, G. and Meyer, A. (1993) Revised phylogeny of whales suggested by mitochondrial ribosomal DNA sequences. *Nature*, **361**: 346-348.

Mohsin, A.K.M., and Ambak, M.A. (1983) *Freshwater fishes of Peninsular Malaysia: Pertanian, Malaysia*. Penerbit Universiti. pp. 284

Nei, M. (1972) Genetic distance between populations. *American Naturalist*, **106**: 283-292.

Nei, M. (1978) Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics*, **89**: 583-590.

Neigel, J.E. (1997) A comparison of alternative strategies for estimating gene flow from genetic markers. *Annual Review of Ecology and Systematics*, **28**: 105-128.

Ng, P.K.L., and Lim, K.K.P. (1990) Snakeheads (Pisces: Channidae): Natural, history, biology and economic importance: Essays in Zoology, Papers Commemorating the 40th Anniversary of the Department of Zoology, National University of Singapore, pp. 127-152.

Nielsen, E.E., Hansen, M.M., and Loeschke, V. (1997) Analysis of microsatellite DNA from old scale samples of Atlantic salmon: A comparison of genetic composition over sixty years. *Molecular Ecology*, **6**: 487-492.

Nielsen, E.E., Hansen, M.M., and Loeschke, V. (1999) Genetic variation in time and space: Microsatellite analysis of extinct and extant populations of Atlantic salmon. *Evolution*, **53**: 261-268.

Norainy, M.H. (2007) Morphological and Genetic Variability of Malaysian Channa spp. based on Morphometric and RAPD techniques. MSc thesis, University Science Malaysia.

Nsengimana, J., Baret, P., Haley, C.S. and Visscher, P.M. (2004) Linkage disequilibrium in the domesticated pig. *Genetics*, **166**: 1395–1404.

O'Connell, M. and Wright, J.M. (1997) Microsatellite DNA in fishes. *Reviews in Fish Biology and Fisheries*, **7**: 331-363.

O'Reilly, P. and Wright, J.M. (1995) The evolving technology of DNA fingerprinting and its application to fisheries and aquaculture. *Journal of Fish Biology*, **47**: 29-55.

O'Reilly, P.T., Herbiner, C. and Wright, J.M. (1998) Analysis of parentage determination in Atlantic salmon (*Salmo salar*) using microsatellites. *Animal Genetics*, **29**: 363-370.

Ohta, T. (1982a) Linkage disequilibrium with the island model. *Genetics*, **101**: 139-155.

Ohta, T. (1982b) Linkage disequilibrium due to random genetic drift in finite subdivided populations. *Proceedings of the National Academy of Sciences*, **76**: 1940-1944.

Orti, G., Pearse, D.E. and Avise, J.C. (1997) Phylogenetic assessment of length variation at a microsatellite locus. *Proceedings of the National Academy of Sciences*, **94**: 10745-10749.

Pemberton, J.M., Slate, J., Bancroft, D.R. and Barrett, J.A. (1995) Non-amplifying alleles at microsatellite loci: a caution for parentage and population studies. *Molecular Ecology*, **4**: 249-252.

Phon, M.K. (2008) Population genetics of wild stock Channa striata using long primer random amplified polymorphic DNA markers and cross amplified microsatellite primers. BSc thesis, University Putra Malaysia.

Pritchard, J. K., Stephens, M., and Donnelly, P. (2000) Inference of population structure using multilocus genotype data. *Genetics*, **155**: 945-959.

Pritchard, J. K., and Wen, W. (2003) Documentation for Structure Software, Version 2 (<http://pritch.bsd.Uchicago.edu>).

Reilly, A., Elliott, N.G., Grewe, P.M., Clabby, C., Powell, R. and Ward, R.D. (1999) Genetic differentiation between Tasmanian cultured Atlantic salmon (*Salmo salar* L.) and their ancestral Canadian population: comparison of microsatellite DNA and allozyme and mitochondrial DNA variation. *Aquaculture*, **173**: 459–469.

Rousset, F. (1997) Genetic differentiation and estimation of gene flow from F- statistics under isolation by distance. *Genetics*, **145**: 1219-1228.

Rumbak F., Freid, D.G. and Thomas, R.H. (1994) Reconstruction of phylogeny of 11 species of Littorina (Gastropoda: Littorinidae) using mitochondrial DNA sequence data. *Nautilus (suppl)*, **2**: 91-97.

Ruzzante, D.E., Hansen, M.M. and Meldrup, D. (2001) Distribution of individual inbreeding coefficients, relatedness and influence of stocking on native anadromous brown trout (*Salmo trutta*) population structure. *Molecular Ecology*, **10**: 2107-2128.

Schlöttere, C., Amos, B. and Tautz, D. (1991) Conservation of polymorphic simple sequence loci in cetacean species. *Nature*, **354**: 63-65.

Schneider, S, Roessli, D, & Excoffier, L. (2000) Arlequin ver. 2.000: A software for population genetics data analysis', Genetics and Biometry Laboratory, University of Geneva, Switzerland.

Scribner, K.T., Gust, J.R. and Fields, R.L. (1996) Isolation and characterization of novel salmon microsatellite loci: cross-species amplification and population genetic applications. *Canadian Journal of Fisheries and Aquatic Science*, **53**: 833-841.

See, L.M. (2008) Isolation and Characterization of Microsatellite loci from the giant freshwater Prawns. Msc thesis, Universiti Putra Malaysia.

Shaklee, J.B. and Bentzen, P. (1998) Genetic identification of stocks of marine fish and shellfish. *Bulletin of Marine Science*, **62**: 589-621

Skibinski, D.O.F. (1994) The potential of DNA techniques in the population and evolutionary genetics of aquatic invertebrates. In: *Proceedings of the Conference on Genetics and Evolution of Aquatic Organisms, Bangor, UK, 10–16 Sept 1992*, (Beaumont, A., ed.). Chapman and Hall, London. pp. 177-199.

Stalling, R.L. Ford, A.F., Nelson, D., Torney, D.C., Hildebrand, C.E., and Moyzis, R.K. (1999) Evolution and distribution of (GT)<sub>n</sub> repetitive sequences in mammalian genomes. *Genomics*, **10**: 807-815.

State, J. and Pemberton, J.M. (2002) Comparing molecular measures for detecting inbreeding depression. *Journal of Evolutionary Biology*, **15**: 20-31.

Tautz, D. (1989) Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Nucleic Acids Research*, **17**: 6463-6471.

Tan, M.P., Siti Azizah, M. N., Jamsari, A. F. J., Thuy, T. T. Nguyen. (2009) Genetic Structure of Snakehead, *Channa striata* (Bloch, 1793) Populations in Peninsular Malaysia based on COI gene. Unpublished.

Tessier, N., Bernatchez, L. and Wright, J.M. (1997) Population structure and impact of supportive breeding inferred from mitochondrial and microsatellite DNA analyses in land-locked Atlantic salmon *Salmo salar* L. *Molecular Ecology*, **6**: 735–750.

Tong, J., Wang, Z., Yu, X., Wu, Q. and Chu, K.H. (2002) Cross-species amplification in silver carp and bighead carp with microsatellite primers of common carp. *Molecular Ecology Notes*, **2**: 245-247.

Usmani, S., Tan, S.G., Siraj, S.S. and Yusoff, K. (2001) Isolation and characterization of microsatellite in the Southeast Asian river catfish *Mystus nemurus*. *Molecular Ecology Notes*, **1**: 264-266.

Waples, R.S. (1989) A generalized approach for estimating effective population size from temporal changes in allele frequency. *Genetics*, **121**: 379-391.

Weber, J.L. (1990) Informativeness of human (dC-dA)<sub>n</sub> (dG-dT)<sub>n</sub> polymorphisms. *Genomics*, **7**: 524-530.

Weber, J.L. and May, P.E. (1989) Abundant class of human DNA polymorphisms which can be typed using the polymerase chain reaction. *American Journal of Human Genetics*, **44**: 388-396.

Wee, K.L. (1982) Snakeheads - Their biology and culture. In: Muir, J.F., and Roberts, R.J. (eds.) *Recent advances in aquaculture*. Westview Press, Boulder, Colorado. pp. 180-213.

Weir, B.S. (1996) Intraspecific differentiation. In: Hillis, D.M., Moritz, C. and Mable, B.K. (eds) *Molecular Systematics*. 2nd ed. Sinauer, Sunderland, MA. pp. 385–405.

Weir, B.S. and Cockerham, C.C. (1984) Estimating F-statistics for the analysis of population Structure. *Evolution*, **38**: 1358-1370.

Wigginton, J.E., Cutler, D.J., and Abecasis, G.R. (2005) A note on exact tests of Hardy-Weinberg equilibrium. *The American Journal of Human Genetics*, **76**: 887-893.

Williams, J.G.K., Kubelik, A.R., Livak, K.J., Rafalski, J.A. and Tingey, S.V. (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research*, **18**: 6531-6535.

Williamson, J.H. (2001) Broodstock management for imperilled and other fishes. In: G.A. Wedemeyer (ed.) *Fish hatchery management*. 2nd ed. American Fisheries Society, Bethesda, Maryland. pp. 397-482.

Wright, S. (1951) The genetical structure of populations. *Annals of Eugenics*, **15**: 323-354.

Wright, S. (1978) *Evolution and the genetics of population, Variability among and within natural populations*. University of Chicago Press, Chicago, **4**: 79-103.

Wright, J.M. and Bentzen, P. (1994) Microsatellites: genetic markers for the future. In: G.R. Carvalho and T.J. Pitcher (eds.) *Reviews in fish biology and fisheries*. Chapman and Hall, London, **4**: 384–388.

Wu, K., Jones, R., Danneberger, L., Scolnik, P.A. (1994) Detection of microsatellite polymorphisms without cloning. *Nucleic Acids Research*, **22**: 3257-3258.

Yang, R. (1998) Estimating hierarchical F-statistics. *Evolution*, **52**: 950-956.

Zhang, D.X. and Hewitt, G.M. (2003) Nuclear DNA analyses in genetic studies of populations: practice, problems and prospects. *Molecular Ecology*, **12**: 563–584.

Zheng, W., Stacey, E. and Coffin, J. (1995) Isolation and characterization of microsatellite loci in the goldfish *Carassius auratus*. *Molecular Ecology*, **4**: 791–792.