

LIST OF TABLES

Tables		Pages
3.1	Sampling sites, location and quantity of samples	16
3.2	List of Random Amplified Microsatellites (RAMs) primers	19
3.3	PCR condition for colony PCR	22
3.4	Standard PCR profile for optimization of the primers	25
3.5	Touchdown PCR profile	26
4.1	RAMs primer and its annealing temperature, T_m	30
4.2	Optimal PCR conditions for seven microsatellite loci	39
4.3	Characterization of seven <i>Channa striata</i> microsatellite loci	40
4.4	Stuttering bands, alleles drop out, and null alleles for polymorphic loci	44
4.5	F_{IS} values for 30 individuals	45
4.6	Allele frequency	47
4.7	Optimized temperatures and product size of each primer used	48
4.8	Polymorphisms at the microsatellite loci amplified for six populations of <i>Channa striata</i>	53
4.9	Genotype frequencies of 10 microsatellite loci in six populations of <i>Channa striata</i>	57
4.10	The polymorphisms observed at the microsatellite loci	

	in six populations of <i>Channa striata</i> .	60
4.11	Number of observed and expected alleles and the value of F_{IS} for all loci.	61
4.12	Level of heterozygosity of six populations of <i>Channa striata</i>	64
4.13	Locus MFW5 shows null allele in Negeri Sembilan population.	70
4.14	Locus MFW15 shows null allele in Penang population.	70
4.15	P-value for locus pair in all populations	72
4.16	The values of genetic distance between six populations of <i>Channa striata</i>	74
4.17	The values of genetic distance between six populations of <i>Channa striata</i> (after removing null allele)	76
4.18	Amova design and results	77
4.19	Amova design and results (after removing null allele)	78
4.20	Population pairwise F_{ST} distance	79
4.21	Population pairwise F_{ST} distance (after removing null allele)	80