

---

<b>Figure 2.1</b>	Different bonding modes of carboxylate ligands to metal	28
<b>Figure 2.2</b>	Quadruple splitting of $^{119}\text{Sn}$	43
<b>Figure 3.1</b>	Electronic structure of a semiconductor	54
<b>Figure 3.2</b>	Comparison of band gap of different type of semiconductors	54
<b>Figure 3.3</b>	Oxidation, reductions by photocatalyst	55
<b>Figure 3.4</b>	General processes for the photo- oxidation mineralization of organic compounds by $\text{TiO}_2 + \text{O}_2$	63
<b>Figure 3.5</b>	Structures for different types of $\text{TiO}_2$	66
<b>Figure 3.6</b>	Doping metal on semiconductor	67
<b>Figure 3.7</b>	Coupling two semiconductors	68
<b>Figure 3.8</b>	Dye sensitizing of $\text{TiO}_2$	69
<b>Figure4.1</b>	Kinetic plot of TBT photo-degradation	82
<b>Figure4.2</b>	Variation of the concentration of TBTCl with irradiation time	83
<b>Figure4.3</b>	Variation of the concentration of TBTCl with irradiation time for different types of nano $\text{TiO}_2$	85
<b>Figure4.4</b>	Kinetic plot of TBT photo-degradation by different types of $\text{TiO}_2$	86
<b>Figure 4.5</b>	Photo-catalytic cycle of degradation of organic compounds	92
<b>Figure 5.1</b>	Structure formula of of di-nbutyl bis(2-hydroxybenzoate )tin (IV)	114
<b>Figure 5.2</b>	Molecular structure of di-nbutyl bis( 2-hydroxy benzoate tin(IV)	115
<b>Figure 5.3</b>	Structure formula of bis[1,1,3,3-tetrabutyl-1,3-bis [2- hydroxybenzoato) 1, 3-distannoxane]	117
<b>Figure 5.4</b>	Molecular structure of bis[1,1,3,3-tetrabutyl-1,3- bis[2-hydroxybenzoato) 1,3- distannoxane]	118

---

<b>Figure 5.5</b>	structure formula of 5-chloro [1-hydroxy-2-hydroxymethyl-2-(2-oxidobenzylideneamino)-3-oxidomethylpropane]dibutyltin(IV)	135
<b>Figure 5.6</b>	Molecular structure of 5-chloro [1-hydroxy-2-hydroxymethyl-2-(2-oxidobenzylideneamino)-3-oxidomethylpropane]dibutyltin(IV)	140