

LIST OF FIGURES

- Figure 2.1** Different molecular arrangement in (a) solid; (b) liquid crystal; and (c) liquid
- Figure 2.2** The nematic phase: (left) an illustration of the orientation of the molecules, and (right) an optical micrograph of the characteristic Schlieren texture
- Figure 2.3** A molecular arrangement of: (a) S_A phase; and (b) S_C phase
- Figure 2.4** An illustration of smectic A phase (*left*), and the optical texture of the phase viewed under a polarizing microscope (*right*)
- Figure 2.5** An illustration of smectic C phase (*left*) and the optical texture of the phase viewed under a polarizing microscope (*right*)
- Figure 2.6** Examples of calamitic metallomesogens
- Figure 2.7** Molecular structure of the first series of discotic liquid crystals discovered: benzene-hexa-*n*-alkanoate derivatives ($R = C_4H_9$ to C_9H_{19})
- Figure 2.8** An illustration of a molecular arrangement of N_D phase
- Figure 2.9** (a) The general structure of Col phases; (b) a representation of Col_r ; and (c) a representation of Col_h
- Figure 2.10** Examples of disc-like metallomesogens: (a) four-chain -diketonates where $R = C_n$ and $R' = C_m$; $n = 7, m = 13$; $n = m = 10, n = m = 12$; and (b) eight-chain -diketonates
- Figure 2.11** Lantern or paddle-wheel structure of $[Cu_2(CH_3COO)_4(H_2O)_2]$
- Figure 2.12** Lantern or paddle wheel structure of the dicopper(II) tetraalkanoate, stacking in the two possible ways for the formation of the columnar phase: (a) tilted to the columnar axis for the tetragonal phase; and (b) parallel to the columnar axis for the hexagonal phase
- Figure 2.13** Applications of ionic liquids

- Figure 2.14** Examples of cations of ionic liquids
- Figure 2.15** An example of metal-containing ionic liquids [12]
- Figure 2.16** General structure of polynuclear copper(II) carboxylates: **I**, stepped without additional ligands; **II**, ligands bridging two Cu; **III**, ligands bridging Cu and O
- Figure 2.17** The enol (top) and enolate (bottom) mechanisms of aldol condensation of carbonyl compounds
- Figure 2.18** Coordination of carboxylate ligand: (I) monodentate, (II) chelating, (III) bridging bidentate (a) *syn-syn*, (b) *syn-anti*, and (c) *anti-anti*
- Figure 2.19** The colour wheel
- Figure 2.20** Shapes (angular dependence functions) of the *d* orbitals
- Figure 2.21** Splitting of *d* orbitals for common stereochemistries
- Figure 2.22** Crystal-field splitting of *d* orbitals for complexes of various geometry
- Figure 2.23** TG and DTA curves of $[\text{Cu}_2(\text{C}_6\text{H}_5\text{COO})_4(\text{C}_2\text{H}_5\text{OH})_2]$
- Figure 2.24** An example of a DSC curve
- Figure 2.25** OPM of liquid crystals phase structures (a) N droplets; (b) nematic LC; (c) cholesteric LC; (d) smectic A LC; (e) hexagonal columnar liquid crystalline and (f) spherulite texture of crystalline phase of a LC
- Figure 2.26** A schematic view of the columnar mesophase of fatty acid copper(II) complexes. Each column is made of stacked dicopper tetracarboxylate units
- Figure 2.27** OPM of (a) copper(II) butanoate, showing tetragonal domains at 184.1°C; (b) typical columnar developable domains for copper(II) butanoate at 164.7°C; (c) copper(II) pentanoate, showing hexagonal homeotropic domains at 98.9°C; (d) developable domains at 118.9°C for copper(II) pentanoate; (e) copper(II) hexanoate, showing hexagonal

homeotropic domains at 135°C; and (f) developable domains for copper(II) hexanoate at 152°C

- Figure 2.28** Random orientation of magnetic moments in paramagnetic materials
- Figure 2.29** Parallel alignment of magnetic moment in ferromagnetic materials
- Figure 2.30** Alignment of magnetic moment in antiferromagnetic materials
- Figure 2.31** Alignment of magnetic moment in ferrimagnetic materials
- Figure 2.32** Cyclic voltammogram of a reversible process
- Figure 4.1** Proposed structure of **Complex 1** (K^+ ions and H_2O solvate molecules are not shown)
- Figure 4.2** FTIR spectrum of **Complex 1**
- Figure 4.3** FTIR spectrum of $CH_3(CH_2)_{14}COOH$
- Figure 4.4** FTIR spectrum of *p*- HOC_6H_4COOH
- Figure 4.5** UV-vis spectrum of **Complex 1** (solid)
- Figure 4.6** Resonance structure of **Complex 1** (only a partial structure is shown)
- Figure 4.7** UV-vis spectrum of **Complex 1** (solution)
- Figure 4.8** Thermogram of **Complex 1**
- Figure 4.9** DSC curve of **Complex 1**
- Figure 4.10** OPM of **Complex 1** at: (a) 25°C; (b) 206°C
- Figure 4.11** Cyclic voltamogram of **Complex 1**
- Figure 4.12** Proposed structural formula of $[Cu_2(p-HOC_6H_4COO)_4]$ (H_2O solvate molecules are not shown)
- Figure 4.13** FTIR spectrum of $[Cu_2(HOC_6H_4COO)_4] \cdot 8H_2O$
- Figure 4.14** UV-vis spectrum of $[Cu_2(p-HOC_6H_4COO)_4] \cdot 8H_2O$ (solution)
- Figure 4.15** Proposed structural formula of $[Cu_2(CH_3(CH_2)_{14}COO)_4]$
- Figure 4.16** FTIR spectrum of $[Cu_2(CH_3(CH_2)_{14}COO)_4]$
- Figure 4.17** UV-vis spectrum of $[Cu_2(CH_3(CH_2)_{14}COO)_4]$ (solution)

- Figure 4.18** Proposed structural formula of **Complex 2** ($\text{CH}_3\text{CH}_2\text{OH}$ solvate is not shown)
- Figure 4.19** FTIR spectrum of **Complex 2**
- Figure 4.20** UV-vis spectrum of **Complex 2** (solution)
- Figure 4.21** Thermogram of **Complex 2**
- Figure 4.22** OPM micrographs of **Complex 2** on heating: (a) 25°C ; (b) 140°C ; (c) 180°C ; and (d) 84°C
- Figure 4.23** DSC curve of **Complex 2**
- Figure 4.24** Cyclic voltamogram of **Complex 2**
- Figure 4.25** The FTIR spectrum of the product isolated from the carbonate-base-acid reaction
- Figure 4.26** The UV-Vis spectrum of the product isolated from the carbonate-base-acid reaction
- Figure 4.27** Proposed structure of **Complex 3** (H_2O solvate molecules are not shown)
- Figure 4.28** FTIR spectrum of **Complex 3**
- Figure 4.29** FTIR spectrum of *p*- $\text{H}_2\text{NC}_6\text{H}_4\text{COOH}$
- Figure 4.30** UV-vis spectrum of **Complex 3** (solid)
- Figure 4.31** UV-visible spectrum of **Complex 3** (solution)
- Figure 4.32** Thermogram of **Complex 3**
- Figure 4.33** DSC curve of **Complex 3**
- Figure 4.34** Cyclic voltammogram of **Complex 3**
- Figure 4.35** The FTIR spectrum of the product isolated from the ligand-exchange reaction
- Figure 4.36** The UV-vis spectrum of the product isolated from the ligand-exchange reaction

- Figure 4.37** The FTIR spectrum of the product isolated from the carbonate-base-acid reaction
- Figure 4.38** The UV-Vis spectrum of the product isolated from the carbonate-base-acid reaction
- Figure 4.39** Proposed structure of **Complex 4** (K^+ ion is not shown)
- Figure 4.40** FTIR spectrum of **Complex 4**
- Figure 4.41** UV-vis spectrum of **Complex 4** (solid)
- Figure 4.42** UV-vis spectrum of **Complex 4** (solution)
- Figure 4.43** TGA of **Complex 4**
- Figure 4.44** DSC of **Complex 4**
- Figure 4.45** Cyclic voltammogram of **Complex 4**
- Figure 4.46** Proposed structure of **Complex 5**
- Figure 4.47** FTIR spectrum of **Complex 5**
- Figure 4.48** UV-vis spectrum of **Complex 5** (solid)
- Figure 4.49** UV-vis spectrum of **Complex 5** (solution)
- Figure 4.50** Thermogram of **Complex 5**
- Figure 4.51** OPM of **Complex 5** on heating: (a) 147°C; (b) 161°C; (c) 208°C; and (d) 217°C
- Figure 4.52** OPM of **Complex 5** on cooling from the isotropic liquid phase: (a) 195°C; (b) 98°C
- Figure 4.53** DSC curve of **Complex 5**
- Figure 4.54** Cyclic voltammogram of **Complex 5**
- Figure 4.55** The FTIR spectrum
- Figure 4.56** The UV-vis spectrum (solution)
- Figure 4.57** FTIR spectrum of $[Cu_2(CH_3(CH_2)_{10}COO)_4]$
- Figure 4.58** UV-visible spectrum of $[Cu_2(CH_3(CH_2)_{10}COO)_4]$ (solid)

- Figure 4.59** FTIR spectrum of **Complex 6**
- Figure 4.60** UV-vis spectrum of **Complex 6** (solid)
- Figure 4.61** UV-vis spectrum of **Complex 6** (solution)
- Figure 4.62** Thermogram of **Complex 6**
- Figure 4.63** Optical structure of **Complex 6** on heating at (a) 107°C; and (b) 152°C.
- Figure 4.64** Optical structure of **Complex 6** on cooling at (a) 161°C; and (b) 78°C
- Figure 4.65** DSC curve of **Complex 6**
- Figure 4.66** Cyclic Voltammogram of **Complex 6**
- Figure 4.67** FTIR spectrum of $[\text{Cu}_2(\text{CH}_3(\text{CH}_2)_8\text{COO})_4]$
- Figure 4.68** UV-visible spectrum of $[\text{Cu}_2(\text{CH}_3(\text{CH}_2)_8\text{COO})_4]$ (solid)
- Figure 4.69** An ORTEP presentation of $[\text{Cu}_2(\text{CH}_3(\text{CH}_2)_8\text{COO})_4]$
- Figure 4.70** The packing pattern of the peacock-blue crystal of $[\text{Cu}_2(\text{CH}_3(\text{CH}_2)_8\text{COO})_4]$
- Figure 4.71** FTIR of peacock-blue crystals
- Figure 4.72** FTIR spectrum of **Complex 7**
- Figure 4.73** UV-vis spectrum of **Complex 7** (solid)
- Figure 4.74** UV-vis spectrum of **Complex 7** (solution)
- Figure 4.75** Thermogram of **Complex 7**
- Figure 4.76** Optical structure of **Complex 6** on heating at (a) 110°C; and (b) 145°C
- Figure 4.77** Optical structure of **Complex 7** on cooling at: (a) 70°C; (b) 51°C and (c) 31°C
- Figure 4.78** DSC curve of **Complex 7**
- Figure 4.79** Plot of melting temperature vs. n for $[\text{Cu}_2(p\text{HOC}_6\text{H}_4\text{COO})(\text{CH}_3(\text{CH}_2)_n\text{COO})_3]$
- Figure 4.80** Cyclic voltammogram of **Complex 7**
- Figure 4.81** FTIR spectrum of $[\text{Cu}_2(\text{CH}_3(\text{CH}_2)_6\text{COO})_4]$

- Figure 4.82** UV-visible spectrum of $[\text{Cu}_2(\text{CH}_3(\text{CH}_2)_6\text{COO})_4]$ in solution
- Figure 4.83** An ORTEP presentation of the peacock-blue crystal of $[\text{Cu}_2(\text{CH}_3(\text{CH}_2)_6\text{COO})_4]$
- Figure 4.84** The crystal packing of the peacock-blue crystal of $\text{Cu}_2(\text{CH}_3(\text{CH}_2)_6\text{COO})_4]$
- Figure 4.85** FTIR spectrum of single crystal of $[\text{Cu}_2(\text{CH}_3(\text{CH}_2)_6\text{COO})_4]$
- Figure 4.86** FTIR spectrum of **Complex 8**
- Figure 4.87** UV-vis spectrum of **Complex 8** (solution)
- Figure 4.88** Thermogram of **Complex 8**
- Figure 4.89** OPM of **Complex 8** on heating: (a) 27°C; (b) 50°C and (c) 101°C
- Figure 4.90** OPM of **Complex 8** on cooling: (a) 148°C; (b) 145°C; (c) 80°C; and (d) 32°C
- Figure 4.91** DSC curve of **Complex 8**
- Figure 4.92** Cyclic voltammogram of **Complex 8**
- Figure 4.93** FTIR spectrum of **Complex 9**
- Figure 4.94** UV-vis spectrum of **Complex 9** (solution)
- Figure 4.95** Proposed structure of **Complex 9**; solvated $\text{C}_2\text{H}_5\text{OH}$ molecules are not shown
- Figure 4.96** Thermogram of **Complex 9**
- Figure 4.97** Optical structure of **Complex 9** on heating at (a) 30°C; (b) 162°C; and (c) 175°C
- Figure 4.98** Optical structures of **Complex 9** on cooling at (a) 105°C; and (b) 96°C; (c) 25°C and (d) 24°C
- Figure 4.99** DSC curve of **Complex 9**
- Figure 4.100** FTIR spectrum of **Complex 10**
- Figure 4.101** UV-vis spectrum of **Complex 10**

- Figure 4.102** Proposed structure of **Complex 10** (K^+ is not shown)
- Figure 4.103** Thermogram of **Complex 10** (K^+ is not shown)
- Figure 4.104** DSC curve of **Complex 10**
- Figure 4.105** FTIR spectrum of **Complex 11** ($n = 8$)
- Figure 4.106** UV-vis spectrum of **Complex 11** ($n = 8$)
- Figure 4.107** Thermogram of **Complex 11** ($n = 8$)
- Figure 4.108** DSC curve of **Complex 11** ($n = 8$)
- Figure 4.109** FTIR spectrum of **Complex 12**
- Figure 4.110** UV-vis spectrum of **Complex 12**
- Figure 4.111** Thermogram of **Complex 12**
- Figure 4.112** DSC curve of **Complex 12**
- Figure 4.113** Chromatogram of dark brown liquid (**A1**)
- Figure 4.114** Chromatogram of yellow liquid (**A2**)
- Figure 4.115** Mass spectrum of molecules at R_t (min): (a) 4.1, (b) 4.4, (c) 5.8, (d) 6.7, and (e) 6.9
- Figure 4.116** Structural formulas of the molecules eluted at R_t (a) 4.1 min and (b) 6.7 min
- Figure 4.117** Structural formulas of the molecules eluted at R_t (a) 4.4 min, and (b) 6.9 min
- Figure 4.118** Chromatogram of dark brown liquid (**B1**)
- Figure 4.119** Chromatogram of yellow liquid (**B2**)
- Figure 4.120** Mass spectrum of the product at $R_t = 9.1$ min