

REFERENCES

- Affan, M. A., Shamsuddin, M. B., Yusof, M. S. M. and Yamin, B. M. (2004). Dichloromethyl[1-(2-pyridyl)ethanone benzoylhydrazonato]tin(IV). *Acta Cryst.* **E60**, m126-m128
- Akhmed, N. A. and Aleksandrov, G. G. (1970). Theoretical and experimental determination of the crystal structure of tetraphenyltin. *J. of Structural Chem.* **11(5)**, 824-827
- Ainscough, E. W., Brodie, A. M., Ranford, J. D. and Waters, J. M. (1995). Hexafluorosilicate coordination to the antitumour copper(II) salicylaldehyde benzoylhydrazone (H₂L) system: single-crystal X-ray structure of [$\{\text{Cu}(\text{HL})\text{H}_2\text{O}\}_2\text{SiF}_6$] $\cdot 2\text{H}_2\text{O}$. *Inorg. Chim. Acta* **236**, 83-88
- Ainscough, E. W., Brodie, A. M., Dobbs, A. J., Ranford, J. D. and Waters, J. M. (1998). Antitumour copper(II) salicylaldehyde benzoylhydrazone (H₂sb) complexes: physicochemical properties and the single-crystal X-ray structures of [$\{\text{Cu}(\text{H}_2\text{sb})(\text{CCl}_3\text{CO}_2)_2\}_2$] and [$\{\text{Cu}(\text{Hsb})(\text{ClO}_4)(\text{C}_2\text{H}_5\text{OH})\}_2$] and the related salicylaldehyde acetylhydrazone (H₂sa) complex, [$\text{Cu}(\text{Hsa})\text{Cl}(\text{H}_2\text{O})$] $\cdot \text{H}_2\text{O}$. *Inorg. Chim. Acta* **276**, 27-38
- Ainscough, E. W., Brodie, A. M., Denny, W. A., Finlay, G. J., Gothe, S. A. and Ranford, J. D. (1999). Cytotoxicity of salicylaldehyde benzoylhydrazone analogs and their transition metal complexes: quantitative structure-activity relationships. *J. Inorg. Biochem.* **77**, 125-133
- Alcock, N. W. and Timms, R. E. (1968). Organo-tin acetates. Part I. The crystal and molecular structure of tribenzyltin acetate. *J. Chem. Soc. A* **1968**, 1873-1876
- Alcock, N. W. and Timms, R. E. (1968). Organo-tin acetates. Part II. The crystal and molecular structure of tricyclohexyltin acetate. *J. Chem. Soc. A* **1968**, 1876-1878
- Alhadi, A. A., Ali, H. M., Puvaneswary, S., Robinson, W. T. and Ng, S. W. (2008). 2-Hydroxy-5-nitrobenzaldehyde thiosemicarbazone. *Acta Cryst.* **E64**, o1606
- Ali, H. M., Zuraini, K., Wan Jeffrey, B., Rizal, M. R. and Ng, S. W. (2008). 1-(2-Hydroxy-5-methylphenyl)ethanone [(1*H*-indol-3-yl)acetyl]hydrazone. *Acta Cryst.* **E64**, o912
- Asgebom, G., Sreedhara, A. and Rao, C. P. (1995). Oxovanadium(V) Schiff base complexes of trishydroxymethylaminomethane with salicylaldehyde and its derivatives: synthesis, characterization and redox reactivity. *Polyhedron* **14(13-14)**, 1873-1879
- Asgedom, G., Sreedhara, A., Kivikoski, J., Valkonen, J., Kolehmainen, E. and Rao, C. P. (1996). Alkoxo bound monooxo- and dioxovanadium(V) complexes: synthesis, characterization, X-ray crystal structures, and solution reactivity studies. *Inorg. Chem.* **35(19)**, 5674-5683
- Barbour, L. J. (2001). X-Seed - A Software Tool for Supramolecular Crystallography *J. Supramol. Chem.* **1(4-6)**, 189-191

Bella, S. D., Fragala, I., Ledoux, I., Diaz-Garcia, M. A. and Marks, T. J. (1997). Synthesis, characterization, optical spectroscopic, electronic structure, and a second-order nonlinear optical (NLO) properties of a novel class of donor-acceptor *bis*(salicylaldiminato)nickel(II) Schiff base NLO chromophores. *J. Am. Chem. Soc.* **119**, 9550-9557

Bellamy, L. J. (1958). The infrared spectra of complex molecules. New York: John Wiley & Sons.

Bertleff, W., Roeper, M. and Sava, X. (2007). "Carbonylation" Ullmann's Encyclopedia of Industrial Chemistry, Wiley-VCH

Bubb, W. A., Berthon, H. A. and Kuchel, P. W. (1995). Tris buffer reactivity with low-molecular-weight aldehydes: NMR characterization of the reactions of glyceraldehyde-3-phosphate. *Bioorg. Chem.* **23**, 119-130

Calligaris, M., Randaccio, L., Barbieri, R. and Pellerito, L. (1974). Crystal structure of the molecular adduct of dimethyltin(IV) chloride with *N,N'*-ethylenebis(salicylideneiminato)nickel(II). *J. Organomet. Chem.* **76**, C56-C58

Cannadine, J. C., Corden, J. P., Errington, W., Moore, P. and Wallbridge, M. G. H. (1996). Two Schiff base ligands derived from 1,2-diaminocyclohexane. *Acta Cryst.* **C52**, m1014-1017

Chan, S. C., Koh, L. L., Leung, P. H., Ranford, J. D. and Sim, K. Y. (1995). Copper(II) complexes of the antitumour-related ligand salicylaldehyde acetylhydrazone (H₂L) and the single-crystal X-ray structures of [$\{\text{Cu}(\text{HL})\text{H}_2\text{O}\}_2\] \cdot 2(\text{NO}_3)$ and [$\{\text{Cu}(\text{HL})(\text{pyridine})(\text{NO}_3)\}_2$]. *Inorg. Chim. Acta* **236**, 139-145

Chumakov, Y. M., Antosyak, B. Y., Mazus, M. D., Tsapkov, V. I. and Samus, M. N. (2003). Crystal structures of *N*-(salicylidene)*tris*(hydroxymethyl)methylamine and *N*-(5-chlorosalicylidene)*tris*(hydroxymethyl)methylamine. *Kristallografiya almeza* **45**, 1025-1030

Chumakov, Y. M., Tsapkov, V. I., Starikova, Z. A., Vorontsov, I. I., Korylukov, A. A., Antosyak, B. Y. and Perrin, M. (2003). The structure and antimycotic activity of condensation products of some aminoalcohols with salicylaldehyde and its derivatives. *J. Mol. Struct.* **647**, 269-274

Chumakov, Y. M., Tsapkov, V. I., Simonov, Y. A., Antosyak, B. Y., Bocelli, G, Perrin, M., Starikova, Z. A., Samus, N., M. and Gulea, A. P. (2005). Coordination compounds of copper(II) nitrate with products of aminotris(hydroxymethyl)methane condensation with salicyl- and 5-nitrosalicylaldehydes: synthesis and crystal structures. *Russian J. Coord. Chem.* **31(8)**, 621-629

Collins, M., Ewing, D., Mackenzie, G., Sinn, E., Sandbhor, U., Padhye, S. and Padhye, S. (2000). Metal complexes as anticancer agents: 2. Synthesis, spectroscopy, magnetism, electrochemistry, X-ray crystal structure and antimelanomal activity of the copper (II) complex of 5-amino-1-tolylimidazole-4-carboxylate in B16F10 mouse melanoma cells. *Inorg. Chem. Comm.* **3(9)**, 453-457

Crowe, A. J. and P. J. Smith. (1982). The synthesis and tin-119m mössbauer spectra of some diorganotin dihalide and dipseudohalide complexes with nitrogen- and oxygen-donor ligands. *J. Organometal. Chem.* **224**, 223-235

Crowe, A. J., Smith, P. J., Cardin, C. J., Parge, H. E. and Smith, F. E. (1984). Possible pre-dissociation of diorganotin dihalide complexes: relationship between antitumour activity and structure. *Cancer Lett.* **24**, 45-48

Cunningham, D. and Little, M. (1977). The structures of organotin(IV) trichloride adducts with *N,N'*-ethylenebis(salicylideneiminato)-nickel(II) and -copper(II) by Mössbauer spectroscopy. *J. Organomet. Chem.* **142(3)**, C58-C60

de Sousa, G. F., Hernández-Ortega, S., Jancik, V. and Valdés-Martínez, J. (2006). *trans*-(2-Acetylpyridine- κ N2-furylhydrazonato- κ^2 N¹,O)dichlorophenyltin(IV) dichloromethane solvate. *Acta Cryst.* **E62**, m819-m821

Dey, D. K., Saha, M. K., Das, M. K., Bhartiya, N. B., Bansal, R. K., Rosair, G. and Mitra, S. (1999). Synthesis and characterization of diorganotin(IV) complexes of tetradentate Schiff bases: crystal structure of *n*-Bu₂Sn(Vanophen). *Polyhedron* **18**, 2687-2696

Dey, M., Rao, C. P., Saarenketo, P. K. and Rissanen, K. (2002). Synthesis, structural diversity, inter-conversion and reactivity of Cu(II) complexes of hydroxy-rich molecules. *Inorg. Chem. Comm.* **5**, 380-383

Doidge-Harrison, S. M. V., Nowell, I. W., Cox, P. J., Howie, R. A., Taylor, O. J. and Wardell, J. L. (1991). C-Stannylated carbohydrate derivatives: IV. Structure and reactivity of methyl 4,6-*O*-benzylidene-3-deoxy-3-triphenylstannyl- α -*D*-altryopyranoside. *J. Organomet. Chem.* **401**, 273-282

Dogan, H. N., Duran, A., Rollas, S., Şener, G., Armutak, Y. and Uysal, M. K. (1998). Synthesis and structure elucidation of some new hydrazones and oxalidones: anticonvulsant activities of 2-(3-acetyloxy-2-naphthyl)-4-acetyl-5-substituted-1,3,4-oxadiazolines. *Med. Sc. Res.* **26**, 755-758

Dogan, H. N., Rollas, S. and Erdeniz, H. (1998). Synthesis, structure elucidation and antimicrobial activity of some 3-hydroxy-2-naphthoic acid hydrazide derivatives. *II Farmaco* **53**, 462-467

Dubey, S. K. and Roy, U. (2002). Review: Biodegradation of tributyltins (organotins) by marine bacteria. *Appl. Org. Chem.* **17(1)**, 3-8

Dutta, S., Bendre, R., Padhye, S., Ahmed, F., Sarkar, F. (2005). Synthesis, antioxidant properties and antiproliferative activities of tetrameric copper and copper-zinc metal complexes of catecholamine Schiff base ligand. *Synth. React. Inorg. Met-Org. Chem.* **35(1)**, 3-10

Faniran, J. A., Patel, K. S., Bailar, J. C. Jr (1974). Infrared spectra of *N,N'*-bis(salicylidene)-1,1-(dimethyl)ethylenediamine and its metal complexes. *J. Inorg. Nucl. Chem.* **36**, 1547-1551

Fun, H. K., Patil, P. S., Jebas, S. R., Sujith, K. V. and Kalluraya, B. (2008). 4-Chloro-*N'*-[(*Z*)-4-(dimethylamino)benzylidene]benzohydrazide monohydrate. *Acta Cryst.* **E64**, o1594-o1595.

Fun, H. K., Quah, C. K., Sujith, S. K. and Kalluraya, B. (2009). *N'*-[(*E*)-4-bromobenzylidene]-2-(4-isobutylphenyl)propanohydrazide. *Acta Cryst.* **E65**, o1184-o1185

Fun, H. K., Yeap, C. S., Sujith, K. V. and Kalluraya, B. (2009). *N'*-[(*E*)-4-chlorobenzylidene]-2-(4-isobutylphenyl)propanohydrazide. *Acta Cryst.* **E65**, o1196-o1197

Gielen, M. (1994). Tin-based antitumour drugs. *Metal-based Drugs* **1**, 213-219

Gielen, M. (1996). Tin-based antitumour drugs. *Coord. Chem. Rev.* **151**, 41-51

Gökçe, M., Utku, S., Gür, S., Özkul, A. and Gümüş, F. (2005). Synthesis, in vitro cytotoxic and antiviral activity of *cis*-[Pt(R(-) and S(+)-2- α -hydroxybenzylbenzimidazole)₂Cl₂] complexes. *Eur. J. Med. Chem.* **40(2)**, 135-141

Gravert, D. J. and Griffin, J. H. (1993). Specific DNA cleavage mediated by manganese complex [SalenMn(III)]⁺. *J. Org. Chem.* **58**, 820-822

Han, Y. Y., Li, Y. H. and Zhao, Q. R. (2010). *N'*-(2,5-Dihydroxybenzylidene)-2-hydroxy-3-methylbenzohydrazide. *Acta Cryst.* **E66**, o1091

Harrison, P. G and Mangia, A. (1976). Structural studies in main group chemistry : XVII. The crystal and molecular structure of phenylchlorobis(diethyldithiocarbamato)-tin(IV). *J. Organomet. Chem.* **120**, 211-216

Holeček, J., Nádvorník, M., Handlíř, K. and Lyčka, A. (1983). ¹³C and ¹¹⁹Sn NMR study of some four- and five-coordinate triphenyltin(IV) compounds. *J. Organomet. Chem.* **241(2)**, 177-184

Holeček, J., Handlíř, K., Nádvorník M. and Lyčka, A. (1983). ¹³C and ¹¹⁹Sn NMR study of some triphenyltin(IV) carboxylates. *J. Organomet. Chem.* **258(2)**, 147-153

Holeček, N., M., J., Handlíř, K. and Lyčka, A. (1986). ¹³C and ¹¹⁹Sn spectra of Di-*n*-butyltin(IV) compounds. *J. Organomet. Chem.* **315**, 299-308

Holeček, J., Handlíř, K., Černý, V., Nádvorník, M. and Lyčka, A. (1987). ¹¹⁹Sn NMR study of the equilibrium of formation of triorganotin(IV) chloride complexes in coordinating solvents. *Polyhedron* **6(5)**, 1037-1039

Holeček, J., Handlíř, K., Nádvorník, M., Teleb, S. M. and Lyčka, A. (1988). ¹³C and ¹¹⁹Sn NMR spectra of divinyltin(IV) compounds. *J. Organomet. Chem.* **339**, 61-68.

Hong, M., Yin, H., Chen, S., Wang, D. (2010). Synthesis and structural characterization of organotin(IV) compounds derived from the self-assembly of hydrazone Schiff base series and various alkyltin salts. *J. Organomet. Chem.* **695**, 653-662

- Kallet, R. H., Jasmer, R. M., Luce, J. M., Lin, L. H. and Marks, J. D. (2000). The treatment of acidosis in acute lung injury with tris-hydroxymethyl aminomethane (THAM). *Am. J. Respir. Crit. Care Med.* **161**, 1149-1153
- Karipides, A. and Wolfe, K. (1975). The crystal structure of tetra(4-methylphenyl)tin. *Acta Cryst.* **B31**, 605-608
- Karipides, A. and Oertel, M. (1977). The crystal structure of tetra(3-methylphenyl)tin. *Acta Cryst.* **B33**, 683-687
- Karipides, A., Reed, A. T., Haller, D. A. and Hayes, F. (1977). The crystal structures of the tetra(2-thienyl) derivatives of germanium, tin and lead. *Acta Cryst.* **B33**, 950-951
- Khoo, L. E. and Smith, F. E. (1982). Tin-oxygen coordination in dimethyltin dichloride *N*-arylsalicylideneimine complexes. *Polyhedron* **1(2)**, 213-214
- Khoo, L. E., Charland, J. P., Gabe, E. J. and Smith, F. E. (1987). Synthesis and structural studies for a series of aromatic Schiff base complexes with triphenyltin chloride and triphenyltin isothiocyanate. *Inorg. Chim. Acta* **128**, 139-145
- Koh, L. L., Kon, O. L., Loh, K. W., Long, Y. C., Ranford, J. D., Tan, A. L.C. and Tjan, Y. Y. (1998). Complexes of salicylaldehyde acylhydrazones: cytotoxicity, QSAR and crystal structure of the sterically hindered *t*-butyl dimer. *J. Inorg. Biochem.* **72**, 155-162
- Koon, Y. C., Lo, K. M. and Ng, S. W. (2009). *Bis*[4-(dimethylamino)pyridinium] tetrabromidobis(3,4-dichlorophenyl)stannate(IV)-1-bromo-3,4-dichlorobenzene(1/1). *Acta Cryst.* **E65**, m663
- Kumar. Das, V. G., Lo, K. M., Chen, W., Blunden, W. and Mak, T. C. W. (1987). Mössbauer and NMR studies of intramolecular coordination in thienyltetraorganotin(IV) compounds, and the x-ray crystal structure of {*C,N*-[3-(2-pyridyl)-2-thienyl]} tri(*p*-tolyl)tin(IV). *J. Organomet. Chem.* **322**, 163-175
- Kumar. Das, V. G., Lo, K. M., Chen, W. and Mak, T. C. W. (1987). Synthesis, spectroscopic study, and x-ray crystal structure of *bis*[3-(2-pyridyl)-2-thienyl-*C,N*]-diphenyltin(IV): the first example of a six-coordinate tetraorganotin compound. *Organometallics* **6**, 10-14
- Liu, F., Wang, J., Wang, R., Wang, H. and, Yao, X. (1989). Syntheses and molecular structures of complexes formed in the reaction of dichlorobis[π -cyclopentadienyl]-dicarbonyliron]tin(IV) with polydentate *o*-(nuclear substituted α -hydroxybenzylidenehydrazonomethyl)phenols. *J. Organomet. Chem.* **371(1)**, 35-41
- Lo, K. M., Ng, S. W. and Kumar Das, V. G. (1993). *Bis*[3-(2-pyridyl)-2-thienyl]di-*p*-tolylstannane, an example of a six-coordinate tetraorganotin compound. *J. Cryst. Spec. Res.* **23(12)**, 933-936
- Lo, K. M. and Ng, S. W. (2008). *Bis*[4-(dimethylamino)pyridinium] tribromidochloridodimethylstannate(IV). *Acta Cryst.* **E64**, m834

- Lyčka, A., Holeček, J., Nádvořník M. and Handlíř, K. (1985). ^{13}C , ^{15}N and ^{119}Sn NMR spectral evidence for tin five-coordination in triorganotin(IV) oxinates. *J. Organomet. Chem.* **280**, 323-329
- Lyčka, A., Jirman, J. and Koloničný, A. (1987). ^{13}C and ^{119}Sn NMR spectra of some tribenzyltin(IV) compounds. *J. Organomet. Chem.* **333**, 305-315
- Mikhaylova, Y., Adam, G., Häussler, L., Eichhorn, K. J. and Voit, B. (2006). Temperature-dependent FTIR spectroscopic and thermoanalytic studies of hydrogen bonding of hydroxyl (phenolic group) terminated hyperbranched aromatic polyesters. *J. Mol. Struct.* **788**, 80-88
- Mohd Ali, H., M. Zain, S., Basirun, W. J., Rahuma, S. M., Sharifah Rohaiza, S. O., Abdullah, N. and Teoh, H. L. (2004). Synthesis, structural characterization, electrochemical and anti-microbial studies of complexes from the NO ligating Schiff base complexes of furoic acid hydrazide. *Malaysian J. of Sc.* **23(1)**, 119-127
- Mohd Ali, H., M. Zain, S., Basirun, W. J., Rahuma, S. M., Sharifah Rohaiza, S. O., Abdullah, N. and Teoh, H. L. (2004). Spectral, electrochemical and anti-microbial studies of nicotinic acid hydrazide Schiff base complexes of cobalt (II) and manganese (II). *Malaysian J. of Sc.* **23(1)**, 129-136
- Molloy, K. C., Purcell, T. G. and Quill, K. (1984). Organotin biocides : I. The structure of triphenyltin acetate. *J. Organomet. Chem.* **267(3)**, 237-247
- Nádvořník M., Holeček, J., Handlíř, K., and Lyčka, A. (1984). ^{13}C and ^{119}Sn NMR spectra of some four- and five-coordinate tri-*n*-butyltin(IV) compounds. *J. Organomet. Chem.* **275**, 43-51
- Narang, K. K., Rao, T. R., Shrestha and S., Shrestha, S. (2000). Synthesis, characterization, thermal and electrical properties of yttrium(III) and lanthanide(III) complexes of salicylaldehyde benzoyl hydrazone. *Synth. React. Inorg. Met-Org. Chem.* **30(5)**, 931 – 954
- Nelson, D. L. and Cox, M. M. L. (2000). Principles of Biochemistry, 3rd Ed. Worth Publishing: New York
- Ng, S. W. (1997). Tetra(*p*-chlorophenyl)tin. *Acta Cryst.* **C53**, 273-274
- Ng, S. W. (2008). Low-temperature redetermination of 4-chloro-2-[tris(hydroxymethyl)methyliminomethyl]phenol as zwitterionic 4-chloro-2-[tris(hydroxymethyl)methyliminomethyl]phenolate. *Acta Cryst.* **C64**, o2455
- Odabaşoğlu, M., Albayrak, Ç., Büyükgüngör, O. and Lönnecke. P. (2003). 2-[[Tris(hydroxymethyl)methyl]aminomethylene}cyclohexa-3,5-dien-1(2*H*)-one and its 6-hydroxy and 6-methoxy derivatives. *Acta Cryst.* **C59**, o616-o619
- Odabaşoğlu, M., Albayrak, C., Özkanca, R., Aykan, F. Z. and Lonecke. P. (2003). Some polyhydroxy azo-azomethine derivatives of salicylaldehyde: synthesis, characterization, spectroscopic, molecular structure and antimicrobial activity studies. *J. Mol. Struct.* **840**, 71-89

- Ogawa, K. and Harada, J. (2003). Aggregation controlled proton tautomerization in salicylideneanilines. *J. Mol. Struct.* **647**, 211-216
- Pan, H, Willem, R., Meunier-Piret, J. and Gielen, M. (1990). Intramolecular O.fwdarw. Sn coordination in (Z)-17-[2-(triphenylstannyl)vinyl]-4-estren-17-ol: evidence by x-ray diffraction analysis and iodo demetalation. *Organometallics* **9**, 2199–2201
- Panneerselvam, P., Nair, R. R., Vijayalakshmi, G., Subramanian, E. H. and Sridhar, S. K. (2005). Synthesis of Schiff bases of 4-(4-aminophenyl)-morpholine as potential antimicrobial agents. *Eur. J. Med. Chem.* **40(2)**, 225-229
- Pavia, D. L., Lampman, G. M. and Kriz, G. S. (2001). Introduction to spectroscopy, 3rd Edition. Thomson learning, Inc. USA.
- Pawanoji, A. A. and Mehta, B. H. (2009). Synthesis, characterization and x-ray diffraction studies of Co(II), Ni(II) and Pd(II) complexes of Schiff base derived from 3-hydroxy benzaldehyde and 3-hydroxy-2-naphthoic hydrazide. *Asian J. Chem.* **21(9)**, 6869-6876
- Pellerito, L., Cefalú, R., Silvestri, A., Di Blanca, F., Barbieri, R., Haupt, H. J., Preut, H. and Huber, F. (1974). Complexes of organometallic compounds : XXXIX. synthesis, infrared and Mössbauer studies on monoorganotin(IV) complexes with tridentate ligands. *J. Organomet. Chem.* **78(1)**, 101-106
- Pettinari, C., Marchetti, F., Pettinari, R., Martini, D., Drozodov, A. and Troyanov, S. (2001). Synthesis and characterisation of tin(IV) and organotin(IV) derivatives 2-[[2-(2-hydroxyphenyl)imino]methyl]phenol. *Inorg. Chim. Acta* **325**, 103-114
- Poller, R. C. (1970). The chemistry of organotin compounds. New York: Academic Press
- Pfeiffer, P., Buchholz, E., Bauer, O. (1931). Innere komplexsalze von oxyaldiminen und Oxyketiminen. *J. Prakt. Chem.* **129**, 163-177
- Pfeiffer, P. and Pfitzner, H. (1936). Hochatomare nebenvalenzringe mit meta- und para-Kondensatio. *J. Prakt. Chem.* **145**, 243-256
- Pfeiffer, P., Hesse, Th., Pfitzner, H., Scholl, W. and Thielert, H. (1937). Innere Komplexsalze der Aldimin- und Azoreihe. *J. Prakt. Chem.* **149**, 217-296
- Qiu, X. Y., Fang, X. N., Liu, W. S. and Zhu, H. L. (2006). (E)-N'-(3-Chlorobenzylidene)isonicotinohydrazide. *Acta Cryst.* **E62**, o2685-o2686
- Qiu, X. Y., Luo, Q. Y., Yang, S. L. and Liu, W. S. (2006). (E)-2-Hydroxy-N'-[1-(4-methoxyphenyl)-ethylidene]benzohydrazide. *Acta Cryst.* **E62**, o4291-o4292.
- Qiu, X. Y., Luo, Z. G., Yang, S. L. and Liu, W. S. (2006). (E)-2-Hydroxy-N'-(2-naphthylmethylene)benzohydrazide. *Acta Cryst.* **E62**, o3531-o3532.

- Raparti, V., Chitre, T., Bothara, K., Kumar, V., Dangre, S., Khachane, C. Gore, S. and Deshmane, B. (2009). Novel 4-(morpholin-4-yl)-N'-(arylidene)benzohydrazides: synthesis, antimycobacterial activity and QSAR investigations. *Eur. J. Med. Chem.* **44**, 3954-3960
- Routier, S., Bernier, J., Waring, M. J., Colson, P., Houssier, C. and Bailly, C. (1996). synthesis of a functionalized Salen-copper complex and its interaction with DNA. *J. Org. Chem.* **61**, 2326–2331
- Sandbhor, U., Padhye, S. and Sinn, E. (2002). *cis* di-oxomolybdenum(VI) complexes with a tridentate *ONO* donor ligand; synthesis, X-ray crystal structure, spectroscopic properties and oxotransfer chemistry. *Trans. Met. Chem.* **27**, 681-685
- Saxena, A. K. (1987). Organotin compounds: toxicology and biomedical applications. *Appl. Organomet. Chem.*, 1(1), 39-56
- Saxena, A. K. and F. Huber, F. (1989). Organotin compounds and cancer chemotherapy. *Coord. Chem. Rev.* **95**, 109-123
- Sengupta, A., Chattopadhyay, T. K. and Majee, B. (1982). Organotin(benzeneazo)phenoxy acetates-examples of seven-coordinated triorganotin chelates. *Indian J. Chem.* **21A**, 1090-1095
- Sheldrick, G. M. (1993). SHELXL-93, Program for structure refinement, University of Göttingen, Germany
- Sheldrick, G. M. (1997). SHELXL-97, Crystal structure analysis program, University of Göttingen, Germany
- Shujha, S., Shah, A., Rehman, Z., Muhammad, N., Ali, S., Qureshi, R., Khalid, N. And Meetsma, A. (2010). Diorganotin(IV) derivatives of *ONO* tridentate Schiff base: Synthesis, crystal structure, *in vitro* antimicrobial, anti-leishmanial and DNA binding studies. *Eur. J. Med. Chem.* **45**, 2902-2911
- Signorini, O., Dockal, E. R., Castellano, G. and Oliva, G. (1996). Synthesis and characterization of aqua[*N,N'*-ethylenebis(3-ethoxysalicylideneaminato)]dioxouranium (VI). *Polyhedron* **15(2)**, 245-255
- Singh, H. L., Varshney, S. and Varshney, A. K. (1999). Organotin(IV) complexes of biologically active Schiff bases derived from heterocyclic ketones and sulphur drugs. *Appl. Organomet. Chem.* **13(9)**, 637–641
- Sisido, K., Takeda, Y. and Kinugawa, Z. (1961). Direct synthesis of organotin compounds. I. Di- and tribenzyltin chlorides. *J. Am. Chem. Soc.* **83**, 538-541
- Stern, A. and Becker, E. I. (1964). Synthesis of Triaryltin Hydrides. *J. Org. Chem.* **29**, 3221-3225

- Sui, Y., Zeng, X., Fang, X., Fu, X., Xiao, Y., Chen, L., Li, M. and Cheng, S. (2007). Syntheses, structure, redox and catalytic epoxidation properties of dioxomolybdenum(VI) complexes with Schiff base ligands derived from tris(hydroxymethyl)aminomethane. *J. of Molecular Catalysis A: Chemical* **270**, 61-67
- Tai, X. S., He, Y., Feng, Y. M. (2007). Crystal structure of (*E*)-*N'*-(4-hydroxybenzylidene)benzohydrazide monohydrate, C₁₄H₁₂N₂O₂·H₂O. *Z. Kristallogr.* **222**, 343-344
- Tatar, L., Nazır, H., Gümüşer, M., Kale, C. and Atakol, O. (2005). Synthesis, crystal structure and electrochemical behaviour of water soluble Schiff bases. *Z. Kristallogr.* **220**, 639-642
- Tiekink, E. R. T. (1991). Structural chemistry of organotin carboxylates: a review of the crystallographic literature. *Appl. Organomet. Chem.* **5**, 1-23
- Tzschach, A. and Jurkschart, K. (1986). Pentacoordinated tetraorganotin and silicon compounds-synthesis and structures. *Pure Appl. Chem.* **58(4)**, 639-646
- Van der Kerk, G. J. M. and Luitjen, J. G. A. (1954). The biocidal properties of organotin compounds. *Appl. Chem.* **4**, 314-319
- Wang, J., Liu, F., Zhang, Y., Wang, R., Wang, H. and Yao, X. (1989). Synthesis and structure of pentacoordinate tin(IV) complexes. *J. Organomet. Chem.* **375(2)**, 173-181
- Wu, W. S., Chen, J., Dai, J. C., Lin, J. M., Lan, X. R. and Huang, T. T. (2005). Crystal structure and fluorescence of *N*-picoloylhydrazide and its binuclear palladium complex. *Acta Chim. Sinica.* **63**, 1967-1973
- Yap, Q. L., Lo, K. M. and Ng, S. W. (2008). Bis[4-(dimethylamino)pyridinium] tetrabromidodiphenylstannate(IV). *Acta Cryst.* **E64**, m696
- Yap, Q. L., Lo, K. M. and Ng, S. W. (2008). Tetrakis(4-methyl-2-thienyl)tin(IV). *Acta Cryst.* **E64**, m1410
- Yaul, S. R., Yaul, A. R., Pethe, G. B. and Aswar, A. S. (2009). Synthesis and characterization of transition metal complexes with *N,O*-chelating hydrazone Schiff base ligand. *American-Eurasian J. Sci. Res.*, **4(4)**, 229-234
- Yin, H., Hong, M. and Wang, Q. B. (2004). Synthesis and characterization of organotin(IV) compounds with Schiff base ligand pyruvic acid isonicotinyl hydrazone and crystal structure of $[n\text{-Bu}_2\text{Sn}(\text{C}_9\text{H}_8\text{N}_3\text{O}_3)(\text{H}_2\text{O})]_2$ and $\{[n\text{-Bu}_2\text{Sn}(\text{C}_9\text{H}_8\text{N}_3\text{O}_3)\text{O}]\text{Sn}(\text{Bu-}n)_3\}_2 \cdot 2\text{H}_2\text{O}$. *Indian J. Chem.* **43A**, 2301-2306
- Yin, H., Hong, M., Xu, H., Gao, Z., Li, G. and Wang, D. (2005). Self-assembly of organotin(IV) moieties with the Schiff-base ligands pyruvic acid isonicotinyl hydrazone and pyruvic acid salicylhydrazone: synthesis, characterization, and crystal structures of monomeric or polymeric complexes. *Eur. J. Inorg. Chem.* **22**, 4572-4581

- Yin, H. D., Hong, M., Wang, Q. B., Xue, S. C. and Wang, D. Q. (2005). Synthesis and structural characterization of diorganotin(IV) esters with pyruvic acid isonicotinyl hydrazone and pyruvic acid salicylhydrazone Schiff bases. *J. Organomet. Chem.* **690**, 1669-1676.
- Yin, H. D., Hong, M., Li, G. and Wang, D. Q. (2005). Synthesis, characterization and structural studies of diorganotin(IV) complexes with Schiff base ligand salicylaldehyde isonicotinylhydrazone. *J. Organomet. Chem.* **690**, 3714-3719
- Yin, H. and Chen, S. W. (2006). Synthesis and characterization of di- and tri-organotin(IV) complexes with Schiff base ligand pyruvic acid 3-hydroxy-2-naphthoyl hydrazone. *Inorg. Chim. Acta* **359**, 3330-3338
- Yin, H. D. and Chen, S. W. (2006). Synthesis and characterization of organotin(IV) compounds with Schiff base of *o*-vanilin-2-thiophenoylhydrazone. *J. Organomet. Chem.* **691**, 3103-3108
- Yin, H., Chen, S., Li, W. and Wang, D. (2007). Synthesis, characterization and crystal structures of the organotin(IV) compounds with the Schiff base ligands of pyruvic acid thiophene-2-carboxylic hydrazone and salicylaldehyde thiophene-2-carboxylic hydrazone. *Inorg. Chim. Acta* **360**, 2215-2223
- Yin, H., Cui, J. and Qiao, Y. (2008). The first 1D tetranuclear organotin(IV) complex with *N'*-acylsalicylhydrazide: synthesis, characterization and crystal structure. *Inorg. Chem. Comm.* **11**, 684-686
- Zhang, J., Wu, L., Zhuang, L. and Wang, G. (2009). 3-Methoxybenzaldehyde thiosemicarbazone. *Acta Cryst.* **E65**, o884