

## LIST OF FIGURES

<b>Figures</b>	<b>Captions</b>	<b>Pages</b>
Figure 2.1	Ring opening of poly( $\epsilon$ -caprolactone)	19
Figure 2.2	Schematic representation of type A2 chains	20
Figure 2.3	Chemical structure of poly( $\epsilon$ -caprolactone)	21
Figure 3.1	Illustration of polymer electrolyte preparation	29
Figure 3.2	Schematic diagram of a DSC thermogram	33
Figure 3.3	DSC machine: TA Instruments DQ200	34
Figure 3.4	SEM machine: Leica's model S440	36
Figure 3.5	Schematic diagram of XRD pattern of polyethylene [Lagaron et al., 2000]	37
Figure 3.6	XRD machine: SIEMENS D5000	39
Figure 3.7	FTIR machine: Thermo Scientific Nicolet iS10	41
Figure 3.8	EIS instrument: HIOKI 3532 Z LCR Hi-tester	42
Figure 3.9	Some impedance plots and the corresponding equivalent circuits	42
Figure 4.1	DSC thermograms of PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN. Inset shows the magnified thermograms indicating $T_m$	49
Figure 4.2	Magnified DSC thermograms of PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN, showing $T_g$	50
Figure 4.3	Relative degree of crystallinity as a function of salt content for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN	53
Figure 4.4	DSC thermograms of pure EC, pure PCL and PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC	55
Figure 4.5	Magnified DSC thermograms of PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC, showing $T_g$	56
Figure 4.6	Magnified DSC thermograms of PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC, showing $T_m$	57

Figure 4.7	Relative degree of crystallinity as a function of EC content for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC	59
Figure 5.1	Schematic diagram of spherulite morphology	63
Figure 5.2	Micrographs (300x) of PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature	66
Figure 5.3	Salt concentration dependence of average diameter of spherulites from micrograph of PCL-NH <sub>4</sub> SCN films	68
Figure 5.4	Micrographs (500x) of PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature	70
Figure 5.5	XRD pattern of pure PCL membrane without ammonium salt	72
Figure 5.6	XRD patterns of (a) pure NH <sub>4</sub> SCN salt and PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature	73
Figure 5.7	The curve fitting XRD patterns of PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN	76
Figure 5.8	Relative degree of crystallinity as a function of salt concentration, calculated from XRD pattern	77
Figure 5.9	XRD patterns of (a) pure NH <sub>4</sub> SCN, (b) pure PCL and PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature	79
Figure 5.10	The curve fitting XRD patterns of PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC	82
Figure 5.11	Relative degree of crystallinity as a function of EC concentration, calculated from XRD pattern	82
Figure 6.1	Room temperature FTIR spectra and curve fitting results of poly(oxyethylene)/siloxane-Mg(CF <sub>3</sub> SO <sub>3</sub> ) <sub>2</sub> in the asymmetrical stretching of SO <sub>3</sub> region	86
Figure 6.2	Room temperature IR spectra of pure PCL film in the region between 650 and 4000 cm <sup>-1</sup> . Inset presents the chemical structure of PCL	87
Figure 6.3	Magnified IR spectra of pure PCL film in the region between 1100 and 1900 cm <sup>-1</sup>	88
Figure 6.4	Room temperature IR spectra of (a) NH <sub>4</sub> SCN and (b) THF in the region between 650 and 4000 cm <sup>-1</sup> . Insets present the respective chemical structures	90
Figure 6.5	Evolution of room temperature IR spectra of (a) NH <sub>4</sub> SCN and PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN	92

	in the region between 700 and 1500 $\text{cm}^{-1}$	
Figure 6.6	Evolution of room temperature IR spectra of (a) $\text{NH}_4\text{SCN}$ and PCL- $\text{NH}_4\text{SCN}$ films added with 0 to 32 wt.% $\text{NH}_4\text{SCN}$ displaying N-H stretching and carbonyl stretching regions	93
Figure 6.7	Schematic representation of the cation coordination in PCL- $\text{NH}_4\text{SCN}$ polymer complex. The dotted line represents the ion-dipole interaction between cation and PCL. Arrow shows the direction of the cation movement	95
Figure 6.8	The curve-fitting IR spectra of PCL- $\text{NH}_4\text{SCN}$ films added with 0 to 32 wt.% $\text{NH}_4\text{SCN}$ displaying CN stretching modes	99
Figure 6.9	Integral area percentage of free ions, contact ion pairs and ion aggregates as a function of salt concentration for PCL- $\text{NH}_4\text{SCN}$ system	100
Figure 6.10	Room temperature IR spectra of pure EC in the region between 650 and 2150 $\text{cm}^{-1}$ . Inset presents the chemical structure of EC	102
Figure 6.11	Room temperature IR spectra of (a) pure EC, (b) pure PCL and PCL-EC films added with (c) 10 wt.% and (d) 30 wt.% EC in the region between 650 and 2150 $\text{cm}^{-1}$	103
Figure 6.12	Magnified IR spectra of (a) pure PCL, PCL-EC films added with (b) 10 wt.% and (c) 30 wt.% EC in the region between 1750 and 1850 $\text{cm}^{-1}$	104
Figure 6.13	Room temperature IR spectra of (a) $\text{NH}_4\text{SCN}$ , (b) pure EC and EC- $\text{NH}_4\text{SCN}$ electrolyte added with (c) 10 wt.% and (d) 30 wt.% $\text{NH}_4\text{SCN}$	105
Figure 6.14	Magnified IR spectra of (a) $\text{NH}_4\text{SCN}$ , (b) pure EC and EC- $\text{NH}_4\text{SCN}$ electrolyte added with (c) 10 wt.% and (d) 30 wt.% $\text{NH}_4\text{SCN}$ in the region between 1600 and 1800 $\text{cm}^{-1}$	106
Figure 6.15	Magnified IR spectra of (a) $\text{NH}_4\text{SCN}$ , (b) pure EC and EC- $\text{NH}_4\text{SCN}$ electrolyte added with (c) 10 wt.% and (d) 30 wt.% $\text{NH}_4\text{SCN}$ in the region between 1200 and 1600 $\text{cm}^{-1}$	106
Figure 6.16	Magnified IR spectra of (a) $\text{NH}_4\text{SCN}$ , (b) pure EC and EC- $\text{NH}_4\text{SCN}$ electrolyte added with (c) 10 wt.% and (d) 30 wt.% $\text{NH}_4\text{SCN}$ in the region between 690 and 1190 $\text{cm}^{-1}$	107
Figure 6.17	Evolution of room temperature IR spectra of (a) $\text{NH}_4\text{SCN}$ , (b) pure EC, (c) pure PCL and PCL- $\text{NH}_4\text{SCN}$ -EC films added with 0 to 50 wt.% EC	109
Figure 6.18	Magnified IR spectra of PCL- $\text{NH}_4\text{SCN}$ -EC films added with 0 to 50 wt.% EC in the region between 650 and 1150 $\text{cm}^{-1}$	110

Figure 6.19	Magnified IR spectra of PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC in the region between 1500 and 1900 cm <sup>-1</sup>	112
Figure 6.20	Schematic representation of the cation coordination in PCL-NH <sub>4</sub> SCN-EC polymer complex. The dotted line represents the ion-dipole interaction between cation and PCL and the stripe line represent the dipole-dipole interaction between PCL and EC. Arrow shows the direction of the cation movement	114
Figure 6.21	The curve-fitting IR spectra of PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC displaying CN stretching modes	117
Figure 6.22	Integral area percentage of free ions, contact ion pairs and ion aggregates as a function of EC concentration for PCL-NH <sub>4</sub> SCN-EC system	118
Figure 7.1	Impedance plot for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature	124
Figure 7.2	Fitting of the equivalent circuit to the experimental data of PCL-NH <sub>4</sub> SCN films added with 26 wt.% NH <sub>4</sub> SCN	126
Figure 7.3	Salt concentration dependence of dc conductivity for PCL-NH <sub>4</sub> SCN films at room temperature. The line is just a guide to the eye	127
Figure 7.4	Variation of $\epsilon'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature. Inset displays the $\epsilon'$ versus $\log f$ for 0 wt.% salt.	129
Figure 7.5	Salt concentration dependence of real relative permittivity for PCL-NH <sub>4</sub> SCN films at selected frequencies	131
Figure 7.6	Conductivity as a function of real relative permittivity for PCL-NH <sub>4</sub> SCN films at selected frequencies. Salt concentrations in wt.% are labeled within the figures.	132
Figure 7.7	Variation of $\epsilon''$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature. Inset displays the magnified graph	133
Figure 7.8	Variation of $M'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature. Inset shows the magnified graph	135
Figure 7.9	Variation of $M''$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature	136
Figure 7.10	Argand plot of $M''$ versus $M'$ for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN. Inset shows the	138

	magnified graph	
Figure 7.11	Variation of loss tangent versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature.	139
Figure 7.12	Salt concentration dependence of relaxation time at room temperature. The line is just a guide to the eye	140
Figure 7.13	Variation of $\log \sigma'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN at room temperature	141
Figure 7.14	Impedance plot for PCL-NH <sub>4</sub> SCN films added with 10 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C.	143
Figure 7.15	The temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN films added with 10 wt.% NH <sub>4</sub> SCN.	144
Figure 7.16	Impedance plot for PCL-NH <sub>4</sub> SCN films added with 20 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C.	145
Figure 7.17	The temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN films added with 20 wt.% NH <sub>4</sub> SCN.	145
Figure 7.18	Impedance plot for PCL-NH <sub>4</sub> SCN films added with 26 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C.	146
Figure 7.19	The temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN films added with 26 wt.% NH <sub>4</sub> SCN.	146
Figure 7.20	Impedance plot for PCL-NH <sub>4</sub> SCN films added with 32 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C.	147
Figure 7.21	The temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN films added with 32 wt.% NH <sub>4</sub> SCN.	147
Figure 7.22	The VTF temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN films added with 0 to 32 wt.% NH <sub>4</sub> SCN.	149
Figure 7.23	Variation of $\log \epsilon'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 32 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C	151
Figure 7.24	Variation of $\epsilon''$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 32 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C. Inset displays the magnified graph	151
Figure 7.25	Variation of $M'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 32 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C	152
Figure 7.26	Variation of $M''$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 32 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C	153

Figure 7.27	Variation of loss tangent versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 32 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C.	154
Figure 7.28	Variation of $\log \sigma'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN films added with 32 wt.% NH <sub>4</sub> SCN at temperature from 25 to 55 °C	154
Figure 7.29	Impedance plot for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature.	156
Figure 7.30	Fitting of the equivalent circuit to the experimental data of PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC.	157
Figure 7.31	EC concentration dependence of dc conductivity for PCL-NH <sub>4</sub> SCN-EC films at room temperature.	158
Figure 7.32	Variation of $\log \epsilon'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature	159
Figure 7.33	EC concentration dependence of $\log \epsilon'$ for PCL-NH <sub>4</sub> SCN-EC films at selected frequencies	160
Figure 7.34	Conductivity as a function of $\epsilon'$ at selected frequencies. EC concentrations in wt.% are labeled within the figures The line is just a guide to the eye	162
Figure 7.35	Variation of $\epsilon''$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature. Inset displays the magnified graph	163
Figure 7.36	Variation of $M'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature	163
Figure 7.37	Variation of $M''$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature	164
Figure 7.38	Argand plot of $M''$ versus $M'$ for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC. Inset shows the magnified graph for 40 wt.% EC	165
Figure 7.39	Variation of loss tangent versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature	166
Figure 7.40	EC concentration dependence of relaxation time and conductivity at room temperature.	166
Figure 7.41	Variation of $\log \sigma'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC at room temperature	167
Figure 7.42	Impedance plot for PCL-NH <sub>4</sub> SCN-EC films added with 20 wt.% EC at temperature from 25 to 50 °C.	169

---

Figure 7.43	The temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN-EC films added with 20 wt.% EC.	169
Figure 7.44	Impedance plot for PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC at temperature from 25 to 45 °C.	170
Figure 7.45	The temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC.	170
Figure 7.46	Impedance plot for PCL-NH <sub>4</sub> SCN-EC films added with 40 wt.% EC at temperature from 25 to 45 °C.	171
Figure 7.47	The temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN-EC films added with 40 wt.% EC.	171
Figure 7.48	The VTF temperature dependence of conductivity for PCL-NH <sub>4</sub> SCN-EC films added with 0 to 50 wt.% EC.	172
Figure 7.49	Variation of $\log \epsilon'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC at temperature from 25 to 45 °C. Inset displays the magnified graph	174
Figure 7.50	Variation of $\epsilon''$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC at temperature from 25 to 45 °C. Inset displays the magnified graph	174
Figure 7.51	Variation of $M'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC at temperature from 25 to 45 °C	175
Figure 7.52	Variation of $M''$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC at temperature from 25 to 45 °C	175
Figure 7.53	Variation of loss tangent versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC at temperature from 25 to 45 °C.	176
Figure 7.54	Variation of $\log \sigma'$ versus $\log f$ for PCL-NH <sub>4</sub> SCN-EC films added with 30 wt.% EC at temperature from 25 to 45 °C	177