

LIST OF FIGURES

| Figure | Caption | Page |
|---------------|--|-------------|
| Figure 2.1 | Schematic diagram of amorphous and crystalline regions of a polymer | 9 |
| Figure 2.2 | Chemical structure of (a) LiTf and (b) delocalization of negative charge in triflate anion | 15 |
| Figure 2.3 | Schematic diagram showing effect of IL doping into polymer electrolyte matrix where K^+ is potassium cation and IL is ionic liquid [Taken from Singh et al. (2008)] | 25 |
| Figure 2.4 | Chemical structures of (a) BMII and (b) BMITf | 26 |
| Figure 2.5 | Temperature dependence plots of the conductivity of PAN–LiClO ₄ containing 0 (CPE–0), 1 (CPE–1) and 3 (CPE–3) wt.% SAP [Taken from Chen et al. (2011)] | 30 |
| Figure 2.6 | Cation motion in a polymer electrolyte via (a) intrachain hopping, (b) interchain hopping, (c) intrachain hopping via ion cluster and (d) intercluster hopping [Taken from Gray, 1997] | 31 |
| Figure 2.7 | Temperature dependence plots of the conductivity of PEO–LiTf comprised of various EO/Li ratios. [Taken from Karan et al. (2008)] | 32 |
| Figure 3.1 | Image of reflux set-up during preparation of polymer electrolytes | 35 |
| Figure 3.2 | Flow chart showing the preparation procedure of PEMA/PVdF–HFP–LiTf polymer electrolyte films | 36 |
| Figure 3.3 | Image of transparent PEMA/PVdF–HFP based polymer electrolytes | 36 |
| Figure 3.4 | Flow chart for the preparation procedure of PEMA/PVdF–HFP–LiTf–EC polymer electrolyte films | 38 |
| Figure 3.5 | Flow chart for the preparation procedure of PEMA/PVdF–HFP–LiTf–PC polymer electrolyte films | 40 |
| Figure 3.6 | Flow chart for the preparation procedure PEMA/PVdF–HFP–LiTf–BMII polymer electrolyte films | 41 |

| | | |
|-------------|---|----|
| Figure 3.7 | Image of semi-transparent PEMA/PVdF-HFP-LiTf-BMII based polymer electrolytes | 42 |
| Figure 3.8 | Flow chart for the preparation procedure of PEMA/PVdF-HFP-LiTf-BMITf polymer electrolyte films | 43 |
| Figure 3.9 | FTIR spectra of (a) pure PEO, (b) pure LiTf and (c) PEO:LiTf (95:5; wt.:wt.) sample [Taken from Ramesh et al., 2008] | 44 |
| Figure 3.10 | TGA curves of PMMA and its nanocomposites [Taken from Costache et al., 2006] | 48 |
| Figure 3.11 | TGA thermograms of PVDF-HFP films containing 0 and 16.3 wt.% PEG [Taken from Chung et al., 2003] | 48 |
| Figure 4.1 | FTIR spectra in the region (a) 3800–2800 cm^{-1} , (b) 1800–800 cm^{-1} and (c) 1000–800 cm^{-1} (enlarged) of i. PEMA, ii. PVdF-HFP, iii. S-0, iv. S-10, v. S-20, vi. S-30, vii. S-40 and viii. LiTf | 52 |
| Figure 4.2 | FTIR spectra in the region (a) 3200–2600 cm^{-1} and (b) 1820–720 cm^{-1} of i. PEMA, ii. PVdF-HFP and iii. PEMA/PVdF-HFP(70:30) or S-0 | 53 |
| Figure 4.3 | Deconvolution and band-fitting components in the region (a) 1440–1340 cm^{-1} , (b) 1220–1100 cm^{-1} and (c) 1080–980 cm^{-1} of i. PEMA, ii. PVdF-HFP and iii. S-0 | 54 |
| Figure 4.4 | Schematic diagram of possible interactions in (a) PEMA, (b) PVdF-HFP and (c) S-0 (where••••represents interactions within a polymer and ■■■represents intermolecular interactions between PEMA and PVdF-HFP) | 57 |
| Figure 4.5 | FTIR spectra in the region 1800–1600 cm^{-1} for (a) S-0, (b) S-10 (Inset: Enlarged deconvoluted bands of S-10), (c) S-20, (d) S-30, (e) S-40, (f) 70 wt.% PEMA-30 wt.% LiTf, (g) 70 wt.% PVdF-HFP-30 wt.% LiTf and (h) LiTf | 59 |
| Figure 4.6 | Deconvoluted FTIR spectra in the region 1220–1120 cm^{-1} for (a) S-0, (b) S-10, (c) S-20, (d) S-30 and (e) S-40 | 61 |
| Figure 4.7 | FTIR spectra in the region 1200–1100 cm^{-1} of (a) PEMA-based film, (b) PVdF-HFP based film containing i. 0 and ii. 30 wt. % of LiTf | 62 |
| Figure 4.8 | Deconvoluted FTIR spectra between 1340–1200 cm^{-1} for (a) S-0, (b) S-10, (c) S-20, (d) S-30 and (e) S-40 | 63 |

| | | |
|-------------|--|----|
| Figure 4.9 | FTIR spectra in the region (a) 1450–1350 cm^{-1} and (b) 1100–1000 cm^{-1} for i. PVdF–HFP and ii. 70 wt.% PVdF–HFP–30 wt.% LiTf | 64 |
| Figure 4.10 | Schematic diagram of possible interactions in PEMA/PVdF–HFP–LiTf system (where ■■■ represents intermolecular interactions between PEMA and PVdF–HFP and ●●● represents coordinate bonds between Li^+ ions and PEMA/PVdF–HFP) | 65 |
| Figure 4.11 | Deconvolution between 1080 and 980 cm^{-1} of (a) S–10, (b) S–20, (c) S–30 and (d) S–40 | 66 |
| Figure 4.12 | Area % of the different ionic species of the triflate anion with respect to wt. % of LiTf in PEMA/PVdF–HFP (70:30) blend | 67 |
| Figure 4.13 | FTIR spectra in the region (a) 3800–2800 cm^{-1} and (b) 1800–800 cm^{-1} of i. EC–0, ii. EC–2, iii. EC–4, iv. EC–6, v. EC–8, vi. EC–10 and vii. EC | 69 |
| Figure 4.14 | FTIR spectra in the region 1850–1600 cm^{-1} for (a) EC–0, (b) EC–2, (c) EC–4, (d) EC–6, (e) EC–8, (f) EC–10, (g) 90 wt.% PEMA–10 wt.% EC, (h) 90 wt.% PVdF–HFP–10 wt.% EC, (i) S–0–EC, (j) EC and (k) LiTf–EC | 71 |
| Figure 4.15 | Deconvoluted FTIR spectra of (a) EC–0, (b) EC–2, (c) EC–4, (d) EC–6, (e) EC–8 and (f) EC–10 in the region 1340–1200 cm^{-1} | 72 |
| Figure 4.16 | Deconvoluted FTIR spectra in the region between 1220 and 1120 cm^{-1} of (a) EC–0, (b) EC, (c) EC–2, (d) EC–4, (e) EC–6, (f) EC–8 and (g) EC–10 between 1220–1120 cm^{-1} | 74 |
| Figure 4.17 | Deconvoluted FTIR spectra of (a) 6 wt.% LiTf–94 wt.% EC, (b) 90 wt.% PEMA–10 wt.% EC, (c) 90 wt.% PVdF–HFP–10 wt.% EC (in the region 1320–1120 cm^{-1}) and (d) 90 wt.% S–0–10 wt.% EC in the region 1220–1120 cm^{-1} | 75 |
| Figure 4.18 | Schematic diagram of possible interactions in PEMA/PVdF–HFP–LiTf–EC system (where ●●● represents coordinate bonds with Li^+ ions, ●●● represents electrostatic interactions between Li^+ and EC and ■■■ represents intermolecular interactions between EC and PEMA/PVdF–HFP) | 76 |
| Figure 4.19 | IR deconvoluted bands between 1080 and 950 cm^{-1} for samples (a) EC–2, (b) EC–4, (c) EC–6, (d) EC–8 and (e) EC–10 | 77 |
| Figure 4.20 | Area % of free ions and ion pairs with respect of EC content in 70 wt.% [PEMA/PVdF–HFP]–30 wt.% LiTf polymer | 78 |

| | | |
|-------------|---|----|
| | electrolytes | |
| Figure 4.21 | FTIR spectra between (a) 3800–2800 cm^{-1} and (b) 1850–650 cm^{-1} of i. PC–0, ii. PC–2, iii. PC–4, iv. PC–6, v. PC–8, vi. PC–10 and vii. PC | 81 |
| Figure 4.22 | FTIR spectra in the region 1850–1600 cm^{-1} for (a) PC–0, (b) PC–2, (c) PC–4, (d) PC–6, (e) PC–8, (f) PC–10, (g) 90 wt.% PEMA–10 wt.% PC, (h) 90 wt.% PVdF–HFP–10 wt.% PC, (i) S–0–PC, (j) PC and (k) 6 wt.% LiTf–94 wt.% PC | 82 |
| Figure 4.23 | IR spectra in the wavenumber region between 1850 and 1700 cm^{-1} of (a) PC and (b) 6 wt.% LiTf–94 wt.% PC | 83 |
| Figure 4.24 | Deconvoluted FTIR spectra of (a) PC–0, (b) PC, (c) PC–2, (d) PC–4, (e) PC–6, (f) PC–8 and (g) PC–10 | 84 |
| Figure 4.25 | Deconvoluted FTIR spectra of (a) PC–0, (b) PC, (c) PC–2, (d) PC–4, (e) PC–6, (f) PC–8 and (g) PC–10 | 86 |
| Figure 4.26 | Deconvoluted FTIR spectra of (a) 6 wt.% LiTf–94 wt.% PC, (b) 90 wt.% PEMA–10 wt.% PC, (c) 90 wt.% PVdF–HFP–10 wt.% PC (between 1320 and 1120 cm^{-1}) and (d) 90 wt.% S–0–10 wt.% PC in the IR region between 1220 and 1120 cm^{-1} | 87 |
| Figure 4.27 | Schematic diagram of possible interactions in PEMA/PVdF–HFP–LiTf–PC system (where $\bullet\bullet\bullet$ represents coordinate bonds with Li^+ ions, $\bullet\bullet\bullet$ represents electrostatic interactions between Li^+ and PC and $\blacksquare\blacksquare\blacksquare$ represents intermolecular interactions between PC and PEMA/PVdF–HFP) | 88 |
| Figure 4.28 | IR deconvoluted bands in the region 1080–980 cm^{-1} for (a) PC–2, (b) PC–4, (c) PC–6, (d) PC–8 and (e) PC–10 | 89 |
| Figure 4.29 | Area % of free ions and ion pairs with respect to PC content in 70 wt.% [PEMA/PVdF–HFP]–30 wt.% LiTf polymer electrolytes | 90 |
| Figure 4.30 | FTIR spectra between (a) 3800–2800 cm^{-1} and (b) 1850–650 cm^{-1} of i. BI–0, ii. BI–5, iii. BI–10, iv. BI–12.5, v. BI–15, vi. BI–17.5, vii. BI–20 and viii. BMII | 93 |
| Figure 4.31 | FTIR spectra in the region 1850–1600 cm^{-1} for (a) BI–0, (b) BI–5, (c) BI–10, (d) BI–12.5, (e) BI–15, (f) BI–17.5, (g) BI–20, (h) 90 wt.% PEMA–10 wt.% BMII, (i) 90 wt.% PVdF–HFP–10 wt.% BMII and (j) S–0–BMII and (k) BMII | 94 |

| | | |
|-------------|---|-----|
| Figure 4.32 | FTIR spectra in the region between 1340 and 1200 cm^{-1} of (a) BI-0, (b) BI-5, (c) BI-10, (d) BI-12.5, (e) BI-15, (f) BI-17.5 and (g) BI-20 | 95 |
| Figure 4.33 | FTIR spectra in the region between 1220 and 1120 cm^{-1} of (a) BI-0, (b) BMII, (c) BI-5, (d) BI-10, (e) BI-12.5, (f) BI-15, (g) BI-17.5 and (h) BI-20 | 96 |
| Figure 4.34 | FTIR spectra in the region between 1220 and 1120 cm^{-1} of (a) 90 wt.% PEMA-10 wt.% BMII and (b) 90 wt.% S-0-10 wt.% BMII | 97 |
| Figure 4.35 | Schematic diagram of possible interactions in PEMA/PVdF-HFP-LiTf-BMII system (where $\bullet\bullet\bullet$ represents coordinate bonds with Li^+ ions and $\blacksquare\blacksquare\blacksquare$ represents intermolecular interactions between BMII and PEMA/PVdF-HFP) | 98 |
| Figure 4.36 | Deconvolution between 1100 and 980 cm^{-1} of (a) BI-5, (b) BI-10, (c) BI-12.5, (d) BI-15 (e) BI-17.5 and (f) BI-20 | 100 |
| Figure 4.37 | Area % of free ion, ion pair and ion aggregate as a function of BMII content | 100 |
| Figure 4.38 | FTIR spectra wavenumber region (a) 3800-2800 cm^{-1} and (b) 1850-650 cm^{-1} of i. BT-0, ii. BT-10, iii. BT-20, iv. BT-30, v. BT-40, vi. BT-50, vii. BT-60 and viii. BMITf | 103 |
| Figure 4.39 | FTIR spectra in the region 1850-1600 cm^{-1} for (a) BT-0, (b) BT-10, (c) BT-20, (d) BT-30, (e) BT-40, (f) BT-50, (g) BT-60, (h) 90 wt.% PEMA-10 wt.% BMITf, (i) 90 wt.% PVdF-HFP-10 wt.% BMITf and (j) S-0-BMITf and (k) BMITf | 105 |
| Figure 4.40 | FTIR spectra in the region between 1350 and 1200 cm^{-1} of (a) BT-0, (b) BMITf, (c) BT-10, (d) BT-20, (e) BT-30, (f) BT-40, (g) BT-50 and (h) BT-60 | 106 |
| Figure 4.41 | FTIR spectra in the region between 1220 and 1120 cm^{-1} of (a) BT-0, (b) BMITf, (c) BT-10, (d) BT-20, (e) BT-30, (f) BT-40, (g) BT-50 and (h) BT-60 | 108 |
| Figure 4.42 | Schematic diagram of possible interactions in PEMA/PVdF-HFP-LiTf-BMITf system (where $\bullet\bullet\bullet$ represents coordinate bonds with Li^+ ions and $\blacksquare\blacksquare\blacksquare$ represents intermolecular interactions between BMITf and PEMA/PVdF-HFP) | 109 |
| Figure 4.43 | Deconvolution between 1080 and 980 cm^{-1} of (a) BT-10, (b) BT-20, (c) BT-30, (d) BT-40, (e) BT-50 and (f) BT-60 | 111 |

| | | |
|-------------|---|-----|
| Figure 4.44 | Area % of free ions, ion pairs and ion aggregates as a function of BMITf content | 113 |
| Figure 5.1 | Nyquist plots of (a) S-0, (b) S-5, (c) S-10, (d) S-12.5, (e) S-15, (f) S-20, (g) S-25, (h) S-30, (i) S-35, (j) S-40 films | 116 |
| Figure 5.2 | Effect of LiTf content on the ionic conductivity at 298 K of PEMA/PVdF-HFP-based polymer electrolytes | 119 |
| Figure 5.3 | Nyquist plots of S-30 at various temperatures | 121 |
| Figure 5.4 | Plot of $\log \sigma$ versus $1000/T$ of PEMA/PVdF-HFP-LiTf polymer electrolytes | 122 |
| Figure 5.5 | Plot of number density and mobility in PEMA/PVdF-HFP-LiTf system | 123 |
| Figure 5.6 | Nyquist plots of (a) EC-2, (b) EC-4, (c) EC-6, (d) EC-8 and (e) EC-10 films | 124 |
| Figure 5.7 | Effect of EC content on the ionic conductivity at 298 K of PEMA/PVdF-HFP-LiTf based polymer electrolytes | 126 |
| Figure 5.8 | Nyquist plots of EC-6 at various temperatures | 127 |
| Figure 5.9 | Plot of $\log \sigma$ versus $1000/T$ of PEMA/PVdF-HFP-LiTf-EC polymer electrolytes | 128 |
| Figure 5.10 | Plot of $\log (\sigma T^{1/2})$ versus $1000/(T-T_0)$ of PEMA/PVdF-HFP-LiTf-EC polymer electrolytes | 128 |
| Figure 5.11 | Plot of number density and mobility in PEMA/PVdF-HFP-LiTf-EC system | 129 |
| Figure 5.12 | Nyquist plots of (a) PC-2, (b) PC-4, (c) PC-6, (d) PC-8 and (e) PC-10 films | 130 |
| Figure 5.13 | Effect of PC content on the ionic conductivity at 298 K of PEMA/PVdF-HFP-LiTf based polymer electrolytes | 132 |
| Figure 5.14 | Nyquist plots of PC-6 at various temperatures | 133 |
| Figure 5.15 | Plot of $\log \sigma$ versus $1000/T$ of PEMA/PVdF-HFP-LiTf-PC polymer electrolytes | 134 |
| Figure 5.16 | Plot of $\log (\sigma T^{1/2})$ versus $1000/(T-T_0)$ of PEMA/PVdF-HFP-LiTf-PC polymer electrolytes | 134 |

| | | |
|-------------|---|-----|
| Figure 5.17 | Plot of number density and mobility in PEMA/PVdF–HFP–LiTf–PC system | 135 |
| Figure 5.18 | Nyquist plots of (a) BI–5, (b) BI–10, (c) BI–12.5, (d) BI–15, (e) BI–17.5 and (f) BI–20 films | 136 |
| Figure 5.19 | Effect of BMII content on the ionic conductivity at 298 K of PEMA/PVdF–HFP–LiTf based polymer electrolytes | 137 |
| Figure 5.20 | Nyquist plots of BI–12.5 at various temperatures | 138 |
| Figure 5.21 | Plot of $\log \sigma$ versus $1000/T$ of PEMA/PVdF–HFP–LiTf–BMII polymer electrolytes | 140 |
| Figure 5.22 | Plot of number density and mobility in PEMA/PVdF–HFP–LiTf–BMII system | 141 |
| Figure 5.23 | Nyquist plots of (a) BT–10, (b) BT–20, (c) BT–30, (d) BT–40, (e) BT–50 and (f) BT–60 films | 142 |
| Figure 5.24 | Plot of $\log \sigma$ vs BMITf content at 298 K | 144 |
| Figure 5.25 | Nyquist plots of BT–60 at various temperatures | 145 |
| Figure 5.26 | Plot of $\log \sigma$ versus $1000/T$ of PEMA/PVdF–HFP–LiTf–BMITf polymer electrolytes | 146 |
| Figure 5.27 | Plot of number density and mobility in PEMA/PVdF–HFP–LiTf–BMITf system | 147 |
| Figure 6.1 | TGA plots of PEMA, PVdF–HFP and S–0 | 150 |
| Figure 6.2 | Decomposition of PEMA through (a) head–to–head linkage scission, (b) scission at vinylidene chain–end units and (c) random scission | 152 |
| Figure 6.3 | Formation of EMA monomers by depropagation and termination of all radicals | 153 |
| Figure 6.4 | Decomposition of PVdF–HFP by random scission by different pathways <i>a</i> and <i>b</i> | 154 |
| Figure 6.5 | Decomposition of LiTf [Taken from Ohtani et al., 2008] | 155 |
| Figure 6.6 | TGA plots of PEMA/PVdF–HFP–LiTf polymer electrolytes | 156 |
| Figure 6.7 | TGA plots of PEMA/PVdF–HFP–LiTf–EC polymer electrolytes | 158 |
| Figure 6.8 | TGA plots of PEMA/PVdF–HFP–LiTf–PC polymer electrolytes | 160 |

| | | |
|-------------|--|-----|
| Figure 6.9 | TGA plots of PEMA/PVdF–HFP–LiTf–BMII polymer electrolytes | 163 |
| Figure 6.10 | Thermal decomposition pathways of BMII | 164 |
| Figure 6.11 | TGA plots of PEMA/PVdF–HFP–LiTf–BMITf polymer electrolytes | 167 |
| Figure 6.12 | Thermal decomposition mechanisms of (a) 1-butyl-3-methylimidazolium and (b) triflate of BMITf [Taken from Ohtani et al., 2008] | 168 |
| Figure 7.1 | Possible ion transport mechanism in PEMA/PVdF–HFP–LiTf–EC system involving the formation of ionic clusters | 182 |