

## REFERENCES

- Abbott, I. A. (1988). Introduction. In I. A. Abbott (Ed.), *Taxonomy of economic seaweeds, with reference to some Pacific and Caribbean species: Vol. II* (pp. 127-129): California Sea Grant College Program, University of California.
- Abbott, I. A. (1994). New records and a reassessment of *Gracilaria* (Rhodophyta) from the Philippines. *Taxonomy of economic seaweeds with reference to some Pacific species*, 4.
- Abbott, I. A., Zhang, J., & Xia, B. M. (1991). *Gracilaria mixta*, sp. nov. and other western Pacific species of the genus (Rhodophyta: Gracilariaceae). *Pacific Science*, 45, 12-27.
- Adams, R. I., Brown, K. M., & Hamilton, M. B. (2004). The impact of microsatellite electromorph size homoplasy on multilocus population structure estimates in a tropical tree (*Corythophora alta*) and an anadromous fish (*Morone saxatilis*). *Molecular Ecology*, 13, 2579-2588.
- Agardh, J. G. (1852). *Species genera et ordines algarum, seu descriptiones succinctae specierum, generum et ordinum, quibus algarum regnum constituitur. Volumis secundi: Algas florideas complectens. 2: 1.* pp. 701-786.
- Aggarwal, R., Sharma, V., Kharbikar, L. L., & Renu. (2008). Molecular characterization of *Chaetomium* species using URP-PCR. *Genetics and Molecular Biology*, 31, 943-946.
- Andreakis, N., Kooistra, W. H. C. F., & Procaccini, G. (2007). Microsatellite markers in an invasive strain of *Asparagopsis taxiformis* (Bonnemaisoniales, Rhodophyta): insights in ploidy level and sexual reproduction. *Gene*, 406, 144-151.
- Armisen, R. (1995). World-wide use and importance of *Gracilaria*. *Journal of Applied Phycology*, 7, 231-243.
- Armour, J. A. L., Alegre, S. A., Miles, S., Williams, L. J., & Badge, R. M. (1999). Minisatellites and mutation processes in tandemly repetitive DNA. In D. B. Goldstein & C. Schlötterer (Eds.), *Microsatellites: evolution and applications* (pp. 24-33). Oxford: Oxford University Press.

- Ashley, M. V., & Dow, B. D. (1994). The use of microsatellite analysis in population biology: background, methods and potential applications. *EXS*, 69, 185-201.
- Asif, M., Mehboob-ur-Rahman, Mirza, J. I., & Zafar, Y. (2008). High resolution metaphor agarose gel electrophoresis for genotyping with microsatellite markers. *Pakistan Journal of Agricultural Sciences*, 45, 75-79.
- Awasthi, D. K. (2009). *Cryptogams: Algae, Bryophyta and Pterldophyta*. Meerut, Delhi: Krishna Prakashan Media.
- Ayeh, K. O. (2008). Expressed sequence tags (ESTs) and single nucleotide polymorphisms (SNPs): Emerging molecular marker tools for improving agronomic traits in plan biotechnology. *African Journal of Biotechnology*, 7, 331-341.
- Badger, J. H., Zhang, T., Fang, Y., Wang, X., Deng, X., Zhang, X., . . . Yu, J. (2012). The complete chloroplast and mitochondrial genome sequences of *Boea hygrometrica*: Insights into the evolution of plant organellar genomes. *PLoS ONE*, 7, e30531.
- Bao, S., Corke, H., & Sun, M. (2002). Microsatellites in starch-synthesizing genes in relation to starch physicochemical properties in waxy rice (*Oryza sativa* L.). *Theoretical and Applied Genetics*, 105, 898-905.
- Bartsch, I., Wiencke, C., Bischof, K., Buchholz, C. M., Buck, B. H., Eggert, A., . . . Wiese, J. (2008). The genus *Laminaria* sensu lato: recent insights and developments. *European Journal of Phycology*, 43, 1-86.
- Barufi, J. B., Cabral de Oliveira, E., Plastino, E. M., & Cabral de Oliveira, M. (2010). Life history, morphological variability and growth rates of the life phases of *Gracilaria tenuistipitata* (Rhodophyta: Gracilariales) in vitro. *Scientia Marina*, 74, 297-303.
- Batley, J., Barker, G., Sullivan, H. O., Edwards, K. J., & Edwards, D. (2003). Mining for single nucleotide polymorphisms and insertions/deletions in maize expressed sequence tag data. *Plant Physiology*, 132, 84-91.
- Beckman, J. S., & Weber, J. L. (1992). Survey of human and rat microsatellites. *Genomics*, 12, 627-631.

- Beebe T. & Rowe, G. (2008). *An Introduction to Molecular Ecology*, 2nd ed. New York, NY: Oxford University Press.
- Bell, P. R., & Hemsley, A. R. (2000). *Green Plants: Their Origin and Diversity*. U. K.: Cambridge University Press.
- Bellorin, A. M., Buriyo, A., Sohrabipour, J., Oliveira, M. C., & Oliveira, E. C. (2008). *Gracilariopsis mclachlanii* sp. nov. and *Gracilariopsis persica* sp. nov. of the Gracilariaceae (Gracilariales, Rhodophyceae) from the Indian Ocean. *Journal of Phycology*, 44, 1022-1032.
- Bellorin, A. M., Oliveira, M. C., & Oliveira, E. C. (2002). Phylogeny and systematics of the marine algal family Gracilariaceae (Gracilariales, Rhodophyta) based on small subunit rDNA and ITS sequences of Atlantic and Pacific species. *Journal of Phycology*, 38, 551-563.
- Bhattacharya, D., Elwood, H. J., Goff, L. J., & Sogin, M. L. (1990). Phylogeny of *Gracilaria lemaneiformis* (Rhodophyta) based on sequence analysis of its small subunit ribosomal RNA coding region. *Journal of Phycology*, 26, 181-186.
- Bhattacharya, D., & Medlin, L. (1998). Algal phylogeny and the origin of land plants. *Plant Physiology*, 116, 9-15.
- Billot, C., Rousvoal, S., Estoup, A., Epplen, J. T., Saumitou-Laprade, P., Valero, M., & Kloareg, B. (1998). Isolation and characterization of microsatellite markers in the nuclear genome of the brown alga *Laminaria digitata* (Phaeophyceae). *Molecular Ecology*, 7, 1778-1780.
- Bird, C. J., Rice, E. L., Murphy, C. A., & Ragan, M. A. (1992). Phylogenetic relationships in the Gracilariales (Rhodophyta) as determined by 18S rDNA sequences. *Phycologia*, 31, 510-522.
- Bird, C. J., & van der Meer, J. P. (1993). Systematics of economically important marine algae: a Canadian perspective. *Canadian Journal of Botany*, 71, 361-369.
- Bixler, H. J., & Porse, H. (2010). A decade of change in the seaweed hydrocolloids industry. *Journal of Applied Phycology*, 23(3), 321-335.

- Botstein, D., White, R. L., Skolnick, M., & Davis, R. W. (1980). Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American Journal of Human Genetics*, 32, 314-331.
- Bouck, A., & Vision, T. (2007). The molecular ecologist's guide to expressed sequence tags. *Molecular Ecology*, 16, 907-924.
- Brasch, D. J., Chang, H. M., Chuah, C. T., & Melton, L. D. (1981). The galactan sulfate from the edible, red alga *Porphyra columbina*. *Carbohydrate Research*, 97, 113-125.
- Brauckmann, B. M. (1990). Industrial solution amenable to biosorption. In B. Volesky (Ed.), *Biosorption or heavy metals* (pp. 52-60). Boca Raton, Florida: CRC Press.
- Bremer, K., Humphries, C. J., Mishler, B. D., & Churchill, S. P. (1987). On cladistic relationships in green plants. *Taxon*, 36, 339-349.
- Brodie, J. A., & Irvine, L. M. (2003). *Seaweeds of the British Isles: Volume 1 Rhodophyta Part 3B Bangiophycidae*: Intercept, Hampshire, UK.
- Buchholz, C., Krause, G., & Buck, B. H. (2012). Seaweed and Man. In C. Wiencke & K. Bischof (Eds.), *Seaweed Biology: Novel Insights into Ecophysiology, Ecology and Utilization* (pp. 509). Heidelberg, Berlin: Springer.
- Burger, G., Saint-Louis, D., Gray, M. W., & Lang, B. F. (1999). Complete sequence of the mitochondrial DNA of the red alga *Porphyra purpurea*: cyanobacterial introns and shared ancestry of red and green algae. *The Plant Cell*, 11, 1675-1694.
- Butterfield, N. J. (2000). *Bangiomorpha pubescens* n. gen., n. sp.: implications for the evolution of sex, multicellularity, and the Mesoproterozoic-Neoproterozoic radiation of eukaryotes. *Paleobiology*, 26, 386-404.
- Caetano-Anollés, G., Bassam, B. J., & Gresshoff, P. M. (1992). Primer-template interactions during DNA amplification fingerprinting with single arbitrary oligonucleotides. *Molecular and General Genetics*, 235, 157-165.

- Callen, D. F., Thompson, A. D., Shen, Y., Phillips, H. A., Richards, R. I., Mulley, J. C., & Sutherland, G. R. (1993). Incidence and origin of "null" alleles in the (AC)<sub>n</sub> microsatellite markers. *The American Journal of Human Genetics*, 52, 922-927.
- Canovas, F., Mota, C., Ferreira-Costa, J., Serrão, E., Coyer, J., Olsen, J., & Pearson, G. (2011). Development and characterization of 35 single nucleotide polymorphism markers for the brown alga *Fucus vesiculosus*. *European Journal of Phycology*, 46, 342-351.
- Casas-Valdez, M., Portillo-Clark, G., Aguila-Ramirez, N., Rodriguez-Astudillo, S., Sanchez-Rodriguez, I., & Carillo-Dominguez, S. (2006). Effect of the marine algae *Sargassum* spp. on the productive parameters and cholesterol content of the brown shrimp, *Farfantepenaeus californiensis* (Holmes, 1900). *Revista de Biología Marina y Oceanografía*, 41, 97-105.
- Chambers, G. K., Cordiner, S. J., Buckleton, J. S., Robertson, B., & Vignaux, G. A. (1997). Forensic DNA profiling: The importance of giving accurate answers to the right questions. *Criminal Law Forum*, 8, 445-459.
- Chambers, G. K., & MacAvoy, E. S. (2000). Microsatellites: consensus and controversy. *Comparative Biochemistry and Physiology - Part B*, 126, 455-476.
- Chan, C. X., Ho, C. L., & Phang, S. M. (2006). Trends in seaweed research. *Trends in plant science*, 11, 165-166.
- Chang, C. F., & Xia, B. M. (1963). *Polycavernosa*, a new genus in the Gracilariaceae. *Studia Marina Sinica*, 3, 119-126.
- Chang, C. F., & Xia, B. M. (1988). On two new *Gracilaria* (Gigartinales, Rhodophyta) from South China. In I. A. Abbott (Ed.), *Taxonomy of economic seaweeds with reference to some Pacific and Caribbean species* (Vol. 2, pp. 127-129): California Sea Grant College Program.
- Chapman, V.J. (1970). *Seaweeds and their uses*. Methuen, London. pp. 304.
- Chase, M., Salamin, N., Wilkinson, M., Dunwell, J., Kesanakurthi, R., Haidar, N., & Savolainen, V. (2005). Land plants and DNA barcodes: short-term and long-term goals. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 360, 1889-1895.

- Cardlea, L., Ramsaya, L., Milbourne, D., Macaulaya, M., Marshalla, D., & Waugh, R. (2000). Computational and experimental characterization of physically clustered simple sequence repeats in plants. *Genetics*, *156*, 847-854.
- Cheney, D. & Kurtzman, A. (1992). Progress in protoplast fusion and gene transfer in red algae. *Proceedings of the Fourteenth International Seaweed Symposium*, *14*, 68.
- Cheng, Y. J., Meng, H. J., Guo, W. W., & Deng, X. X. (2006). Universal chloroplast primer pairs for simple sequence repeat analysis in diverse genera of fruit crops. *Journal of Horticultural Science & Biotechnology*, *81*, 132-138.
- Chesnick, J. M., & Cattolico, R. A. (1993). Isolation of DNA from eucaryotic algae. *Methods in Enzymology*, *224*, 168-176.
- Ching, A., Caldwell, K. S., Jung, M., Dolan, M., Smith, O. S. H., Tingey, S., . . . Rafalski, A. J. (2002). SNP frequency, haplotype structure and linkage disequilibrium in elite maize inbred lines. *BMC Genetics*, *3*, 1-14.
- Cho, Y. G., Ishii, T., Temnykh, S., Chen, X., Lipovich, L., McCouch, S. R., . . . Cartinhour, S. (2000). Diversity of microsatellites derived from genomic libraries and GenBank sequences in rice (*Oryza sativa* L.). *Theoretical and Applied Genetics*, *100*, 713-722.
- Chung, S. M., & Staub, J. E. (2003). The development and evaluation of consensus chloroplast primer pairs that possess highly variable sequence regions in a diverse array of plant taxa. *Theoretical and Applied Genetics*, *107*, 757-767.
- Cipriani, G., Lot, G., Huang, W. G., Marrazzo, M. T., Peterlunger, E., & Testolin, R. (1999). AC/GT and AG/CT microsatellite repeats in peach [*Prunus presica* (L) batsch]: isolation, characterisation and cross-species amplification in *Prunus*. *Theoretical and Applied Genetics*, *99*, 65-72.
- Coleman, A. W., & Goff, L. J. (1991). DNA analysis of eukaryotic algal species. *Journal of Phycology*, *27*, 463-473.
- Coll n, J., Pinto, E., Peders n, M., & Colepicolo, P. (2003). Induction of Oxidative Stress in the Red Macroalga *Gracilaria tenuistipitata* by Pollutant Metals. *Archives of Environmental Contamination and Toxicology*, *45*, 337-342.

- Collín, P. N., Collín, J., Reis, M. S., Pedersen, M., Setubal, J. C., Varani, A. M., . . . Oliveira, M. C. (2012). Analysis of expressed sequence tags from the agarophyte *Gracilaria tenuistipitata* (Rhodophyta). *Journal of Applied Phycology*, *24*, 641-647.
- Cordeiro, G. M., Casu, R., McIntyre, C. L., Manners, J. M., & Henry, R. J. (2001). Microsatellite markers from sugarcane (*Saccharum* spp.) ESTs cross transferable to erianthus and sorghum. *Plant Science*, *160*, 1115-1123.
- Corley-Smith, G. E., Lim, C. J., Kalmar, G. B., & Brandhorst, B. P. (1997). Efficient detection of DNA polymorphisms by fluorescent RAPD analysis. *Biotechniques*, *22*, 690-692.
- Cosmetic Ingredient Dictionary (2002–2011) Algae.  
(<http://www.cosmeticscop.com/cosmetic-ingredient-dictionary/A.aspx>)
- Coyer, J. A., Hoarau, G., Beszteri, B., Pearson, G., & Olsen, J. L. (2008). Expressed sequence tag-derived polymorphic SSR markers for *Fucus serratus* and amplification in other species of *Fucus*. *Molecular Ecology Resources*, *9*, 168-170.
- Coyer, J. A., Veldsink, J. H., Jones, K., Stam, W. T., & Olsen, J. L. (2002). Characterization of microsatellite loci in the marine seaweeds, *Fucus serratus* and *F. evanescens* (Heterokontophyta; Fucaceae). *Molecular Ecology Notes*, *2*, 35-37.
- Craig, J., Fowler, S., Burgoyne, L. A., Scott, A. C., & Harding, H. W. J. (1988). Repetitive deoxyribonucleic acid (DNA) and human genome variation: A concise review relevant to forensic biology. *Journal of Forensic Sciences*, *33*, 1111-1126.
- Crossland, S., Coates, D., Grahame, J., & Mill, P. J. (1993). Use of random amplified polymorphic DNAs (RAPDs) in separating two sibling species of *Littorina*. *Marine Ecology Progress Series*, *96*, 301-305.
- Damonte, E. B., Neyts, J., Pujol, C. A., Snoeck, R., Andrei, G., Ikeda, S., . . . De Clercq, E. (1994). Antiviral activity of a sulphated polysaccharide from the red seaweed *Nothogenia fastigiata*. *Biochemical Pharmacology*, *47*, 2187-2192.
- Dawson, E. Y. (1949). Studies of the northeast Pacific Gracilariaceae. *Occasional Papers of the Allan Hancock Foundation*, *7*, 1-105.

- De Clercq, E. (1996). Chemotherapy of human immunodeficiency virus (HIV) infection: anti-HIV agents targeted at early stages in the virus replicative cycle. *Biomedicine & Pharmacotherapy*, 50, 207-215.
- Dellaporta, S. L., Wood, V. P., & Hicks, J. B. (1983). A plant DNA mini-preparation: version II. *Plant Molecular Biology Reporter*, 1, 19-21.
- De Ruiter, G. A., & Rudolph, B. (1997). Carrageenan biotechnology. *Trends in Food Science and Technology*, 8, 389-395.
- de Senerpont Domis, L. N., Famà P., Bartlett, A. J., Prud'homme van reine, W. F., Espinosa, C. A., & Trono Jr., G. C. (2003). Defining taxon boundaries in members of the morphologically and genetically plastic genus *Caulerpa* (Caulerpales, Chlorophyta). *Journal of Phycology*, 39, 1019-1037.
- Destombe, C., & Douglas, S. E. (1991). Rubisco spacer sequence divergence in the rhodophyte alga *Gracilaria verrucosa* and closely related species. *Current Genetics*, 19, 395-398.
- DeVerno, L. L., & Mosseler, A. (1997). Genetic variation in red pine (*Pinus resinosa*) revealed by RAPD and RAPD-RFLP analysis. *Canadian Journal of Forest Research*, 27, 1316-1320.
- Dhargalkar, V. K., & Kavlekar, D. (2004). Seaweeds - A field manual. National Institute of Oceanography, Dona Paula, Goa. pp. 36.
- Dillehay, T. D., Ramírez, C., Pino, M., Collins, M. B., Rossen, J., & Pino-Navarro, J. D. (2008). Monte Verde: Seaweed, Food, Medicine, and the Peopling of South America. *Science*, 320, 784-786.
- Dixon, P. S. (1973). *Biology of the Rhodophyta*: Oliver and Boyd, Edinburgh.
- Donaldson, S. L., Chopin, T., & Saunders, G. W. (1998). Amplified fragment length polymorphism (AFLP) as a source of genetic markers for red algae. *Journal of Applied Phycology*, 10, 365-370.
- Doty, M. S. (1986). Estimating farmer returns from producing *Gracilaria* and *Eucheuma* on line farms. *Monografias Biologicas*, 4, 45-62.



- Doyle, J. J., & Doyle, J. (1990). Isolation of plant DNA from fresh tissue. *Focus*, *12*, 13-15.
- Dresselhaus, T., Cordts, S., Heuer, S., Sauter, M., Lörz, H., & Kranz, E. (1999). Novel ribosomal genes from maize are differentially expressed in the zygotic and somatic cell cycles. *Molecular and General Genetics*, *261*, 416-427.
- Drew, K. M. (1954). Studies in the Bangioideae III. The Life-history of *Porphyra umbilicalis* (L.) Kütz. var. *lociniata* (Lightf.) J. Ag. A. The Conchocelis-Phase in culture. *Annals of Botany*, *70*, 184-211.
- Ebert, D., & Peakall, R. O. D. (2009). Chloroplast simple sequence repeats (cpSSRs): technical resources and recommendations for expanding cpSSR discovery and applications to a wide array of plant species. *Molecular Ecology Resources*, *9*, 673-690.
- Echt, C. S., DeVerno, L. L., Anzidei, M., & Vendramin, G. G. (1998). Chloroplast microsatellites reveal population genetic diversity in red pine, *Pinus resinosa* Ait. *Molecular Ecology*, *7*, 307-316.
- Ellis, J. R., & Burke, J. M. (2007). EST-SSRs as a resource for population genetic analyses. *Heredity*, *99*, 125-132.
- Eujayl, I., Sorrells, M., Baum, M., Wolters, P., & Powell, W. (2001). Assessment of genotypic variation among cultivated durum wheat based on EST-SSRs and genomic SSRs. *Euphytica*, *119*, 39-43.
- Fain, S. R., Druehl, L. D., & Baillie, D. L. (1988). Repeat and single copy sequences are differentially conserved in the evolution of kelp chloroplast DNA. *Journal of Applied Phycology*, *24*, 292-302.
- Fama, P., Jousson, O., Zaninetti, L., Meinesz, A., Dini, F., Di Guiseppe, G., . . . Pawlowski, J. (2002). Genetic polymorphism in *Caulerpa taxifolia* (Ulvophyceae) chloroplast DNA revealed by a PCR-based assay of the invasive Mediterranean strain. *Journal of Evolutionary Biology*, *15*, 618-624.
- Fawley, M. W., & Fawley, K. P. (2004). A simple and rapid technique for the isolation of DNA from microalgae. *Phycological Society of America*, *40*, 223-225.

- FAO (2001). The state of world fisheries and aquaculture 2000, *Electronic edition*. <http://fao.org>.
- FAO (2011a) FAO Fisheries Department, Fishery Information, Data and Statistics Unit. FishStatPlus. Universal Software for fishery statistical time series. Version 2.3 in 2000.
- FAO (2012). The state of world fisheries and aquaculture 2012. *Electronic edition*, <http://fao.org>.
- Fleurence, J. (1999). Seaweed proteins: biochemical, nutritional aspects and potential uses. *Trends in Food Science and Technology*, 10, 25-28.
- Fraser, L. G., Harvey, C. F., Crowhurst, R. N., & De Silva, H. N. (2004). EST-derived microsatellites from *Actinidia* species and their potential for mapping. *Theoretical and Applied Genetics*, 108, 1010-1016.
- Fredericq, S., & Hommersand, M. H. (1989). Comparative morphology and taxonomic status of *Gracilariopsis* (Gracilariales, Rhodophyta). *Journal of Phycology*, 25, 228-241.
- Fujiwara-Arasaki, T., Mino, N., & Kuroda, M. (1984). The protein value in human nutrition of edible marine algae in Japan. *Hydrobiologia*, 116-117, 513-516.
- Gan, S. Y., Qin, S., Rofina, Y. O., Yu, D., & Phang, S. M. (2003). Transient expression of lacZ in particle bombarded *Gracilaria changii* (Gracilariales, Rhodophyta). *Journal of Applied Phycology*, 15, 345-349.
- Gao, L., Tang, J., Li, H., & Jia, J. (2003). Analysis of microsatellites in major crops assessed by computational and experimental approaches. *Molecular Breeding*, 12, 245-261.
- Gargiulo, G. M., Morabito, M., Genoves, G., & De Masi, F. (2006). Molecular systematics and phylogenetics of Gracilariacean species from the Mediterranean Sea. *Journal of Applied Phycology*, 18, 497-504.

- Goff, L. J., & Coleman, A. W. (1988). The use of plastid DNA restriction endonuclease patterns in delineating red algal species and populations. *Journal of Phycology*, *24*, 357-368.
- González, M., Montoya, R., Candia, A., Gómez, P., & Cisternas, M. (1996). Organellar DNA restriction fragment length polymorphism (RFLP) and nuclear random amplified polymorphic DNA (RAPD) analyses of morphotypes of *Gracilaria* (Gracilariales, Rhodophyta) from Chile. *Hydrobiologia*, *326-327*, 229-234.
- Graham, L. E., & Wilcox, L. W. (2000). *The algae*. New York: Prentice Hall.
- Greville, R. K. (1830). *Algae Britannicae*. *Edinburgh*. pp. 1-218.
- Grodzicker, T., Williams, J., Sharp, P., & Sambrook, J. (1975). Physical mapping of temperature-sensitive mutations of adenoviruses. *Cold Spring Harbor Symposia on Quantitative Biology*, *39*, 439-446.
- Guillemin, M. L., Destombe, C., Faugeron, S., Correa, J. A., & Valero, M. (2005). Development of microsatellites DNA markers in the cultivated seaweed, *Gracilaria chilensis* (Gracilariales, Rhodophyta). *Molecular Ecology Notes*, *5*, 155-157.
- Gupta, P. K., Rustgi, S., Sharma, S., Singh, R., Kumar, N., & Balyan, H. S. (2003). Transferable EST-SSR markers for the study of polymorphism and genetic diversity in bread wheat. *Molecular Genetics and Genomics*, *270*, 315-323.
- Gurgel, C. F. D., & Fredericq, S. (2004). Systematics of the Gracilariaceae (Gracilariales, Rhodophyta): a critical assessment based on *rbcL* sequence analyses. *Journal of Phycology*, *40*, 138-159.
- Hackauf, B., & Wehling, P. (2002). Identification of microsatellite polymorphisms in an expressed portion of the rye genome. *Plant Breeding*, *121*, 17-25.
- Hagopian, J. C., Reis, M., Kitajima, J. P., Bhattacharya, D., & de Oliveira, M. C. (2004). Comparative analysis of the complete plastid genome sequence of the red alga *Gracilaria tenuistipitata* var. *liui* provides insights into the evolution of rhodoplasts and their relationship to other plastids. *Journal of Molecular Evolution*, *59*, 464-477.

- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, 41, 95-98.
- Han, Z. G., Guo, W. Z., Song, X. L., & Zhang, T. Z. (2004). Genetic mapping of EST-derived microsatellites from the diploid *Gossypium arboreum* in allotetraploid cotton. *Molecular Genetics and Genomics*, 272, 308-327.
- Harris, E. H. (1989). *The Chlamydomonas sourcebook. A Comprehensive Guide to Biology and Laboratory Use*. San Diego: Academic Press.
- Hawkes, M. W. (1990). Reproductive strategies. In K. M. Cole & R. G. Sheath (Eds.), *Biology of the red algae* (pp. 455-476). New York: Cambridge University Press.
- Helentjaris, T., Slocum, M., Wright, S., Schaefer, A., & Nienhuis, J. (1986). Construction of genetic linkage maps in maize and tomato using restriction fragment length polymorphisms. *Theoretical and Applied Genetics*, 72, 761-769.
- Hepperle, D. (2002). Fingerprinter©. A software for the analysis, bin-assignment and 01-matrix construction of molecular fingerprints (AFLPs). *Win32-Version*. Distributed by the author via: <http://science.do-mix.de>.
- Hepperle, D. (2004). TreeMe©. A software for visualization, manipulation, layouting and labelling of phylogenetic trees. *Win32-Version*. Distributed by the author via: <http://www.sequentix.de>.
- Hirschfeld, D., Dhar, A. K., Rask, K., & Alcivar-Warren, A. (1999). Genetic diversity in the eastern oyster (*Crassostrea virginica*) from Massachusetts using RAPD technique. *Journal of Shellfish Research*, 18, 121-125.
- Hong, Y. K., Coury, D. A., Polne-Fuller, M., & Gibor, A. (1992). Lithium chloride extraction of DNA from the seaweed *Porphyra perforata* (Rhodophyta). *Journal of Phycology*, 28, 717-720.
- Hong, Y. K., Sohn, C. H., Lee, K. W., & Kim, H. G. (1997). Nucleic acid extraction conditions from seaweed tissues for polymerase chain reaction. *Journal of Marine Biotechnology*, 5, 95-99.

- Hori, H., Lim, B. L., & Osawa, S. (1985) Evolution of green plants as deduced from 5S rRNA sequences. *Proceedings of the National Academy of Sciences of the USA*, 82, 820–823.
- Hori, H., & Osawa, S. (1987). Origin and evolution of organisms as deduced from 5S ribosomal RNA sequences. *Molecular Biology and Evolution*, 4, 445-472.
- Huang, X., & Madan, A. (1999). CAP3: A DNA sequence assembly program. *Genome Research*, 9, 868-877.
- Huff, D. R., Peakall, R., & Smouse, P. E. (1993). RAPD variation within and among natural populations of outcrossing buffalograss [*Buchloë dactyloides* (Nutt.) Engelm.]. *Theoretical and Applied Genetics*, 86, 927-934.
- Hughes, C. R., & Queller, D. C. (1993). Detection of highly polymorphic microsatellite loci in a species with little allozyme polymorphism. *Molecular Ecology*, 2, 131-137.
- Hunt, G. J., & Page Jr., R. E. (1992). Patterns of inheritance with RAPD molecular markers reveal novel types of polymorphism in the honey bee. *Theoretical and Applied Genetics*, 85, 15-20.
- Indergaard, M., & Østgaard, K. (1991). Polysaccharides for food and pharmaceutical uses. In M. D. Guiry & G. Blunden (Eds.), *Seaweed resources in Europe. Uses and potential*. (pp. 169-183). Chichester: Wiley.
- Isabel, N., Beaulieu, J., Thériault, P., & Bousquet, J. (1999). Direct evidence for biased gene diversity estimates from dominant random amplified polymorphic DNA (RAPD) fingerprints. *Molecular Ecology*, 8, 477-483.
- Iyer, R., Clerck, O., Bolton, J., & Coyne, V. (2005). Morphological and taxonomic studies of *Gracilaria* and *Gracilariopsis* species (Gracilariales, Rhodophyta) from South Africa. *South African Journal of Botany*, 70, 521-539.
- Jacob, H. J., Linderpaintner, K., Lincoln, S. E., Kusumi, K., Bunker, R. K., Mao, Y. P., . . . Lander, E. S. (1991). Genetic Mapping of a gene causing hypertension in the stroke-prone spontaneously rat. *Cell*, 67, 213-224.

- Jander, G., Norris, S. R., Rounsley, S. D., Bush, D. F., Levin, I. M., & Last, R. L. (2002). Arabidopsis Map-Based Cloning in the Post-Genome Era. *Plant Physiology*, 129, 440-450.
- Jin, H.-J., Kim, J. H., Sohn, C. H., DeWreede, R. E., Choi, T. J., Towers, G. H. N., . . . Hong, Y. K. (1997). Inhibition of Taq DNA polymerase by seaweed extracts from British Columbia, Canada and Korea. *Journal of Applied Phycology*, 9, 383-388.
- Jorgensen, R. A., Cuellar, R. E., Thompson, W. F., & Kavanagh, T. A. (1987). Structure and variation in ribosomal RNA genes of pea. *Plant Molecular Biology Reporter*, 8, 3-12.
- John, D. M., & Maggs, C. A. (1997). Species problems in eukaryotic algae: a modern perspective. In M. F. Claridge, H. A. Dawah & M. R. Wilson (Eds.), *The Units of Biodiversity: Species in Practice* (pp. 83-107). London: Chapman and Hall.
- Joshi, S. P., Ranjekar, P. K. & Gupta, V. S. (1999). Molecular markers in plant genome analysis. *Current Science*, 77, 230-240.
- Kang, H.-W., Park, D.-S., Park, Y.-J., Lee, B.-M., Cho, S.-M., Kim, K.-T., . . . Go, S.-J. (2002). PCR based detection of *Phellinus linteus* using specific primers generated from Universal Rice Primer (URP) derived PCR polymorphic band. *Mycobiology*, 30, 202-207.
- Kantety, R. V., La Rota, M., Matthews, D. E., & Sorrells, M. E. (2002). Data mining for simple-sequence repeats in expressed sequence tags from barley, maize, rice, sorghum, and wheat. *Plant Molecular Biology*, 48, 501-510.
- Kawai, T. (2004). Morphology, life history, and molecular phylogeny of *Stschapovia flagellaris* (Tilopteridales, Phaeophyceae) and the erection of the Stschapoviaceae fam. nov. *Journal of Phycology*, 40, 1156-1169.
- Kenrick, P., & Crane, P. R. (1997). The origin and early evolution of plants on land. *Nature*, 389, 33-39.
- Khlestkina, E. K., Myint Than, M. H., Pestsova, E. G., Röder, M. S., Malyshev, S. V., Korzun, V., & Börner, A. (2004). Mapping of 99 new microsatellite-derived loci in rye (*Secale cereale* L.) including 39 expressed sequence tags. *Theoretical and Applied Genetics*, 109, 725-732.

- Kim, K. M., Hoarau, G. G., & Boo, S. M. (2012). Genetic structure and distribution of *Gelidium elegans* (Gelidiales, Rhodophyta) in Korea based on mitochondrial *cox1* sequence data. *Aquatic Botany*, 98, 27-33.
- Kim, M. S., Kim, M., Terada, R., Yang, E. C., & Boo, S. M. (2008a). *Gracilaria parvispora* is the correct name of the species known as *G. bursa-pastoris* in Korea and Japan. *Taxon*, 57, 231-237.
- Kim, M. S., Yang, M. Y., & Cho, G. Y. (2010). Applying DNA barcoding to Korean Gracilariaceae (Rhodophyta). *Cryptogamie, Algologie*, 31, 387-401.
- Klinbunga, S., Ampayup, P., Tassanakajon, A., Jarayabhand, P., & Yoosukh, W. (2000). Development of species-specific markers of the tropical oyster (*Crassostrea belcheri*) in Thailand. *Marine Biotechnology*, 2, 476-484.
- Kolender, A. A., Matulewicz, M. C., & Cerezo, A. S. (1995). Structural analysis of antiviral sulfated alpha-D-(1-->3)-linked mannans. *Carbohydrate Research* 273, 179-185.
- Kostamo, K., Blomster, J., Korpelainen, H., Kelly, J., Maggs, C. A., & Mineur, F. (2008). New microsatellite markers for *Ulva Intestinalis* (Chlorophyta) and the transferability of markers across species of Ulvaceae. *Phycologia*, 47, 580-587.
- Kostrzewa, M., Valentin, K., Maid, U., Radetzky, R., & Zetsche, K. (1990). Structure of the rubisco operon from the multicellular red alga *Antithamnion spec.* *Current Genetics*, 18, 465-469.
- Kota, R., Varshney, R. K., Thiel, T., Dehmer, K. J., & Graner, A. (2001). Generation and comparison of EST-derived SSRs and SNPs in barley (*Hordeum vulgare* L.). *Hereditas*, 135, 145-151.
- Kress, W. J., & Erickson, D. L. (2007). A two-locus global DNA barcode for land plants: the coding *rbcL* gene complements the non-coding *trnH-psbA* spacer region. *PLoS ONE*, 2, 508.
- La Claire II, J. W., & Herrin, D. L. (1997). Co-isolation of highquality DNA and RNA from coenocytic green algae. *Plant Molecular Biology Reporter*, 15, 263-272.

- Landry, B. S. (1994). DNA mapping in plants. In B. R. Glick & J. E. Thompson (Eds.), *Methods in plant molecular biology and biotechnology* (pp. 269-285). Boca Raton, Florida: CRC Press.
- Leblanc, C., Boyen, C., Richard, O., Bonnard, G., Grienenberger, J. M., & Kloareg, B. (1995a). Complete sequence of the mitochondrial DNA of the rhodophyte *Chondrus crispus* (Gigartinales). Gene content and genome organization. *Journal of Molecular Biology*, 250, 484-495.
- Lee, R. E. (1999). *Phycology* (Third edition ed.): Cambridge University Press.
- Lehmann, T., Hawley, W. A., & Collins, F. H. (1996). An evaluation of evolutionary constraints on microsatellite loci using null alleles. *Genetics*, 144, 1155-1163.
- Levinson, G., & Gutman, G. A. (1987). Slipped-strand mispairing: a major mechanism for DNA sequence evolution. *Molecular Biology and Evolution*, 4, 203-221.
- Levring, T., Hoppe, H. A. & Schmid, O. J. (1969). *Marine Algae. A survey of research and utilization*. Cram, De Gruyter and Co., Hamburg, Germany.
- Lewis, L. A., & Flechtner, V. R. (2004). Cryptic species of *Scenedesmus* (Chlorophyta) from desert soil communities of Western North America. *Journal of Phycology*, 40, 1127-1137.
- Lewis, L. A., & McCourt, R. M. (2004). Green algae and the origin of land plants *American Journal of Botany*, 91, 1535-1556.
- Lewmanomont, K. (1994). The species of *Gracilaria* from Thailand. In I. A. Abbott (Ed.), *Taxonomy of economic seaweeds* (Vol. 4, pp. 135-148): La Jolla: California Sea Grant College Program.
- Liewlaksaneeyanawin, C., Ritland, C. E., El-Kassaby, Y. A., & Ritland, K. (2004). Single-copy, species-transferable microsatellite markers developed from loblolly pine ESTs. *Theoretical and Applied Genetics*, 109, 361-369.
- Lim, B. L., Kawai, H., Hori, H., & Osawa, S. (1986). Molecular evolution of 5S ribosomal RNA from red and brown algae. *Japanese Journal of Genetics*, 61, 169-176.



- Lim, P. E., & Phang, S. M. (2004). *Gracilaria* species (Gracilariales, Rhodophyta) of Malaysia including two new records. *Malaysian Journal of Science*, 23, 71-80.
- Lim, P. E., Thong, K. L., & Phang, S. M. (2001). Molecular differentiation of two morphological variants of *Gracilaria salicornia*. *Journal of Applied Phycology*, 13, 335-342.
- Lin, M. C., & Liao, C. M. (1999). 65Zn(II) accumulation in the soft tissue and shell of abalone *Haliotis diversicolor supertexta* via the alga *Gracilaria tenuistipitata* var. *liui* and the ambient water. *Aquaculture*, 178, 89-101.
- Lirasan, T., & Twide, P. (1993). Farming *Eucheuma* in Zanzibar, Tanzania. *Hydrobiologia*, 260-261, 353-355.
- Liu, F., Wang, X., Yao, J., Fu, W., & Duan, D. (2010). Development of expressed sequence tag-derived microsatellite markers for *Saccharina (Laminaria) japonica*. *Journal of Applied Phycology*, 22, 109-111.
- Liu, Z. J., & Cordes, J. F. (2004). DNA marker technologies and their applications in aquaculture genetics. *Aquaculture*, 238, 1-37.
- Lluisma, A. O., & Ragan, M. A. (1997). Expressed sequence tags (ESTs) from the marine red alga *Gracilaria gracilis*. *Journal of Applied Phycology*, 9, 287-293.
- Luxton, D. M. (1993). Aspects of the farming and processing of *Kappaphycus* and *Eucheuma* in Indonesia. *Hydrobiologia*, 260-261, 365-371.
- Lynch, M., & Walsh, B. (1998). *Genetics and analysis of quantitative traits*. Sunderland, MA: Sinauer Associates.
- Maggs, C. A., Verbruggen, H., & De Clerck, O. (2007). Molecular systematics of red algae: building future structures of firm foundations. In J. Brodie & J. Lewis (Eds.), *Unravelling the algae: the past, present and future of algal systematics* (pp. 103-121). Boca Raton, London & New York: CRC Press.
- Manhart, J. R., & McCourt, R. M. (1992). Molecular data and species concepts in the algae. *Journal of Phycology*, 28, 730-737.

- Marino, M. A., Turni, L. A., Del Rio, S. A., Williams, P. E., & Cregan, P. B. (1995). The analysis of simple sequence repeat DNA in soybean by capillary gel electrophoresis. *Applied and Theoretical Electrophoresis*, 5, 1-5.
- McHugh, D. J. (2003). A guide to the seaweed Industry. *FAO Fisheries Technical Paper 441*.
- Meneses, I., & Abbott, I. A. (1987). *Gracilaria* and *Polycavernosa* (Rhodophyta) from Micronesia. *Micronesica*, 20, 187-200.
- Meusnier, I., Valero, M., Olsen, J., & Stam, W. (2004). Analysis of rDNA ITS1 indels in *Caulerpa taxifolia* (Chlorophyta) supports a derived, incipient species status for the invasive strain. *European Journal of Phycology*, 39, 83-92.
- Moen, E., & Ostgaard, K. (1997). Aerobic digestion of Ca-alginate gels studied as a model system of seaweed tissue degradation. *Journal of Applied Phycology*, 9, 261-267.
- Morgante, M., Hanafey, M., & Powell, W. (2002). Microsatellites are preferentially present with non-repetitive DNA in plant genomes. *Nature Genetics*, 30, 194-200.
- Morgante, M., & Olivieri, A. M. (1993). PCR-amplified microsatellites as markers in plant genetics. *The Plant Journal*, 3, 175-182.
- Mosseler, A. (1992). Life history and genetic diversity in red pine: implications for gene conservation in forestry. *The Forestry Chronicle*, 68, 701-708.
- Mosseler, A., Innes, D. J., & Roberts, B. A. (1991). Lack of allozymic variation in disjunct Newfoundland populations of red pine (*Pinus resinosa*). *Canadian Journal of Forest Research*, 21, 525-528.
- Murata, M., & Nakazoe, J. (2001). Production and use of marine algae in Japan. *Japan Agricultural Research Quarterly*, 35, 281-290.
- Nam, K. W., Maggs, C. A., McIvor, L., & Stanhope, M. J. (2000). Taxonomy and phylogeny of *Osmundea* (Rhodophyceae, Rhodophyta) in Atlantic Europe. *Journal of Phycology*, 36, 759-772.

- Nguyen, H. D. (1992). Vietnamese species of *Gracilaria* and *Gracilariopsis*. In I. A. Abbott (Ed.), *Taxonomy of Economic Seaweeds* (Vol. 3, pp. 207-210): La Jolla, California: California Sea Grant College.
- Nicot, N., Chiquet, V., Gandon, B., Amilhat, L., Legeai, F., Leroy, P., . . . Sourdille, P. (2004). Study of simple sequence repeat (SSR) markers from wheat expressed sequence tags (ESTs). *Theoretical and Applied Genetics*, *109*, 800-805.
- Nisizawa, K., Noda, H., Kikuchi, R., & Watanabe, T. (1987). The main seaweeds foods in Japan. *Hydrobiologia*, *151/152*, 5-29.
- Ohno, M., & Largo, D. B. (1998). The seaweed resources of Japan. In M. Ohno & A. T. Critchley (Eds.), *Seaweed Resources of the World* (pp. 1-14): Kanagawa International Fisheries Training Center, Japan International Cooperative Agency.
- Olsen, J. L., Sadowski, G., Stam, W. T., Veldsink, J. H., & Jones, K. (2002). Characterization of microsatellite loci in the marine seaweed *Ascophyllum nodosum* (Phaeophyceae ; Fucales ). *Molecular Ecology Notes*, *2*, 33-34.
- Palmer, J. D. (1991). Plastid chromosomes: structure and evolution. In L. Bogorad & I. K. Vasil (Eds.), *Cell Culture and Somatic Genetics of Plant, Vol. 7A, Molecular Biology of Plastids* (pp. 5-53). San Diego: Academic Press.
- Pammi, S., Schertz, K., Xu, G., Hart, G., & Mullet, J. E. (1994). Random-amplified-polymorphic DNA markers in sorghum. *Theoretical and Applied Genetics*, *89*, 80-88.
- Pang, Q., Sui, Z., Kang, K. H., Kong, F., & Zhang, X. (2010). Application of SSR and AFLP to the analysis of genetic diversity in *Gracilariopsis lemaneiformis* (Rhodophyta). *Journal of Applied Phycology*, *22*, 607-612.
- Papenfuss, G. F. (1967). Notes on algal nomenclature - V. Various Chlorophyceae and Rhodophyceae. *Journal of Phycological Society*, *5*, 95-105.
- Parsons, T. J., Maggs, C. A., & Douglas, S. E. (1990). Plastid DNA restriction analysis links the heteromorphic phases of an apomictic red algal life history. *Journal of Phycology*, *26*, 495-500.

- Partis, L., & Wells, R. J. (1996). Identification of fish species using random amplified polymorphic DNA (RAPD). *Molecular and Cellular Probes*, *10*, 435-441.
- Pemberton, J. M., Slate, J., Bancroft, D. R., & Barrett, J. A. (1995). Nonamplifying alleles at microsatellite loci – a caution for parentage and population studies. *Molecular Ecology*, *4*, 249-252.
- Pepin, L., Amigues, Y., Lepingle, A., Berthier, J., Bensaid, A., & Vaiman, D. (1995). Sequence conservation of microsatellites between *Bos Taurus* (cattle), *Capra hircus* (goat) and related species. Examples of use in parentage testing and phylogeny analysis. *Heredity*, *74*, 53-61.
- Perez de la Vega, M. P. (1993). Biochemical characterization of populations. In M. D. Hayward, N. O. Bosermark & I. Romagosa (Eds.), *Plant breeding: principles and prospects* (pp. 182-201). London: Chapman and Hall.
- Petes, T. D., Greenwell, P. W., & Dominska, M. (1997). Stabilization of microsatellite sequences by variant repeats in the yeast *Saccharomyces cerevisiae*. *Genetics*, *146*, 491-498.
- Phang, S. M. (2006). Seaweed resources in Malaysia: current status and future prospects. *Aquatic Ecosystem Health & Management*, *9*, 185-202.
- Phang, S. M., & Lewmanomont, K. (2001). *Gracilaria changii* (B. M. Xia & I. A. Abbott) I. A. Abbott, C. F. Chang & B. M. Xia. In P. Reine & G. C. J. Trono (Eds.), *Plant Resources of South-East Asia. No. 15(1) Cryptogams: Algae*. (pp. 178-180). Leiden: Backhuys Publishers.
- Phang, S. M., Shahrudin, S., Noraishah, H., & Sasekumar, A. (1996). Studies on *Gracilaria changii* (Gracilariales, Rhodophyta) from Malaysian mangroves. *Hydrobiologia*, *326-327*, 347-352.
- Phang, S. M., & Yoshida, T. (1997). *Sargassum stolonifolium* Phang et Yoshida sp. nov. from Penang Island, Peninsular Malaysia. In I. A. Abbott (Ed.), *Taxonomy of Economic Seaweeds: with reference to some Pacific and Caribbean species. Vol. VI* (pp. 61-73). California: California Sea Grant College.
- Phillips, N., Smith, C. M., & Morden, C. W. (2001). An effective DNA extraction protocol for brown algae. *Phycological Research*, *49*, 97-102.

- Picoult-Newberg, L., Ideker, T. E., Pohl, M. G., Taylor, S. L., Donaldson, M. A., Nickerson, D. A., & Boyce-Jacino, M. (1999). Mining SNPs from EST databases. *Genome Research*, *9*, 167-174.
- Popovich, D. G., Yeo, C. R., & Zhang, W. (2012). Ginsenosides derived from Asian (*Panax ginseng*), American ginseng (*Panax quinquefolius*) and potential cytoactivity. *International Journal of Biomedical and Pharmaceutical Sciences*. 203.183.32.151.
- Potter, R. H., & Jones, M. G. K. (1991). Molecular analysis of genetic stability. In J. H. Dodds (Ed.), *In vitro methods for conservation of plant genetic resources* (pp. 71-91). London: Chapman and Hall.
- Powell, W., Machray, G. C., & Provan, J. (1996). Polymorphism revealed by simple sequence repeats. *Trends in Plant Science*, *1*, 215-222.
- Powell, W., Morgante, M., Andre, C., McNicol, J., Machray, G., Doyle, J., . . . Rafalski, J. (1995a). Hypervariable microsatellites provide a general source of polymorphic DNA markers for the Chloroplast genome. *Current Biology* *5*, 1023-1029.
- Powell, W., Morgante, M., McDevitt, R., Vendramin, G., & Rafalski, J. (1995b). Polymorphic simple sequence repeat regions in chloroplast genomes: applications to the population genetics of pines. *Proceedings of the National Academy of Sciences, USA*, *92*, 7759-7763.
- Provan, J., Kumar, A., Shepherd, L., Powell, W., & Waugh, R. (1996). Analysis of intra-specific somatic hybrids of potato (*Solanum tuberosum*) using simple sequence repeats. *Plant Cell Reports*, *16*, 196-199.
- Provan, J., & Maggs, C. A. (2012). Unique genetic variation at a species's rear edge is under threat from global climate change. *Proceedings of the Royal Society of London Series*, *279*, 39-47.
- Provan, J., Biss, P. M., McMeel, D., & Mathews, S. (2004a). Universal primers for the amplification of chloroplast microsatellites in grasses (*Poaceae*). *Molecular Ecology Notes*, *4*, 262-264.
- Provan, J., Murphy, S., & Maggs, C. (2004b). Universal plastid primers for Chlorophyta and Rhodophyta. *European Journal of Phycology*, *39*, 43-50.

- Provan, J., Powell, W., & Hollingsworth, P. (2001). Chloroplast microsatellites: new tools for studies in plant ecology and evolution. *Trends in Ecology & Evolution*, *16*, 142-147.
- Qin, S., Jiang, P., & Tseng, C. K. (2004). Molecular biotechnology of marine algae in China. *Hydrobiologia*, *512*, 21-26.
- Qureshi, S. N., Saha, S., Kantety, R. V., & Jenkins, J. N. (2004). EST-SSR: A new class of genetic markers in cotton. *Journal of Cotton Science*, *8*, 112-123.
- Rabouam, C., Comes, A. M., Bretagnolle, V. V., Humbert, J. F., Periquet, G., & Bigot, Y. (1999). Features of DNA fragments obtained by random amplified polymorphic DNA (RAPD) assays. *Molecular Ecology*, *8*, 493-503.
- Rafalski, A. (2002). Applications of single nucleotide polymorphism in crop genetics. *Current Opinion in Plant Biology*, *5*, 94-100.
- Ramsay, L., Macaulay, M., Degli Ivanisovich, S., MacLean, K., Cardle, L., Fuller, J., . . . Waugh, R. (2000). A Simple Sequence Repeat-Based Linkage Map of Barley. *Genetics*, *156*, 1997-2005.
- Rassmann, K., Schlötterer, C., & Tautz, D. (1991). Isolation of simple-sequence loci for use in polymerase chain reaction-based DNA fingerprinting. *Electrophoresis*, *12*, 113-118.
- Reith, M., & Munholland, J. (1995). Complete nucleotide sequence of the *Porphyra purpurea* chloroplast genome. *Plant Molecular Biology Reporter*, *13*, 333-335.
- Renn, D. (1997). Biotechnology and the red seaweed polysaccharide industry: status, needs and prospects. *Trends in Biotechnology*, *15*, 9-14.
- Ritland, C., & Ritland, K. (2000). DNA-fragment markers in plants. In A. J. Baker (Ed.), *Molecular methods in ecology* (pp. 208-234). Oxford, UK: Blackwell Science Ltd.
- Robba, L., Russell, S. J., Barker, G. L., & Brodie, J. (2006). Assessing the use of mitochondrial *cox1* marker for use in DNA barcoding of red algae (Rhodophyta). *American Journal of Botany*, *93*, 1101-1108.

- Rongwen, J., Akkaya, M. S., Bhagwat, A. A., Lavi, U., & Cregan, P. B. (1995). The use of microsatellite DNA markers for soybean genotype identification. *Theoretical and Applied Genetics*, 90, 43-48.
- Routtu, J., Hoikkala, A., & Kankare, M. (2007). Microsatellite-based species identification method for *Drosophila virilis* group species. *Hereditas*, 144, 213-221.
- Rozen, S., & Skaletsky, H. J. (2000). *Primer3 on the WWW for general users and for biologist programmers*. Paper presented at the Bioinformatics Methods and Protocols: Methods in Molecular Biology, Humana Press, Totowa, NJ.
- Rudd, S. (2003). Expressed sequence tags: Alternative or complement to whole genome sequences? *Trends in Plant Science*, 8, 321-329.
- Saha, S., Karaca, M., Jenkins, J. N., Zipf, A. E., Reddy, O. U. K., & Kantety, R. V. (2003). Simple sequence repeats as useful resources to study transcribed genes of cotton. *Euphytica*, 130, 355-364.
- Saiki, R. K., Gelfand, D. H., Stoffel, S., Scharf, S. J., Higuchi, R., Horn, G. T., . . . Erlich, H. A. (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science*, 239, 487-491.
- Sánchez-Pérez, R., Ballesterb, J., Dicentaa, F., Arúsb, P., & Martínez-Gómez, P. (2006). Comparison of SSR polymorphisms using automated capillary sequencers, and polyacrylamide and agarose gel electrophoresis: Implications for the assessment of genetic diversity and relatedness in almond. *Scientia Horticulturae*, 108, 310-316.
- Santos, G. A., & Doty, M. S. (1983). Agarose from *Gracilaria cylindrica*. *Botanica Marina*, 26, 31-34.
- Saunders, G. W. (2005). Applying DNA barcoding to red macroalgae: a preliminary appraisal holds promise for future applications. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360, 1879-1888.
- Saunders, G. W. (2009). Routine DNA barcoding of Canadian Gracilariales (Rhodophyta) reveals the invasive species *Gracilaria vermiculophylla* in British Columbia. *Molecular Ecology Resources*, 9, 140-150.

- Schierwater, B., Metzler, D., Krüger, K., & Streit, B. (1996). The effects of nested primer binding sites on the reproducibility of PCR: mathematical modeling and computer simulation studies. *Journal of Computational Biology*, 3, 235-251.
- Schlötterer, C. (1998). Microsatellites. In A. R. Hoelzel (Ed.), *Molecular genetic analysis of populations: A practical approach* (pp. 237-261). Oxford: IRL Press.
- Schlötterer, C., & Pemberton, J. (1994). The use of microsatellites for genetic analysis of natural populations. *EXS*, 69, 203-214.
- Schuelke, M. (2000). An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology*, 18, 233-234.
- Schug, M., Hutter, C., Wetterstrand, K., Gaudette, M., Mackay, T., & Aquadro, C. (1998). The mutation rates of di-, tri- and tetranucleotide repeats in *Drosophila melanogaster*. *Molecular Biology and Evolution*, 15, 1751-1760.
- Scott, K. D. (2001). Microsatellites derived from ESTs and their comparison with those derived from by other methods. In R. J. Henry (Ed.), *Plant genotyping: the DNA fingerprinting of plants* (pp. 225-237). New York: CABI.
- Scribner, K. T., & Pearce, J. M. (2000). Microsatellites: Evolutionary and methodological background and empirical applications at individual, population, and phylogenetic levels. In A. Baker (Ed.), *Molecular Methods in Ecology*. Oxford, UK: Blackwell Science, Ltd.
- Selkoe, K. A., & Toonen, R. J. (2006). Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. *Ecology Letters*, 9, 615-629.
- Semagn, K., Bjørnstad, Å., & Ndjiondjop, M. N. (2006). An overview of molecular marker methods for plants. *African Journal of Biotechnology*, 5, 2540-2568.
- Sharma, A., Namdeo, G., & Mahadik, K. R. (2008). Molecular markers: New prospects in plant genome analysis. *Pharmacognosy Reviews*, 2, 23-24.
- Shi, Y., Yang, G., Liu, Y., Liao, M., Li, X., & Cong, Y. (2007). Development of 18 polymorphic microsatellite DNA markers of *Laminaria japonica* (Phaeophyceae). *Molecular Ecology Notes*, 7, 620-622.



- Shivji, M. S. (1991). Organization of the chloroplast genome in the red alga *Porphyra yezoensis*. *Current Genetics*, *19*, 49-54.
- Sim, M. C., Lim, P. E., Gan, S. Y., & Phang, S. M. (2007). Identification of random amplified polymorphic DNA (RAPD) marker for differentiating male from female and sporophytic thalli of *Gracilaria changii* (Rhodophyta). *Journal of Applied Phycology*, *19*, 763-769.
- Smartfiber (2010). SeaCell® pure—SeaCell® active.  
([http://www.smartfiber.de/index.php?option=com\\_content&view=article&id=49&Itemid=430&lang=en](http://www.smartfiber.de/index.php?option=com_content&view=article&id=49&Itemid=430&lang=en))
- Smit, A. J. (2004). Medicinal and pharmaceutical uses of seaweed natural products: A review. *Journal of Applied Phycology*, *16*, 245-262.
- St ägel, A., Portis, E., Toppino, L., Rotino, G. L., & Lanteri, S. (2008). Gene-based microsatellite development for mapping and phylogeny studies in eggplant. *BMC Genomics*, *9*, 357.
- Stephenson, P., Bryan, G., Kirby, J., Collins, A., Devos, K., Busso, C., & Gale, M. (1998). Fifty new microsatellite loci for the wheat genetic map. *Theoretical and Applied Genetics*, *97*, 946-949.
- Sun, J., Liu, T., Guo, B., Jin, D., Weng, M., Feng, Y., . . . Wang, B. (2006). Development of SSR primers from EST sequences and their application in germplasm identification of *Porphyra* lines (Rhodophyta). *European Journal of Phycology*, *41*, 329-336.
- Sutherland, G. R., & Richards, R. I. (1995). Simple tandem DNA repeats and human genetic disease. *Proceedings of the National Academy of Sciences, U. S. A.*, *92*, 3636-3641.
- Sze, P. (1998). *A biology of the algae*: WCB/McGraw-Hill.
- Takahashi, M. M. (2010). Sequênciamento e anáise do genoma mitocondrial de *Gracilaria tenuistipitata* (Gracilariales, Rhodophyta). Master Thesis, Institute of Biosciences, University of São Paulo.  
(<http://www.teses.usp.br/teses/disponiveis/41/41132/tde-06102010-164456/>)

- Temnykh, S., DeClerck, G., Lukashova, A., Lipovich, L., Cartinhour, S., & McCouch, S. (2001). Computational and Experimental Analysis of Microsatellites in Rice (*Oryza sativa* L.): Frequency, Length Variation, Transposon Associations, and Genetic Marker Potential. *Genome Research*, *11*, 1441-1452.
- Temnykh, S., Park, W. D., Ayres, N., Cartinhour, S., Hauck, N., Lipovich, L., . . . McCouch, S. R. (2000). Mapping and genome organization of microsatellite sequences in rice (*Oryza sativa* L.) *Theoretical and Applied Genetics*, *100*, 697-712.
- Tanaka, H., Fukuda, N., & Shoyama, Y. (2006). Identification and differentiation of *Panax* species using ELISA, RAPD and eastern blotting. *Phytochemical Analysis*, *17*, 46-55.
- Tassanakajon, A., Pongsomboon, S., Jarayabhand, P., Klinbunga, S., & Boonsaeng, V. V. (1998). Genetic structure in wild populations of black tiger shrimp (*Penaeus monodon*) using randomly amplified polymorphic DNA analysis. *Journal of Marine Biotechnology*, *6*, 249-254.
- Terachi, T. (1993). The progress of DNA analyzing techniques and its impact on plant molecular systematics. *Journal of Plant Research*, *106*, 75-79.
- Terada, R., Kawaguchi, S., Masuda, M., & Phang, S. M. (2000). Taxonomic Notes on Marine Algae from Malaysia. III. Seven Species of Rhodophyceae. *Botanica Marina*, *43*, 347-357.
- Therkelsen, G. H. (1993). Carrageenan. In R. L. Whistler & J. N. BeMiller (Eds.), *Industrial gums* (pp. 156-161). New York: NY: Academic Press.
- Thiel, T., Michalek, W., Varshney, R. K., & Graner, A. (2003). Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics*, *106*, 411-422.
- Thomas, M. R., & Scott, N. S. (1993). Microsatellite repeats in grapevine reveal DNA polymorphisms when analyzed as sequence-tagged sites (STSs). *Theoretical and Applied Genetics*, *86*, 985-990.

- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F., & Higgins, D. G. (1997). The Clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*, *24*, 4876-4882.
- Tingey, S. V., Rafalski, J. A., & Williams, J. G. K. (1993). *Application of RAPD technology to plant breeding*. Minnesota: ASHS Publishers.
- Towle, G. A., & Whistler, R. L. (1973). Industrial Gums: Polysaccharides and Their Derivatives. In L. P. Miller (Ed.), *Phytochemistry: Inorganic elements and special groups of chemicals* (pp. 198-248). New York: Van Nostrand Reinhold Co.
- Triboush, S. O., Danilenko, N. G., & Davydenko, O. G. (1998). A method for isolation of chloroplast DNA and mitochondrial DNA from sunflower. *Plant Molecular Biology Reporter*, *16*, 183-189.
- Tseng, C. K., & Xia, B. M. (1999). On the *Gracilaria* in the western Pacific and southeastern Asia region. *Botanica Marina*, *42*, 209-217.
- Tsutsui, I., Huynh, Q. N., Nguyễn, H. D., Arai, S., & Yoshida, T. (2005). The common marine plants of southern Vietnam. *numerous colour photographs. Usa: Japan Seaweed Association*, 1-250 + [251].
- van den Hoek, C., Mann, D. G., & Jahns, H. M. (1995). *Algae: An Introduction to Phycology*: Cambridge University Press, Cambridge.
- van der Strate, H. J., Olsen, J. L., van de Zande, L., Edwards, K. J., & Stam, W. T. (2000). Isolation and characterization of microsatellite loci in the benthic seaweed, *Cladophoropsis membranacea* (Cladophorales, Chlorophyta). *Molecular Ecology*, *9*, 1442-1443.
- Varela-Alvarez, E., Andreakis, N., Lago-Leston, A., Pearson, G. A., Serrao, E. A., Procaccini, G., . . . Marba, N. (2006). Genomic DNA isolation from green and brown algae (Caulerpales and Fucales) for microsatellite library construction. *Journal of Phycology*, *42*(3), 741-745.
- Varshney, R. K., Beier, U., Khlestkina, E., Kota, R., Korzun, V., Röder, M., . . . Börner, A. (2007). Single nucleotide polymorphisms in rye: discovery, frequency and applications for genome mapping and diversity studies. *Theoretical and Applied Genetics*, *114*, 1105-1116.

- Varshney, R. K., Graner, A., & Sorrells, M. E. (2005). Genic microsatellite markers in plants: features and applications. *Trends in Biotechnology*, 23, 48-55.
- Varshney, R. K., Thiel, T., Stein, N., Langridge, P., & Graner, A. (2002). In silico analysis on frequency and distribution of microsatellites in ESTs of some cereal species. *Cellular and Molecular Biology Letters*, 7, 537-546.
- Vejl, P. (1997). Identification of genotypes in hop (*Humulus lupulus*) by RAPD analysis using Gel Manager for Windows. *Rostlinna Vyroba* 43, 325-331.
- Vendramin, G. G., Lelli, L., Rossi, P., & Morgante, M. (1996). A set of primers for the amplification of 20 chloroplast microsatellites in *Pinaceae*. *Molecular Ecology*, 5, 595-598.
- Vidal, R., Meneses, I., & Smith, M. (2008). Phylogeography of the genus *Spongites* (corallinales, rhodophyta) from Chile. *Journal of Phycology*, 44, 173-182.
- Voisin, M., Engel, C. R., & Viard, F. (2005). Differential shuffling of native genetic diversity across introduced regions in a brown alga: aquaculture vs. maritime traffic effects. *Proceedings of National Academy of Sciences of the United States of America*, 102, 5432-5437.
- Vos, P., Hogers, R., Bleeker, M., Reijans, M., Van der Lee, T., Hornes, M., . . . Zabeau, M. (1995). AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research*, 23, 4407-4414.
- Walker, A. B., Fournier, H. R., Neefus, C. D., Nardi, G. C., & Berlinsky, D. L. (2009). Partial replacement of fish meal with laver *Porphyra* spp. in diets for atlantic cod. *North American Journal of Aquaculture* 71, 39-45.
- Wang, D., Wang, X., Li, D., Wang, F., & Duan, D. (2006). The genetic analysis and germplasm identification of the gametophytes of *Undaria pinnatifida* (Phaeophyceae) with RAPD method. *Journal of Applied Phycology*, 18, 801-809.
- Wang, M. L., Mosjidis, J. A., Morris, J. B., Chen, Z. B., Barkley, N. A., & Pederson, G. A. (2009). Evaluation of *Lespedeza* germplasm genetic diversity and its phylogenetic relationship with the genus *Kummerowia*. *Conservation Genetics*, 10, 79-85.

- Wang, Z., Weber, J. L., Zhong, G., & Tanksley, S. D. (1994). Survey of plant short tandem DNA repeats. *Theoretical and Applied Genetics*, 88, 1-6.
- Wattier, R., Dallas, J. F., Destombe, C., Saumitou-Laprade, P., & Valero, M. (1997). Single locus microsatellites in Gracilariales (Rhodophyta): High level of genetic variability within *Gracilaria gracilis* and conservation in related species. *Journal of Phycology*, 33, 868-880.
- Wattier, R., & Maggs, C. A. (2001). Intraspecific variation in seaweeds: the application of new tools and approaches. *Advances in Botanical Research*, 35, 171-212.
- Wattier, R. A., Prod'ohl, P. A., & Maggs, C. A. (2000). DNA isolation protocol for red seaweed (Rhodophyta). *Plant Molecular Biology Reporter*, 18, 275-281.
- Weber, D., & Helentjaris, T. (1989). Mapping RFLP loci in maize using B-a translocations. *Genetics*, 121, 583-590.
- Weber, J. L. (1990). Informativeness of human (dC-dA)<sub>n</sub>(dG-dT)<sub>n</sub> polymorphisms. *Genomics*, 7, 524-530.
- Weller, J. W., & Reddy, A. (1997). Fluorescent detection and analysis of RAPD amplicons using the ABI PRISM DNA sequencers. In M. R. Micheli & R. Bova (Eds.), *Fingerprinting methods based on arbitrarily primed PCR* (pp. 81-92): Springer Lab Manual.
- Welsh, J., & McClelland, M. (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Research*, 18, 7213-7218.
- Whitmer, A. C. (2002). Microsatellite markers for the intertidal kelp *Postelsia palmaeformis* (Heterokontophyta; Laminariales). *Molecular Ecology Notes*, 2, 469-471.
- Williams, J. G., Kubelik, A. R., Livak, K. J., Rafalski, J. A., & Tingey, S. V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research*, 18, 6531-6535.
- Winston, J. E. (1999). Describing species – practical taxonomic procedure for biologists. Columbia University Press, New York.

- Withell, A. F., Millar, A. J. K., & Kraft, G. T. (1994). Taxonomic studies of the genus *Gracilaria* (Gracilariales, Rhodophyta) from Australia. *Australian Systematic Botany*, 7, 281-352.
- Witvrouw, M., Este, J., Mateu, M., Reymen, D., Andrei, G., Snoeck, R., . . . De Clercq, E. (1994). Activity of a sulfated polysaccharide extracted from the red seaweed *Aghardhiella tenera* against human immunodeficiency virus and other enveloped viruses. *Antiviral Chemistry and Chemotherapy*, 5, 297-303.
- Wynne, M. J. (1989). The reinstatement of *Hydropuntia* Montagne (Gracilariaceae, Rhodophyta). *Taxon*, 38, 476-479.
- Wynne, M. J. (1995). Benthic marine algae from the Seychelles collected during the R/V *Te Vega* Indian Ocean Expedition. *Contr. Univ. Michigan Herb*, 20, 261-346.
- Xia, B. M. (1986). On *Gracilaria salicornia* (C. Agardh) Dawson. *Chinese Journal of Oceanology and Limnology*, 4, 100-107.
- Xia, B. M., & Abbott, I. A. (1987). New species of Polycavernosa Chang & Xia (Gracilariaceae, Rhodophyta) from the western Pacific. *Phycologia*, 26, 405-418.
- Xie, C. T., Chen, C. S., Ji, D. H., & Xu, Y. (2009). Characterization, development and exploitation of EST-derived microsatellites in *Porphyra haitanensis* Chang et Zheng (Bangiales, Rhodophyta). *Journal of Applied Phycology*, 21, 367-374.
- Yamamoto, H. (1975). The relationship between *Gracilariopsis* and *Gracilaria* from Japan. *Bulletin of the Faculty of Fisheries Hokkaido University*, 26, 217-222.
- Yamamoto, H. (1978). Systematic and anatomical study of the genus *Gracilaria* in Japan. *Memoirs of the Faculty of Fisheries, Hokkaido University*, 25, 97-152.
- Yamamoto, H., & Trono, G. C., Jr (1994). *Two new species of Gracilaria from the Philippines*. Paper presented at the Taxonomy of Economic Seaweeds with reference to some Pacific species, La Jolla, California: California Sea Grant College.

- Yang, E. C., & Boo, S. M. (2004). Evidence for two independent lineages of *Griffithsia* (Ceramiaceae, Rhodophyta) based on plastid protein-coding *psaA*, *psbA*, and *rbcL* gene sequences. *Molecular Phylogenetics and Evolution*, *31*, 680-688.
- Yang, E. C., Kim, M. S., Geraldino, P. J. L., Sahoo, D., Shin, J. A., & Boo, S. M. (2007). Mitochondrial *cox1* and plastid *rbcL* genes of *Gracilaria vermiculophylla* (Gracilariaceae, Rhodophyta). *Journal of Applied Phycology*, *20*, 161-168.
- Yang, G. P., Saghai Maroof, M. A., Xu, C. G., Zhang, Q., & Biyashev, R. M. (1994). Comparative analysis of microsatellite DNA polymorphism in landraces and cultivars of rice. *Molecular and General Genetics* *245*, 187-194.
- Yoshida, T., Stiger, V., & Horiguchi, T. (2000). *Sargassum boreale* sp. nov. (Fucales, Phaeophyceae) from Hokkaido, Japan. *Phycologia Research*, *48*, 125-131.
- Yow, Y. Y., Lim, P. E., & Phang, S. M. (2011). Genetic diversity of *Gracilaria changii* (Gracilariaceae, Rhodophyta) from west coast, Peninsular Malaysia based on mitochondrial *cox1* gene analysis. *Journal of Applied Phycology*, *23*, 219-226.
- Yow Y.Y., Lim P.E., & Phang, S.M. (2012). Assessing the use of mitochondrial *cox1* gene and *cox2-3* spacer for genetic diversity study of Malaysian *Gracilaria changii* (Gracilariaceae, Rhodophyta) from Peninsular Malaysia. *Journal of Applied Phycology*.
- Yu, J.-K., Dake, T. M., Singh, S., Benscher, D., Li, W., Gill, B., & Sorrells, M. E. (2004). Development and mapping of EST-derived simple sequence repeat markers for hexaploid wheat. *Genome*, *47*, 805-818.
- Zane, L., Bargelloni, L., & Patarnello, T. (2002). Strategies for microsatellite isolation: a review. *Molecular Ecology*, *11*, 1-16.
- Zhang, T., Fang, Y., Wang, X., Deng, X., Zhang, X., Hu, S., & Yu, J. (2012). The Complete Chloroplast and Mitochondrial Genome Sequences of *Boea hygrometrica*: Insights into the Evolution of Plant Organellar Genomes. *PLoS ONE*, *7*, 1.
- Zhou, Y. H., & Ragan, M. A. (1995). Cloning and characterization of the nuclear gene and cDNAs for triosphosphate isomerase of the marine red algae *Gracilaria verrucosa*. *Current Genetics*, *28*, 317-323.

- Zuccarello, G. C., Burger, G., West, J. A., & King, R. J. (1999). A mitochondrial marker for red algal intraspecific relationships. *Molecular Ecology Notes*, 8, 1443-1447.
- Zuccarello, G. C., Critchley, A. T., Smith, J., Sieber, V., Lhonneur, G. B., & West, J. A. (2006). Systematics and genetic variation in commercial *Kappaphycus* and *Euचेuma* (Solieriaceae, Rhodophyta). *Journal of Applied Phycology*, 1, 417-425.
- Zuccarello, G. C., & West, J. A. (2003). Multiple cryptic species: molecular diversity and reproductive isolation in the *Bostrychia radicans*/*B. moritziana* complex (Rhodomelaceae, Rhodophyta) with focus on North American isolates. *Journal of Phycology*, 39, 948-959.