
REFERENCES

- Apocada, C. K., & Chapman, L. J. (2004). Adult damselflies of Kibale National Park, Uganda: Habitat associations and seasonal occurrence. *Odonatologica*, 33, 129-146.
- Artiss, T., Schultz, T. R., Polhemus, D. A., & Simon, C. (2001). Molecular phylogenetic analysis of the dragonfly genera *Libellula*, *Ladona*, and *Plathemis* (Odonata: Libellulidae) based on mitochondrial *cytochrome oxidase I* and 16S rRNA sequence data. *Molecular Phylogenetics and Evolution*, 18, 348-361.
- Asahina, S. (1993). *A list of the Odonata from Thailand (Parts I – XXI)* (Bro. A. Pinratana, ed.), Bangkok.
- Askew, R. R. (1998). *The dragonflies of Europe*. Harley books, Colchester, 291 pp.
- Avise, J.C. (2004). *Molecular markers, natural history, and evolution*. (2nd ed). Sunderland : Sinauer Associates. Inc.
- Bechly, G. (1996). Morphologische untersuchungen am flugelgeader der rezenten Libellen und deren stammgruppenvertreter (Insecta: Odonatoptera), unter besonderer berucksichtigung der phylogenetischen systematic und des grundplanes der Odonata. *Petalura, spec*, 2, 402.
- Bechly, G., Nel, A., Martinez-Delclos, X., & Fleck, G. (1998). Four new dragonflies from the upper Jurassic of Germany and the lower Cretaceous of Mongolia (Anisoptera: Hemeroscopidae, Sonidae, and Proterogomphidae fam. nov.). *Odonatologica*, 27(2), 149 – 187.

-
- Bechly, G. (2002). The phylogenetic systematics of Odonata. Retrieved from <http://www.bechly.de/phylosys.htm>.
- Bely, A. E., & Weisblat, D. A. (2006). Lessons from leeches: a call for DNA barcoding in the lab. *Evolution and Development*, 8, 491-501.
- Borror, D. J., & White, R. E. (1970). A field guide to insects America north of Mexico. Boston, MA: Houghton Mifflin Company.
- Bridges, C. A. (1993). *Catalogue of the family-group, genus-group, and species group names of the Odonata of the world*. (2nd ed.) Urbana, IL.: Charles A. Bridges.
- Bridges, C. A. (1994). In: C. A. Bridges (Ed.), *Catalogue of the Family-Group, Genus-Group, and Species-Group Names of the Odonata of the World*, third ed., Urbana, IL, USA.
- Brown, J. M., McPeck, M. A., & May, M. L. (2000). A phylogenetic perspective on habitat shifts and diversity in the North American *Enallagma* damselflies. *Systematic Biology*, 49, 697 – 712.
- Bull, J. J., Huelsenbeck, J. P., Cunningham, C. W., Swofford, D. L., & Waddell, P. J. (1993). Partitioning and combining data in phylogenetic analysis. *Systematic Biology*, 42, 384 – 397.
- Burmeister, H. (1839). *Handbuch der Entomologie*. 2(2), 2. Berlin: Enslin.
- Carle, F. L. (1979). Environmental monitoring potential of the Odonata, with a list of rare and endangered Anisoptera of Virginia, USA. *Odonatologica*, 8, 319 – 323.

-
- Carle, F. L. (1982). The wing vein homologies and phylogeny of the Odonata: a continuing debate. *Soc. Int. Odonatol. (Rapid Commun.)*, 4, x-66.
- Carle, F. L. (1995). Evolution, taxonomy, and biogeography of ancient gondwanian libelluloides, with comments on anisopteroid evolution and phylogenetic systematics (Anisoptera: Libelluloidea). *Odonatologica*, 24(4), 383 – 506.
- Castella, E. (1987). Larval Odonata distribution as a descriptor of fluvial ecosystems: the Rhône and Ain Rivers, France. *Advances in Odonatology*, 3, 23 – 40.
- Caterino, S., Cho, S., & Sperling, F. A. H. (2000). The current state of insect molecular systematic: a thriving tower of Babel. *Annual Review of Entomology*, 45, 1-54.
- Chandra, S. B. C., Vlk, J. L., & Kapatral, V. (2006). Comparative insect mitochondrial genomes: Differences despite conserved genome synteny. *African Journal of Biotechnology*, 5(14), 1308 – 1318.
- Che Salmah, M. R. (1996). Some aspects of biology and ecology of *Neurothemis tullia* (Drury) (Odonata: Libellulidae) in the laboratory and rain fed rice field in Peninsular Malaysia. Ph.D. Thesis, Universiti Pertanian Malaysia, Serdang, Selangor, Malaysia.
- Chippindale, P. T., & Weins, J. J. (1994). Weighting, partitioning, and combining characters in phylogenetic analysis. *Systematic Biology*, 43, 278 – 287.
- Chippindale, P. T., Davén, V. K., Whitmore, D. H., & Robinson, J. V. (1999). Phylogenetic relationships of north American damselflies of the genus *Ischura* (Odonata: Zygoptera: Coenagrionidae) based on sequences of three mitochondrial genes. *Molecular Phylogenetics and Evolution*, 11, 110 – 121.

-
- Chovanec, A., & Raab, R. (1997). Dragonflies (Insecta, Odonata) and the ecological status of newly created wetlands – examples for long term bioindication programmes. *Limnology*, 27, 381 – 392.
- Clark, T. E., Samways, M. J. (1996). Dragonflies (Odonata) as indicators of biotope quality in the Krüger National Park, South Africa. *Journal of Applied Ecology*, 33, 1001–1012.
- Cook, L. M. (2008). Diversity and evenness from sequential sightings. *Insect Conservation and Diversity*. 1– 3.
- Corbet, P. S. (1980). Biology of Odonata. *Annual Reviews Entomology*, 25, 189- 717.
- Corbet, P. S. (1993). Are Odonata useful as bioindicators? *Libellula*, 12, 91–102.
- Corbet, P. S. (1999). Dragonflies: Behaviour and ecology of Odonata. Colchester, UK: Harley Books. 829 p.
- Crozier, R. H., & Crozier, Y. C. (1993). The mitochondrial genome of the honeybee *Apis mellifera*: complete sequence and genome organization, *Genetics*, 133, 97-117.
- Davies, D. A. L., & Tobin, P. (1985). The Dragonflies of the World: a Systematic List of the Extant Species of Odonata, Vol. 2, Anisoptera. *Societas Internationalis Odonatologica, Rapid Communications, Suppl. 5*, 151.
- de Quieroz, A., Donoghue, M. J., & Kim, J. Y. (1995). Separate versus combined analysis of phylogenetic evidence. *Annual Review of Ecology and Systematics*, 26, 657 – 681.

-
- Dietrich, C. H., Whitcomb, R. F., & Black, W. C. IV (1997). Phylogeny of the grassland leafhopper genus *Flexamia* (Homoptera: Cicadellidae) based on mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution*, 8, 139 – 149.
- Dijkstra, K. –D. B., Groeneveld, L. F., Clausnitzer, V., & Hadrys, H. (2007). The *Pseudagrion* split: Molecular phylogeny confirms the morphological and ecological dichotomy of Africa's most diverse genus of Odonata (Caenagrionidae). *International Journal of Odonatology*, 10, 31-41.
- Eck, R. V., & Dayhoff, M. O. (1966). *Atlas of protein sequence and structure*. Maryland : National Biomedical Research Foundation.
- Eyre, M. D., Ball, S. G., & Foster, G. N. (1986). An initial classification of the habits of aquatic Coleoptera in North-east England. *Journal Applied Ecology*, 23, 841 – 852.
- Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39, 783-791.
- Fincke, O. (1992). Behavioural ecology of the giant damselflies of Barro Colorado Island, Panama (Odonata: Zygoptera: Pseudostigmatidae). In D. Quintero, & A. Aoello. (Eds.) *Insects of Panama and Mesoamerica* (pp. 103–113). Oxford: Oxford University Press.
- Fleck, G., Brenk, M., & Misof, B. (2006). Larval and molecular characters help to solve phylogenetic puzzles in the highly diverse dragonfly family Libellulidae (Insecta: Odonata: Anisoptera): The Tetrathemistinae are a polyphyletic group. *Organisms Diversity Evolution*, 8, 1 – 16.

-
- Flook, P. K., Klee, S., & Rowell, C. H. F. (1999). Combined molecular phylogenetic analysis of the Orthoptera (Arthropoda, Insecta) and implications for their higher systematics *Systematic Biology*, *48*, 233 – 253.
- Fraser, F. C. (1957). *A reclassification of the order Odonata*. Sydney, Australia: Royal Zoological Society of New South Wales.
- Fulan, J. A., & Henry, R. (2006). The Odonata (Insecta) assemblage on *Eichhornia azurea* (Sw.) Kunth (Pontederiaceae) stands in Camargo Lake, a lateral lake on the Paranapanema River (state of São Paulo, Brazil), after an extreme inundation episode. *Acta Limnologica Brasiliensia*, *18*(4), 423-431.
- Fulan, J. A., Raimundo, R., & Figueiredo, D. (2008). Habitat characteristics and dragonflies diversity and abundance in the Guadiana River, eastern of the Alentejo, Portugal. *Boletín de la Asociación Española de Entomología*, *32*, 327-340.
- Furse, M. T., Morse, D., Wright, J. F., & Armitage, P. D. (1984). The influence of seasonal and taxonomic factors on the prediction to their macro-invertebrate communities. *Freshwater Biology*, *14*, 257 – 280.
- Gaufin, A. R. (1973). Use of aquatic invertebrates in the assessment or water quality. In J. J. Cairns, & K. L. Dickson (Eds.), *Biological methods for the assessment of waters quality, STP 528*, (pp. 96–116). Philadelphia: American Society for Testing and Materials.
- Gaufin, A. R., Clubb, R., & Newell, R. (1974). Studies on the tolerance of aquatic insect to low oxygen concentrations. *Great Basin Naturalist*, *34*, 45–59.

-
- Giribet, G., & Wheeler, W. C. (1999). On gaps. *Molecular Phylogenetics and Evolution*, *13*, 132 – 143.
- Gomez, A., Wright, P. J, Lunt, D. H., Canchino, J. M., Calvalho, G. R., & Hughes, R. N. (2007). Mating trials validate the use of DNA barcoding to reveal cryptic speciation of a marine bryozoan taxon. *Proceedings of the Royal Society B: Biological Sciences*, *274*, 199-346.
- Grimaldi, D., & Engel, M. S. (2005). *Evolution of the insects*. New York: Cambridge University Press.
- Groeneveld, L. F., Clausnitzer, V., & Hadryz, H. (2007). Convergent evolution of gigantism in damselflies of Africa and South America? Evidence from nuclear and mitochondrial sequence data. *Molecular Phylogenetics and Evolution*, *42*, 339 – 346.
- Gupta, I. J., De, M. L., & Mitra, T. R. (1995). Conspectus of Odonata fauna of Calcutta, India. *Records of the Zoological Survey of India*, *95*, 107 – 121.
- Hadryz, H., Balick, M., & Schierwater, B. (1992). Applications of random amplified polymorphic DNA (RAPD) in molecular ecology. *Molecular Ecology*, *1*, 55–63.
- Hadryz, H., Clausnitzer, V., & Groeneveld, L. V. (2006). The present role and future promise of conservation genetics for forest Odonates. In A. Rivera (Ed.) *Forest and dragonflies* (pp. 279-299). Sofia, Bulgaria: Pensoft Publishers.
- Hajibabaei, M., Janzen, D. H., Burns, J. M., Hallwachs, W., & Hebert, P. D. (2006). DNA barcodes distinguish species of tropical Lepidoptera. *Proceedings of the National Academy of Sciences, USA*, *103*, 968-971.

-
- Hamalainen, M. (1994), Dragonflies of Mount Kinabalu (the highest mountain in Borneo). *Malangpo*, 11, 77 – 81.
- Hawkes, H. A. (1979). Invertebrates as indicator of river water quality. In *Biological Indicators of Water Quality*, James A, Evison L (eds). John Wiley: Chichester; 17–61.
- Hawking, J. H., & New, T. R. (1999). The distribution patterns of dragonflies (Insecta: Odonata) along the Kiewa River, Australia, and their relevance in conservation assessment. *Hydrobiologia*, 392, 249 – 260.
- Hawking, J. H., & New, T. R. (2002) Interpreting dragonfly diversity to aid in conservation assessment: Lessons from the odonata assemblage at Middle Creek, north eastern Victoria, Australia. *Journal Insect Conservation*, 6, 171 - 178.
- Hayashi, F., Dobata, S., & Futahashi, R. (2004). Macro and microscale distribution patterns of two closely related Japanese *Mnais* species inferred from nuclear ribosomal DNA, ITS sequence and morphology (Zygoptera: Calopterigidae). *Odonatologica*, 33, 399 – 412.
- Hayashi, F., Dobata, S., & Futahashi, R. (2005). Distributed population genetics: Suspected introgressive hybridization between two *Mnais* damselfly species (Odonata). *Zoological Science*, 22, 869 – 881.
- Hebert, P. D., Cywinska, A., Ball S. L, & deWaard, J. R. (2003). Biological identifications through dna barcodes. *Proceedings of the Royal Society B: Biological Sciences*, 270, 313-321.

-
- Hebert, P. D., Penton, E. H., Burns, J. M., Janzen, D. H., & Hallwachs, W. (2004) Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes fulgerator*. *Proceedings of the National Academy of Sciences, USA*. 101, 14812-14817.
- Hellawell, J. M. (1986). Biological Indicators of Freshwater Pollution and Environmental Management, Mellanby K (ed.). Pollution Monitoring Series. Elsevier: Amsterdam.
- Hennig, W. (1969). *Die stammesgeschichte der insekten*. Frankfurt, Germany: Kramer.
- Hennig, W. (1981). *Insect phylogeny*. New York: Wiley.
- Hillis, D. M. (1987). Molecular versus morphological approaches to systematics. *Annual Review of Ecology and Systematics*, 18, 23 – 42.
- Hillis, D. M., Mable, B. K., Larson, A., Davis, S. K., & Zimmer, E. A. (1996). Nucleic acids IV: sequencing and cloning. In D. M. Hillis, M. Craig, M. H. David MH (Eds.), *Molecular systematic* (pp 321 – 339). Sunderland, Mass: Sinauer Associates.
- Hilsenhoff, W. L. (1988). Seasonal correction factors for the biotic index. *Great Lakes Entomologist*, 21, 9 – 13.
- Hoback, W. W., & Stanley, D. W. (2001). Mini review: insects in hypoxia. *Journal of Insect Physiology*, 47, 533-542.
- Hofmann, T. A., & Mason, C. F. (2005). Habitat characteristics and the distribution of Odonata in a lowland river catchment in eastern England. *Hydrobiologia*, 539, 137-147.

-
- Hovmöller, R., & Johansson, F. (2004). A phylogenetic perspective on larval spine morphology in *Leucorrhinia* (Odonata: Libellulidae) based on ITS1, 5.8S, and ITS2 rDNA sequences. *Molecular Phylogenetics and Evolution*, 30, 653 – 662.
- Johansson, F., & Samuelsson, L. (1994). Fish induced variation in abdominal spine length of *Leucorrhinia dubia* (Odonata) larvae? *Oecologia*, 100, 74 – 79.
- Jordan, S., Simon, C., & Polhemus, D. (2003). Molecular systematics and adaptive radiation of Hawaii's endemic damselfly genus *Megalagrion* (Odonata: Coenagrionidae). *Systematic Biology*, 52, 89 – 109.
- Kalkman, V. J., Clausnitzer, V., Dijkstra, K-D. B., Orr, A. G., Paulson, D. R., & van Tol, J. (2008). Global diversity of dragonflies (Odonata) in freshwater. *Hydrobiologia*, 595, 351 – 363.
- Kimura, M. (1980). A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16, 111-120.
- Kiyoshi, T., & Sota, T. (2006). Differentiation of the dragonfly genus *Davidus* (Odonata: Gomphidae) in Japan inferred from mitochondrial and nuclear gene genealogies. *Zoological Science*, 23, 1 – 8.
- Kluge, A. G. (1989). A concern for evidence and a phylogenetic hypothesis of relationships among *Epicrates* (Boidae, Serpentes). *Systematic Zoology*, 38, 7 – 25.

- Kress, W. J., Wurdack, K. J., Zimmer, E. A., Weigt, L. A., & Janzen, D. H. (2005). Use of DNA barcodes to identify flowering plants. *Proceedings of the National Academy of Sciences, USA*, 102, 8369-8374.
- Lenat, D. R. (1993). A biotic index for Southeastern United States: derivation and lists of tolerance value with criteria for assigning water quality ratings. *Journal of the North American Benthological Society*, 12, 279 – 290.
- Lieftinck, M. A. (1954). Handlist of Malaysian Odonata. *Treubia*, 22, 1- 202.
- Lim, R. P., & Furtado, J. I. (1975). Population changes in the aquatic fauna inhabiting the bladderwort, *Utricularia flexuosa* VAHL., in a tropical swamp, Tasek Bera, Malaysia. *Verhandlungen des Internationalen Verein Limnologie*, 19, 1390-1397.
- Lohmann, H. (1996). Das phylogenetische system der Anisoptera (Odonata). *Entomologische Zeitschrift*, 106(6), 209 – 252.
- Loxdale, H. D., & Lushai, G. (1998). Molecular markers in entomology. *Bulletin of Entomological Research*, 88, 577 - 600.
- Ludwig, J. A., & Reynolds, J. F. (1988). Statistical ecology. A primer on method and computing. John Wiley & Sons. Inc.
- Lutz, P. A. (1974). Effects of Temperature and Photoperiod on Larval Development in *Tetragoneuria Cynosura* (Odonata: Libellulidae). *Ecology*, 55(2), 370-377.

-
- Macan, T. T. (1974). Twenty generations of *Pyrrhosoma nymphula* (Sulzer) and *Enallagma cyathigerum* (Charpentier) (Zygoptera: Coenagrionidae). *Odonatologica*, 23, 115 – 124.
- Magurran, A. E. (1988). *Ecological Diversity and its Measurement*. Chapman and Hall, London.
- Metcalf, J. L. (1989). Biological water quality assessment of running waters based on microinvertebrates communities: History and present status in Europe. *Environmental Pollution*, 60, 101 – 139.
- Mikolajewski, D. J., & Johansson, F. (2002). Morphological and behavioral defenses in dragonfly larvae: trait compensation and cospecialization. *Behavioral Ecology* 15(4), 614 – 620.
- Miller, P. L. (1987). *Dragonflies*. Cambridge: Cambridge University Press.
- Misof, B., Rickert, A. M., Buckley, T. R., Fleck, G., & Sauer, K. P. (2001). Phylogenetic signal and its decay in mitochondrial SSU and LSU rRNA gene fragments of Anisoptera. *Molecular Biology and Evolution*, 18, 27-37.
- Misof, B. (2000). Diversity of Anisoptera (Odonata): inferring speciation process from patterns of morphological diversity. *Zoology (Jena)*, 105 (4), 355 – 365.
- Misof, B., Anderson, C. L., & Hadrys, H. (2000). A phylogeny of the damselfly genus *Calopteryx* (Odonata) using mitochondrial 16S rDNA markers. *Molecular Phylogenetics and Evolution*, 15, 5 -14.
- Miyamoto, M. M., & Fitch, W. M. (1995). Testing species phylogenies and phylogenetic methods with congruence. *Systematic Biology*, 44, 64 – 76.

-
- Moore, N. W. (1984). Dragonflies as indicators of environmental health. *IUCN Species Survival Commission Newsletter 1984*, 7 – 8.
- Moore, N. W. (1997). Status survey and conservation action plan: Dragonflies. IUCN/SSC Odonata Specialist Group, IUCN, Gland, Switzerland and Cambridge, UK.
- Needham, J. G., Westfall, Jr, M. J., & May, M. L. (2000). Dragonflies of North America. Scientific Publishers, Gainesville, FL.
- Nei, M., & Kumar, S. (2000). Molecular Evolution and Phylogenetics. Oxford University Press, New York.
- Nel, A., & Paicheler, J. C. (1993). Les Libellulidae fossils. Un inventaire critique (Odon., Anisoptera, Libelluloidea). *Entomologica Gallica*, 4, 166 – 190.
- Nel, A., & Paicheler. (1994). Les Libelluloidea autres que Libellulidae fossils. Un inventaire critique (Odonata, Corduliidae, Macromiidae, Synthemistidae, Chlorogomphidae et Mesophlebiidae). *Nouvelle Revue d'Entomologie (N.S.)*, 11 (4), 321 – 334.
- Norma-Rashid, Y. (1995a). Preliminary odonate survey of Perak, Peninsular Malaysia. *Notulae Odonatologica*, 4, 80-82
- Norma-Rashid, Y. (1995b). A dynamic encounter with the odonates in Sungai Seliah, Kelantan, Peninsular Malaysia. *Wallaceana*. 76, 17-22.
- Norma-Rashid, Y., Zakaria-Ismail, M., & Hamalainen, M. (1996). Odonate fauna from Kelantan, Pahang and Muar drainages, Malaysia. Proc. 5th Nat Biol Symp. UKN, Malaysia.

-
- Norma-Rashid, Y. (1998). Animal of Malaysia: dragonflies and damselflies. In Encyclopedia of Malaysia – *Wildlife of Malaysia*. 3, 104-105.
- Norma-Rashid, Y. (1999). Behavioural ecology of *Tyriobapta torrida* Kirby (Anisoptera: Libellulidae) at the breeding and resting sites. *Odonatologia*. 28, 139-150.
- Norma-Rashid, Y., Mohd-Sofian, A., & Zakaria-Ismail, M. (2001). Diversity and distribution of Odonata (dragonflies and damselflies) in the fresh water swamp lake Tasek Bera, Malaysia. *Hydrobiologia*, 459(12), 135-146.
- Norma-Rashid, Y., Cheong, L. F., Lua, H. K., & Murphy, D. H. (2008). The Dragonflies (Odonata) of Singapore Current Status Records and Collections of the Raffles Museum of Biodiversity Research.
- Norma-Rashid, Y. (2010). Dragonflies (Odonata) of Bachok Coast, Kelantan and Promoting Common Names. *Malaysian Journal of Science*, 29, 73-79.
- Oertli, B., Indermuehle, N., Angélibert, S., Hinden, H., & Stoll, A. (2008). Macroinvertebrate assemblages in 25 high alpine ponds of the Swiss National Park (Cirque of Macun) and relation to environmental variables. *Hydrobiologia*, 597, 29 – 41.
- Olmstead, R. G., & Sweere, J. A. (1994). Combining data in phylogenetic systematics: An empirical approach using three molecular data sets in the Solonaceae. *Systematic Biology*, 42, 467 – 481.

-
- Orr, A. G., Butler, G. G., Hamalainen, M., & Kemp, R. G. (2004). Insecta: Odonata. In M. Y. Catherine & H. S. Yong. *Freshwater Invertebrates of the Malaysian Region* (pp. 409-442). Kuala Lumpur: Academy of Sciences.
- Ott, J. (2001). Expansion of Mediterranean Odonata in Germany and Europe: Consequences of climate changes. In G. -R Walther, C. A. Burga & P. J. Edwards (Eds.). *Fingerprints' of climate change: Adapted behaviour and shifted species ranges* (pp. 89-111). New York: Kluwer Academic/Plenum Publishers..
- Ott, J. (2009). *Dragonflies and climate change*. Sofia, Bulgaria: Pensoft.
- Palmer, M. W. (1995). How should one count species? *Natural Areas Journal*, 15, 124 – 135.
- Palumbi, S. R. (1996). Nucleic acids. II. The polymerase chain reaction. In D. M. Hillis, C. Moritz, & B. K. Mable (Eds.). *Molecular Systematics* (pp. 205–247). Sunderland, MA: Sinauer Associates.
- Pashley, D. P., & Li, D. K. (1992). Sequence evolution in mitochondrial ribosomal and ND-1 genes in Lepidoptera: Implications for phylogenetic analyses. *Molecular Biology and Evolution*, 9(6), 1061 – 1075.
- Pfau, H. K. (1991). Contributions of functional morphology to the phylogenetic systematic of Odonata. *Adv. Odonatol.*, 5, 109-141.
- Pielou, E. C. (1975). *Ecological diversity*. New York: Wiley.
- Pollard, J. B. & Berril, M. (1992). The distribution of dragonfly nymphs across a pH gradient in South-Central Ontario Lakes, *Canadian Journal of Zoology*, 70, 878–885.

-
- Rach, J., DeSalle, R., Sarkar, I. N., Schierwater, B., & Hadrys, H. (2007). Character-based DNA barcoding allows discrimination of genera, species and populations in Odonata. *Proceedings of The Royal Society*, 275, 237-247.
- Rambur, P. (1842) *Histoire naturelle des insectes: Neuroptères. insectes neuroptères*. Paris: Librairie Encyclopédique de Roret.
- Rehn, A.C. (2003). Phylogenetic analysis of higher-level relationships of Odonata. *Systematic Entomology*, 28, 181 – 239.
- Rith-Najarian, J. C. (1998). The influence of forest vegetation variables on the distribution and diversity of dragonflies in a northern Minnesota forest landscape: A preliminary study (Anisoptera). *Odonatologica*, 27, 335 – 351.
- Roux, K. H. (1995). *Optimization and troubleshooting in PCR*. Cold Spring Harbor: Laboratory Press.
- Rubinoff, D. (2006a). DNA barcoding evolves into the familiar. *Conservation Biology*, 20, 1548–1549.
- Rubinoff, D. (2006b). Utility of mitochondrial DNA barcodes in species conservation. *Conservation Biology*, 20, 1026 – 1033.
- Sahlén, G. (1999). The impact of forestry on dragonfly diversity in Central Sweden. *The International Journal of Odonatology*, 2, 177 – 186.
- Sahlén, G., & Katarina, E.(2001). Identification of dragonflies (Odonata) as indicators of general species richness in boreal forest lakes. *Biodiversity and Conservation*, 10, 673 – 690.

-
- Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing trees. *Molecular Biology and Evolution*, *4*, 406 – 425.
- Sarot, E. E. (1958). *Folklore of the dragonfly: A linguistic approach*. Rome: Storia e Letteratura.
- Sato, M., & Riddiford, N. (2007). A preliminary study of the Odonata of S'Albufera Natural Park, Mallorca: Status, conservation priorities and bioindicator potential. *Journal Insect Conservation*, *12*(5), 539 – 548.
- Schinder, M., Fesl, C., & Chovanec, A. (2003). Dragonfly association (Insecta: Odonata) in relation to habitat variables: A multivariate approach. *Hydrobiologia*, *497*, 169 – 180.
- Schmidt, E. (1985). Habitat inventarization, characterization and bioindication by a Representative Spectrum of Odonata species (RSO). *Odonatologica*, *14*, 127 – 133.
- Shelton, M. D., & Edwards, C. R. (1983). Effects of weeds on the diversity and abundance of insects in soybeans. *Environmental Entomology*, *12*, 296 – 299.
- Silsby, J. (2001). *Dragonflies of the world*. Washington D. C.: Smithsonian Institution Press.
- Simaika, J. P., & Samways, M. J. (2001). Comparative assessment of indices of freshwater habitat conditions using different invertebrate taxon sets. *Ecological Indicator*, *11*, 371 – 378.

-
- Simaika, J. P., & Samways, M. J. (2009). An easy-to-use index of ecological integrity for prioritizing freshwater sites and for assessing habitat quality. *Biodiversity Conservation* 18, 1171 – 1185.
- Simmons, M. P., & Ochoterena, H. (2000). Gaps as characters in sequence-based phylogenetic analyses. *Systematic Biology*, 49, 369 – 381.
- Simpson, E. H. (1949). *Measurement of diversity*. London: Nature.
- Smith, B., & Wilson, J. B. (1996). A consumer's guide to evenness indices. *Oikos*, 76, 70 – 82.
- Smith, M. A., Woodley, N. E., Janzen, D. H., Hallwachs, W., & Hebert, P. D. (2006). DNA barcodes reveal cryptic host-specificity within the presumed polyphagous members of a genus of parasitoid flies (Diptera: Tachnidae). *Proceedings of the National Academy of Sciences, USA*, 103, 3657-3662.
- Smolka, G. E., Stewart, P. M., & Swinford, T. O. (1999). Distribution of odonates (Dragonflies and Damselflies) in the Indiana Dunes National Lakeshore and Nearby Lands. *Natural Areas Journal*, 19, 132 – 141.
- Southwood, T. R. E., & Henderson, P. A. (2000). *Ecological Methods*. Oxford, U.K: Blackwell.
- Steinmann, H. (1997). World catalogue of Odonata. Vol. II. Anisoptera. In M. Fischer (Ed.) *The animal kingdom: A compilation and characterization of the recent animal groups*, (Part III). Berlin: Walter de Gruyter.

-
- Tamura, K., Dudley, J., Nei, M., & Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, 24, 1596-1599.
- Teles, L. F. O. (1994). A new methodology for biological water quality assessment *International association of theoretical and applied limnology*, 25, 1942 – 1944.
- Tennessen, K. J. (1997). The rate of species descriptions in Odonata. *Entomological News*, 108, 122 – 126.
- Thompson, J., Higgins, D., & Gibson, T. (1994.) CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22, 4673-4680.
- Tillyard, R. J. (1917). The biology of dragonflies. London: Cambridge University Press.
- Tollyard, R. J. (1935). The biology of dragonflies (Odonata or Paraneuroptera). Cambridge: Cambridge University Press.
- Trueman, J. W. H. (1996). A preliminary cladistic analysis of Odonate wing venation. *Odonologica*, 25, 59-72.
- Tsuda, S. (2000). A distributional list of world Odonata 2000. Osaka: Tsuda. S.
- Turgeon, J., & McPeck, M. A. (2002). Phylogeographic analysis of a recent radiation of *Enballagma* damselflies (Odonata: Coenagrionidae). *Molecular Ecology*, 11, 1989 – 2001.

-
- Volger, A. P., & Welsh, A. (1997). Phylogeny of North American *Cicindela* tiger beetles inferred from multiple mitochondrial DNA sequences. *Molecular Phylogenetics and Evolution*, 8, 225 – 235.
- Voshell, J. R., & Simmons, G. M. (1978). The Odonata of a new Reservoir in the southeastern United States. *Odonatologica*, 7(1), 67-76.
- Ward, J. V. (1992). *Aquatic insect ecology*. New York: John Wiley and Sons. 466 p.
- Ward, R. D., Zemplak, T. S., Innes, B. H., Last, P. R., & Hebert, P. D. (2005). DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360, 1847-1857.
- Ware, J., May, M., & Kjer, K. (2007). Phylogeny of the higher Libelluloidea (Anisoptera: Odonata): An exploration of the most speciose super family of dragonflies. *Molecular Phylogenetics and Evolution*, 45, 289 – 310.
- Watts, P. C., Rouquette, R., Saccheri, J., Kemp, J., & Thompson, D. J. (2004). Molecular and ecological evidence for small-scale isolation by distance in an endangered damselfly, *Coenagrion mercurial*. *Molecular Ecology*, 13, 2931 – 2945.
- Williams, D. D., & Feltmate, B. (1992). *Aquatic insects*. Wallingford, U. K: CAB International.
- William Bouchard, R., & Paul S. (2009). Guide to Aquatic Invertebrate Families of Mongolia. *Odonata (Dragonflies & Damselflies)*. (pp. 64 – 74).

- Witt, J. D, Threlhoff, D. L., Hebert, P. D. (2006). DNA barcoding reveals extraordinary cryptic diversity in an amphipod genus: implications for desert spring conservation. *Molecular Ecology*, 15, 3073-3082.
- Wong, A., Smith, M. L., Forbes, M. R. (2003). Differentiation between sub-populations of a polychromatic damselfly with respect to morph frequencies, but not neutral genetic markers. *Molecular Ecology*, 12, 3505 – 3513.
- Yap, C. K., Rahim Ismail A., Ismail A., & Tan, S. G. (2003). Species diversity of Macrobenthic invertebrates in the Semenyih River, Selangor, Peninsular Malaysia. *Pertanika Journal of Tropical Agricultural Science*, 26, 139 – 146.