

References

- Ait-Si-Ali**, S., Guasconi, V., Fritsch, L., Yahi, H., Sekhri, R., Naguibneva, I., Robin, P., Cabon, F., Polesskaya, A. & Harel-Bellan, A. (2004). A Suv39h-dependent mechanism for silencing S-phase genes in differentiating but not in cycling cells. *Embo J* **23**, 605-15.
- Alunni-Fabbroni, M**, Littlewood T, Deleu L, Caldeira S, Giarre M, Dell' Orco, M and Tommasino M, 2000. Induction of S phase and apoptosis by the human papillomavirus type 16 E7 protein are separable events in immortalized rodent fibroblasts. *Oncogene*, **19**:2277-2285
- Baker**, C. C., Phelps, W. C., Lindgren, V., Braun, M. J., Gonda, M. A. & Howley, P. M. (1987). Structural and transcriptional analysis of human papillomavirus type 16 sequences in cervical carcinoma cell lines. *J Virol* **61**, 962-71.
- Baldi**, A., De Luca, A., Claudio, P. P., Baldi, F., Giordano, G. G., Tommasino, M., Paggi, M. G. & Giordano, A. (1995). The RB2/p130 gene product is a nuclear protein whose phosphorylation is cell cycle regulated. *J Cell Biochem* **59**, 402-8.
- Bandara**, L. R. & La Thangue, N. B. (1991). Adenovirus E1a prevents the retinoblastoma gene product from complexing with a cellular transcription factor. *Nature* **351**, 494-7.
- Banerjee**, N. S., Genovese, N. J., Noya, F., Chien, W. M., Broker, T. R., and Chow, L. T. (2006). Conditionally activated E7 proteins of high-risk and low-risk human papillomaviruses induce S phase in postmitotic, differentiated human keratinocytes. *J Virol* **80**, 6517-6524.
- Barbosa**, M. S., Lowy, D. R. & Schiller, J. T. (1989). Papillomavirus polypeptides E6 and E7 are zinc-binding proteins. *J Virol* **63**, 1404-7.
- Barbosa**, M. S., Vass, W. C., Lowy, D. R. & Schiller, J. T. (1991). In vitro biological activities of the E6 and E7 genes vary among human papillomaviruses of different oncogenic potential. *J Virol* **65**, 292-8
- Beijersbergen**, R. L., Carlee, L., Kerkhoven, R. M. & Bernards, R. (1995). Regulation of the retinoblastoma protein-related p107 by G1 cyclin complexes. *Genes Dev* **9**, 1340-53.

- Beijersbergen**, R. L., Carlee, L., Kerkhoven, R. M. & Bernards, R. (1995). Regulation of the retinoblastoma protein-related p107 by G1 cyclin complexes. *Genes Dev* **9**, 1340-53.
- Berezutskaya**, E., Yu, B., Morozov, A., Raychaudhuri, P. & Bagchi, S. (1997). Differential regulation of the pocket domains of the retinoblastoma family proteins by the HPV16 E7 oncoprotein. *Cell Growth Differ* **8**, 1277-86.
- Bernard H.U.**, Burk R.D., Chen Z., Doorslaer K.V., zur Hausen H., & de Villiers E.M. (2010). Classification of papillomaviruses (PVs) based on 189 PV types and proposal of taxonomic amendments. *Virology* **410**, 70-79.
- Black**, E. P., Huang, E., Dressman, H., Rempel, R., Laakso, N., Asa, S. L., Ishida, S., West, M. & Nevins, J. R. (2003). Distinct gene expression phenotypes of cells lacking Rb and Rb family members. *Cancer Res* **63**, 3716-23.
- Boshart M**, Gissmann L, Ikenberg H, Kleinheinz A, Scheurlen W, zur Hausen H. 1984 A new type of papillomavirus DNA, its presence in genital cancer biopsies and in cell lines derived from cervical cancer. *EMBO J* 3(5):1151-7.
- Boyer**, S. N., Wazer, D. E. & Band, V. (1996). E7 protein of human papilloma virus-16 induces degradation of retinoblastoma protein through the ubiquitin-proteasome pathway. *Cancer Res* **56**, 4620-4.
- Brehm**, A., Nielsen, S. J., Miska, E. A., McCance, D. J., Reid, J. L., Bannister, A. J. & Kouzarides, T. (1999). The E7 oncoprotein associates with Mi2 and histone deacetylase activity to promote cell growth. *Embo J* **18**, 2449-58.
- Buckley**, C. H., Butler, E. B. & Fox, H. (1982). Cervical intraepithelial neoplasia. *J Clin Pathol* **35**, 1-13.
- Burd, E. M.** (2003). Human papillomavirus and cervical cancer. *Clin Microbiol Rev* **16**, 1-17.

- Caldeira S, de Villiers E M and Tommasino M,2000.** Human papillomavirus E7 proteins stimulate proliferation independently of their ability to associate with retinoblastoma protein. *Oncogene*, **19**:821-826
- Canhoto, A. J., Chestukhin, A., Litovchick, L. & DeCaprio, J. A. (2000).** Phosphorylation of the retinoblastoma-related protein p130 in growth-arrested cells. *Oncogene* **19**, 5116-22
- Cates, W., Jr. (1999).** Estimates of the incidence and prevalence of sexually transmitted diseases in the United States. American Social Health Association Panel. *Sex Transm Dis* **26**, S2-7.
- Cheng, M., Olivier, P., Diehl, J. A., Fero, M., Roussel, M. F., Roberts, J. M. & Sherr, C. J. (1999).** The p21(Cip1) and p27(Kip1) CDK 'inhibitors' are essential activators of cyclin D-dependent kinases in murine fibroblasts. *Embo J* **18**, 1571-83.
- Cheng, S., Schmidt-Grimminger, D. C., Murant, T., Broker, T. R. & Chow, L. T. (1995).** Differentiation-dependent up-regulation of the human papillomavirus E7 gene reactivates cellular DNA replication in suprabasal differentiated keratinocytes. *Genes Dev* **9**, 2335-49.
- Chow, K. N. & Dean, D. C. (1996).** Domains A and B in the Rb pocket interact to form a transcriptional repressor motif. *Mol Cell Biol* **16**, 4862-8.
- Ciccolini, F., Di Pasquale, G., Carlotti, F., Crawford, L. & Tommasino, M. (1994).** Functional studies of E7 proteins from different HPV types. *Oncogene* **9**, 2633-8.
- Ciechanover, A., Orian, A., and Schwartz, A. L. (2000).** Ubiquitin-mediated proteolysis: biological regulation via destruction. *Bioessays* **22**, 442-451.
- Ciuffo G.** 1907. Innesto positivo con filtrato di verucca volgare. *Giorn Ital Mal Venereol* **48**, 12-17
- Classon M & Harlow E. (2002).** The retinoblastoma tumour suppressor in development and cancer. *Nat. Rev. Cancer* **12**, 910-917.
- Cobrinik, D. (2005).** Pocket proteins and cell cycle control. *Oncogene* **24**, 2796-2809.

- Collins, A. S., Nakahara, T., Do, A., and Lambert, P. F.** (2005). Interactions with pocket proteins contribute to the role of human papillomavirus type 16 E7 in the papillomavirus life cycle. *J Virol* **79**, 14769-14780.
- Crawford LV and Crawford EM.** 1963. A comparative study of polyoma and papilloma viruses. *Virology* **21**:258-263.
- Davies, R., Hicks, R., Crook, T., Morris, J. & Vousden, K.** (1993). Human papillomavirus type 16 E7 associates with a histone H1 kinase and with p107 through sequences necessary for transformation. *J Virol* **67**, 2521-8.
- Delius, H., and Hofmann, B.** (1994). Primer-directed sequencing of human papillomavirus types. *Curr Top Microbiol Immunol* **186**, 13-31.
- Demers, G. W., Espling, E., Harry, J. B., Etscheid, B. G. & Galloway, D. A.** (1996). Abrogation of growth arrest signals by human papillomavirus type 16 E7 is mediated by sequences required for transformation. *J Virol* **70**, 6862-9
- Deshaias, R. J.** (1999). SCF and Cullin/Ring H2-based ubiquitin ligases. *Annu Rev Cell Dev Biol* **15**, 435-467.
- Dick Frederick A and Dyson Nicholas J** (2002). Three regions of the pRB pocket domain affect its inactivation by human papillomavirus E7 proteins. *J. Virol* **76**:12:6224.
- Dillner, J., Meijer, C. J., von Krogh, G., and Horenblas, S.** (2000). Epidemiology of human papillomavirus infection. *Scand J Urol Nephrol Suppl*, 194-200.
- Dimova, D. K., Stevaux, O., Frolov, M. V. & Dyson, N. J.** (2003). Cell cycle-dependent and cell cycle-independent control of transcription by the Drosophila E2F/RB pathway. *Genes Dev* **17**, 2308-20.
- Dong W L, Caldeira S, Sehr P, Pawlita M and Tommasino M, 2001.** Determination of the binding affinity of different human papillomavirus E7 proteins for the tumour suppressor pRb by a plate-binding assay. *J Virol Methods*, **98**:91-98

- Doorbar J, Elston RC, Napthine S, Raj K, Medcalf E, Jackson D, Coleman N, Griffin HM, Masterson P, Stacey S, Mengistu Y, Dunlop J.** 2000 The E1E4 protein of human papillomavirus type 16 associates with a putative RNA helicase through sequences in its C terminus. *J Virol.* 74(21):10081-95.
- Doorbar J, Ely S, Sterling J, McLean C. and Crawford L.** 1991. Specific interaction between HPV-16 E1-E4 and cytokeratins results in collapse of the epithelial cell intermediate filament network *Nature* 352, 824 - 827.
- Doorbar, J.** (2006). Molecular biology of human papillomavirus infection and cervical cancer. *Clin Sci (Lond)* 110, 525-541.
- Drain, P. K., Holmes, K. K., Hughes, J. P., and Koutsky, L. A.** (2002). Determinants of cervical cancer rates in developing countries. *International Journal of Cancer* 100, 199-205.
- D'Souza, G., Kreimer, A. R., Viscidi, R., Pawlita, M., Fakhry, C., Koch, W. M., Westra, W. H., and Gillison, M. L.** (2007). Case-control study of human papillomavirus and oropharyngeal cancer. *N Engl J Med* 356, 1944-1956.
- Dunaief, J. L., Strober, B. E., Guha, S., Khavari, P. A., Alin, K., Luban, J., Begemann, M., Crabtree, G. R. & Goff, S. P.** (1994). The retinoblastoma protein and BRG1 form a complex and cooperate to induce cell cycle arrest. *Cell* 79, 119-30.
- Durst M, Croce CM, Gissmann L, Schwarz E, Huebner K.** 1987. Papillomavirus sequences integrate near cellular oncogenes in some cervical carcinomas. *Proc Natl Acad Sci U S A.* 84(4):1070-4.
- Dyson, N., and E. Harlow (ed.).** 1992. Adenovirus E1A targets key regulators of cell proliferation. Imperial Cancer Research Fund, London, United Kingdom.
- Dyson, N., Guida, P., Munger, K. & Harlow, E.** (1992). Homologous sequences in adenovirus E1A and human papillomavirus E7 proteins mediate interaction with the same set of cellular proteins. *J Virol* 66, 6893-902.

- Dyson, N.**, Howley, P. M., Munger, K., and Harlow, E. (1989). The human papilloma virus-16 E7 oncoprotein is able to bind to the retinoblastoma gene product. *Science* **243**, 934-937.
- Ferreira, R.**, Magnaghi-Jaulin, L., Robin, P., Harel-Bellan, A. & Trouche, D. (1998). The three members of the pocket proteins family share the ability to repress E2F activity through recruitment of a histone deacetylase. *Proc Natl Acad Sci U S A* **95**, 10493-8.
- Flores ER, Allen-Hoffmann BL, Lee D, Lambert PF.** 2000. The human papillomavirus type 16 E7 oncogene is required for the productive stage of the viral life cycle. *J Virol.* 74(14):6622-31.
- Funk, J. O.**, Waga, S., Harry, J. B., Espling, E., Stillman, B. & Galloway, D. A. (1997). Inhibition of CDK activity and PCNA-dependent DNA replication by p21 is blocked by interaction with the HPV-16 E7 oncoprotein. *Genes Dev* **11**, 2090-100.
- Gage, J. R.**, Meyers, C. & Wettstein, F. O. (1990). The E7 proteins of the nononcogenic human papillomavirus type 6b (HPV-6b) and of the oncogenic HPV-16 differ in retinoblastoma protein binding and other properties. *J Virol* **64**, 723-30.
- Gagriva, S.**, Hauser, S., Kolfschoten, I., Osterloh, L., Agami, R. & Gaubatz, S. (2004). Inhibition of oncogenic transformation by mammalian Lin-9, a pRB-associated protein. *Embo J* **23**, 4627-38
- Genovese N.J.**, Banerjee N.S., Broker T.R & Chow L.T. (2008). Casein kinase II motif-dependent phosphorylation of human papillomavirus E7 protein promotes p130 degradation and S-phase induction in differentiated human keratinocytes. *J. Virol* **82**, 4862-4873.
- Giarrè, M.**, Caldeira, S., Malanchi, I., Ciccolini, F., Leao, M. J. & Tommasino, M. (2001). Induction of pRb degradation by the human papillomavirus type 16 E7 protein is essential to efficiently overcome p16INK4a-imposed G1 cell cycle Arrest. *J Virol* **75**, 4705-12
- Gissmann L**, Pfister H, Zur Hausen H. 1977. Human papilloma viruses (HPV): characterization of four different isolates. *Virology*. 76(2):569-80.

- Glickman, M. H., and Ciechanover, A.** (2002). The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction. *Physiol Rev* **82**, 373-428.
- Gonzalez, S. L., Stremlau, M., He, X., Basile, J. R. & Munger, K.** (2001). Degradation of the retinoblastoma tumor suppressor by the human papillomavirus type 16 E7 oncoprotein is important for functional inactivation and is separable from proteasomal degradation of E7. *J Virol* **75**, 7583-91.
- Gonzalez, S. L., Stremlau, M., He, X., Basile, J. R. & Munger, K.** (2001). Degradation of the retinoblastoma tumor suppressor by the human papillomavirus type 16 E7 oncoprotein is important for functional inactivation and is separable from proteasomal degradation of E7. *J Virol* **75**, 7583-91.
- Halbert, C. L., Demers, G. W. & Galloway, D. A.** (1991). The E7 gene of human papillomavirus type 16 is sufficient for immortalization of human epithelial cells. *J Virol* **65**, 473-8.
- Harbour, J. W., Luo, R. X., Dei Santi, A., Postigo, A. A. & Dean, D. C.** (1999). Cdk phosphorylation triggers sequential intramolecular interactions that progressively block Rb functions as cells move through G1. *Cell* **98**, 859-69.
- Harrison, M. M., Ceol, C. J., Lu, X. & Horvitz, H. R.** (2006). Some *C. elegans* class B synthetic multivulva proteins encode a conserved LIN-35 Rb-containing complex distinct from a NuRD-like complex. *Proc Natl Acad Sci U S A* **103**, 16782-7.
- Hebner, C. M., and Laimins, L. A.** (2006). Human papillomaviruses: basic mechanisms of pathogenesis and oncogenicity. *Rev Med Virol* **16**, 83-97.
- Helt, A. M., Funk, J. O. & Galloway, D. A.** (2002). Inactivation of both the retinoblastoma tumor suppressor and p21 by the human papillomavirus type 16 E7 oncoprotein is necessary to inhibit cell cycle arrest in human epithelial cells. *J Virol* **76**, 10559-68

- Herrera**, R. E., Sah, V. P., Williams, B. O., Makela, T. P., Weinberg, R. A. & Jacks, T. (1996). Altered cell cycle kinetics, gene expression, and G1 restriction point regulation in Rb-deficient fibroblasts. *Mol Cell Biol* **16**, 2402-7.
- Hoeppner, S., Baumli, S., and Cramer, P.** (2005). Structure of the mediator subunit cyclin C and its implications for CDK8 function. *J Mol Biol* **350**, 833-842.
- Hughes FJ, Romanos MA.** 1993. E1 protein of human papillomavirus is a DNA helicase/ATPase. *Nucleic Acids Res.* **21**(25):5817-23.
- Huibregtse, J. M., Scheffner, M., and Howley, P. M.** (1993). Cloning and expression of the cDNA for E6-AP, a protein that mediates the interaction of the human papillomavirus E6 oncoprotein with p53. *Mol Cell Biol* **13**, 775-784.
- Human papillomavirus and related cancer (2010). WHO.
- Hurford, R. K., Jr., Cobrinik, D., Lee, M. H. & Dyson, N.** (1997). pRB and p107/p130 are required for the regulated expression of different sets of E2F responsive genes. *Genes Dev* **11**, 1447-63.
- Ishida, S., Huang, E., Zuzan, H., Spang, R., Leone, G., West, M. & Nevins, J. R.** (2001). Role for E2F in control of both DNA replication and mitotic functions as revealed from DNA microarray analysis. *Mol Cell Biol* **21**, 4684-99.
- Jewers, R. J., Hildebrandt, P., Ludlow, J. W., Kell, B. & McCance, D. J.** (1992). Regions of human papillomavirus type 16 E7 oncoprotein required for immortalization of human keratinocytes. *J Virol* **66**, 1329-35
- Jones, D. L., Alani, R. M. & Munger, K.** (1997). The human papillomavirus E7 oncoprotein can uncouple cellular differentiation and proliferation in human keratinocytes by abrogating p21Cip1-mediated inhibition of cdk2. *Genes Dev* **11**, 2101-11.

- Kim, G. Y., Mercer, S. E., Ewton, D. Z., Yan, Z., Jin, K. & Friedman, E.** (2002). The stress-activated protein kinases p38 alpha and JNK1 stabilize p21(Cip1) by phosphorylation. *J Biol Chem* **277**, 29792-802.
- Knight A S, Notaridou M and Watson R J, 2009.** A Lin9 complex is recruited by B-myb to activate transcription of G2/M genes in undifferentiated embryonal carcinoma cells. *Oncogene* **28**:1737-1747
- Korenjak M, Taylor-Harding B, Binné UK, Satterlee JS, Stevaux O, Aasland R, White-Cooper H, Dyson N, Brehm A** (2004). Native E2F/RBF complexes contain Myb-interacting proteins and repress transcription of developmentally controlled E2F target genes. *Cell* ;119(2):181-93.
- Krek, W., Ewen, M. E., Shirodkar, S., Arany, Z., Kaelin, W. G., Jr. & Livingston, D. M.** (1994). Negative regulation of the growth-promoting transcription factor E2F-1 by a stably bound cyclin A-dependent protein kinase. *Cell* **78**, 161-72.
- Krek, W., Xu, G. & Livingston, D. M.** (1995). Cyclin A-kinase regulation of E2F-1 DNA binding function underlies suppression of an S phase checkpoint. *Cell* **83**, 1149-58
- LaBaer, J., Garrett, M. D., Stevenson, L. F., Slingerland, J. M., Sandhu, C., Chou, H. S., Fattaey, A. & Harlow, E.** (1997). New functional activities for the p21 family of CDK inhibitors. *Genes Dev* **11**, 847-62
- Lam EW; Morris JD; Davies R; Crook T; Watson RJ; Vousden KH. 1994.** HPV16 E7 oncoprotein deregulates B-myb expression: correlation with targeting of p107/E2F complexes. *EMBO J.* **13**:871-878
- Law MF, Lancaster WD, Howley PM.** 1979. Conserved polynucleotide sequences among the genomes of papillomaviruses. *J Virol.* 32(1):199-207.
- Lee D, Sohn H, Kalpana GV, Choe J.** 1999. Interaction of E1 and hSNF5 proteins stimulates replication of human papillomavirus DNA. *Nature.* 399(6735):487-91.

- Lee**, E. Y., Chang, C. Y., Hu, N., Wang, Y. C., Lai, C. C., Herrup, K., Lee, W. H. & Bradley, A. (1992). Mice deficient for Rb are nonviable and show defects in neurogenesis and haematopoiesis. *Nature* **359**, 288-94.
- Lee**, E. Y., Chang, C. Y., Hu, N., Wang, Y. C., Lai, C. C., Herrup, K., Lee, W. H. & Bradley, A. (1992). Mice deficient for Rb are nonviable and show defects in neurogenesis and haematopoiesis. *Nature* **359**, 288-94
- Lee**, J. O., Russo, A. A. & Pavletich, N. P. (1998). Structure of the retinoblastoma tumour-suppressor pocket domain bound to a peptide from HPV E7. *Nature* **391**, 859-65
- Levine, A. J.** (1997). p53, the cellular gatekeeper for growth and division. *Cell* **88**, 323-331.
- Lewis**, P. W., Beall, E. L., Fleischer, T. C., Georlette, D., Link, A. J. & Botchan, M. R. (2004). Identification of a Drosophila Myb-E2F2/RBF transcriptional repressor complex. *Genes Dev* **18**, 2929-40.
- Litovchick L**, Sadasivam S, Florens L, Zhu X, Swanson SK, Velmurugan S, Chen R, Washburn MP, Liu XS, DeCaprio JA (2007). Evolutionarily conserved multisubunit RBL2/p130 and E2F4 protein complex represses human cell cycle-dependent genes in quiescence. *Mol Cell*;26(4):539-51.
- Longworth, M. S., and Laimins, L. A.** (2004). Pathogenesis of human papillomaviruses in differentiating epithelia. *Microbiol Mol Biol Rev* **68**, 362-372.
- Ludlow**, J. W., Glendening, C. L., Livingston, D. M. & DeCaprio, J. A. (1993). Specific enzymatic dephosphorylation of the retinoblastoma protein. *Mol Cell Biol* **13**, 367-72.
- Luo**, R. X., Postigo, A. A. & Dean, D. C. (1998). Rb interacts with histone deacetylase to repress transcription. *Cell* **92**, 463-73
- Magnaghi-Jaulin**, L., Groisman, R., Naguibneva, I., Robin, P., Lorain, S., Le Villain, J. P., Troalen, F., Trouche, D. & Harel-Bellan, A. (1998). Retinoblastoma protein represses transcription by recruiting a histone deacetylase. *Nature* **391**, 601-5.

- Matlashewski, G.**, Schneider, J., Banks, L., Jones, N., Murray, A. & Crawford, L. (1987). Human papillomavirus type 16 DNA cooperates with activated ras in transforming primary cells. *Embo J* **6**, 1741-6.
- Matsukura, T.**, Kanda, T., Furuno, A., Yoshikawa, H., Kawana, T. & Yoshiike, K. (1986). Cloning of monomeric human papillomavirus type 16 DNA integrated within cell DNA from a cervical carcinoma. *J Virol* **58**, 979-82
- Matsushime, H.**, Roussel, M. F., Ashmun, R. A. & Sherr, C. J. (1991). Colony-stimulating factor 1 regulates novel cyclins during the G1 phase of the cell cycle. *Cell* **65**, 701-13.
- Mayol, X.**, Garriga, J. & Grana, X. (1995). Cell cycle-dependent phosphorylation of the retinoblastoma-related protein p130. *Oncogene* **11**, 801-8
- McIntyre, M. C.**, Frattini, M. G., Grossman, S. R. & Laimins, L. A. (1993). Human papillomavirus type 18 E7 protein requires intact Cys-X-X-Cys motifs for zinc binding, dimerization, and transformation but not for Rb binding. *J Virol* **67**, 3142-50.
- Missero, C.**, Di Cunto, F., Kiyokawa, H., Koff, A. & Dotto, G. P. (1996). The absence of p21Cip1/WAF1 alters keratinocyte growth and differentiation and promotes ras-tumor progression. *Genes Dev* **10**, 3065-75
- Moberg, K.**, Starz, M. A. & Lees, J. A. (1996). E2F-4 switches from p130 to p107 and pRB in response to cell cycle reentry. *Mol Cell Biol* **16**, 1436-49.
- Muller, H.**, Bracken, A. P., Vernell, R., Moroni, M. C., Christians, F., Grassilli, E., Prosperini, E., Vigo, E., Oliner, J. D. & Helin, K. (2001). E2Fs regulate the expression of genes involved in differentiation, development, proliferation, and apoptosis. *Genes Dev* **15**, 267-85
- Munger, K. et al. (1989).** The E6 and E7 genes of human papillomavirus type 16 are necessary and sufficient for transformation of primary human keratinocytes. *Journal of Virology*, **63**, 4417-4423.

- Munger, K., and Howley, P. M.** (2002). Human papillomavirus immortalization and transformation functions. *Virus Res* **89**, 213-228.
- Munger, K.**, Yee, C. L., Phelps, W. C., Pietenpol, J. A., Moses, H. L. & Howley, P. M. (1991). Biochemical and biological differences between E7 oncoproteins of the high- and low-risk human papillomavirus types are determined by amino-terminal sequences. *J Virol* **65**, 3943-8
- Narita, M.**, Nunez, S., Heard, E., Narita, M., Lin, A. W., Hearn, S. A., Spector, D. L., Hannon, G. J. & Lowe, S. W. (2003). Rb-mediated heterochromatin formation and silencing of E2F target genes during cellular senescence. *Cell* **113**, 703-16
- Narlikar, G. J.**, Fan, H. Y. & Kingston, R. E. (2002). Cooperation between complexes that regulate chromatin structure and transcription. *Cell* **108**, 475-87.
- Nielsen, S. J.**, Schneider, R., Bauer, U. M., Bannister, A. J., Morrison, A., O'Carroll, D., Firestein, R., Cleary, M., Jenuwein, T., Herrera, R. E. & Kouzarides, T. (2001). Rb targets histone H3 methylation and HP1 to promoters. *Nature* **412**, 561-5.
- Oh, K. J.**, Kalinina, A., Wang, J., Nakayama, K., Nakayama, K. I., and Bagchi, S. (2004a). The papillomavirus E7 oncoprotein is ubiquitinated by UbcH7 and Cullin 1- and Skp2-containing E3 ligase. *J Virol* **78**, 5338-5346.
- Okuda, M.**, Horn, H. F., Tarapore, P., Tokuyama, Y., Smulian, A. G., Chan, P. K., Knudsen, E. S., Hofmann, I. A., Snyder, J. D., Bove, K. E. & Fukasawa, K. (2000). Nucleophosmin/B23 is a target of CDK2/cyclin E in centrosome duplication. *Cell* **103**, 127-40.
- Osterloh, L.**, von Eyss, B., Schmit, F., Rein, L., Hubner, D., Samans, B., Hauser, S. & Gaubatz, S. (2007). The human synMuv-like protein LIN-9 is required for transcription of G2/M genes and for entry into mitosis. *Embo J* **26**, 144-57
- Pardee, A. B.** (1974). A restriction point for control of normal animal cell proliferation. *Proc Natl Acad Sci U S A* **71**, 1286-90.
- Pardee, A. B.** (1989). G1 events and regulation of cell proliferation. *Science* **246**, 603-8.

- Park JS**, Chee YH, Namkoong SE, Han SK, Kim TE, Lee HY, Kim SJ. 1994. Human papillomavirus detection in cervical carcinoma tissues and para-aortic lymph nodes by the polymerase chain reaction. *Gynecol Oncol*. 53(3):344-51.
- Petti L. Nilson L. A. DiMaio D.** 1991 Activation of the platelet-derived growth factor receptor by the bovine papillomavirus E5 transforming protein *EMBO J*. 10 845-855.
- Phelps, W. C., Munger, K., Yee, C. L., Barnes, J. A. & Howley, P. M.** (1992). Structure-function analysis of the human papillomavirus type 16 E7 oncoprotein. *J Virol* **66**, 2418-27.
- Pilkinton M**, Sandoval R, Barrett K, Tian X, Colamonici OR (2007). Mip/LIN-9 can inhibit cell proliferation independent of the pocket proteins. *Blood Cells Mol Dis* ;39(3):272-7.
- Pilkinton M**, Sandoval R, Colamonici OR (2007). Mammalian Mip/LIN-9 interacts with either the p107, p130/E2F4 repressor complex or B-Myb in a cell cycle-phase-dependent context distinct from the Drosophila dREAM complex. *Oncogene* ;26(54):7535-43.
- Pilkinton M**, Sandoval R, Song J, Ness SA, Colamonici OR (2007). Mip/LIN-9 regulates the expression of B-Myb and the induction of cyclin A, cyclin B, and CDK1. *J Biol Chem*. 5;282(1):168-75.
- Pilkinton, M., Sandoval, R., Song, J., Ness, S. A. & Colamonici, O. R.** (2007). Mip/LIN-9 regulates the expression of B-Myb and the induction of cyclin A, cyclin B, and CDK1. *J Biol Chem* **282**, 168-75.
- Polager, S., Kalma, Y., Berkovich, E. & Ginsberg, D.** (2002). E2Fs up-regulate expression of genes involved in DNA replication, DNA repair and mitosis. *Oncogene* **21**, 437-46.
- Rampias T, Sasaki C, Weinberger P and Psyrri A, 2009.** E6 and E7 gene silencing and transformed phenotype of human papillomavirus 16-positive oropharyngeal cancer cells. *J Natl cancer Inst*, **101**:412-423
- Rayman, J. B., Takahashi, Y., Indjeian, V. B., Dannenberg, J. H., Catchpole, S., Watson, R. J., te Riele, H. & Dynlacht, B. D.** (2002). E2F mediates cell cycle-dependent transcriptional

repression in vivo by recruitment of an HDAC1/mSin3B corepressor complex. *Genes Dev* **16**, 933-47.

Ren, S. & Rollins, B. J. (2004). Cyclin C/cdk3 promotes Rb-dependent G0 exit. *Cell* **117**, 239-51

Roman et al. (2006). The E7 proteins of low and high-risk human papillomaviruses share the ability to target the pRB family member p130 for degradation. *PNAS*, **103**, 437-442.

Rous P. and Beard JW 1935. The progression to carcinoma of virus-induced rabbit papilloma (shope). *J. exp. Med.* 79: 511-537.

Rous P. and Beard JW. 1934. Carcinomatous changes in virus-induced papillomas of the skin of the rabbit. *Proc. Soc. Exp. Biol. Med.* 62: 523-548.

Scheffner M, Werness BA, Huibregtse JM, Levine AJ. and Howley PM. 1990. The E6 oncoprotein encoded by human papillomavirus types 16 and 18 promotes the degradation of p53. *Cell*, 63, 1129–113.

Schmit et al. (2007). The human synMuv-like protein LIN-9 is required for transcription of G2/M genes and for entry into mitosis. *The EMBO Journal*, **26**, 144-157.

Schmit F, Korenjak M, Mannefeld M, Schmitt K, Franke C, von Eyss B, 2007. LINC, a human complex that is related to pRB-containing complexes in invertebrates regulates the expression of G2/M genes. *Cell cycle* **6**:1903-1913

Sedman J, Stenlund A. 1995. Co-operative interaction between the initiator E1 and the transcriptional activator E2 is required for replicator specific DNA replication of bovine papillomavirus *in vivo* and *in vitro*. *EMBO J.* 14(24):6218-28.

Seo YS, Muller F, Lusky M. & Hurwitz J. 1993. Bovine papilloma virus (BPV)-encoded E1 protein contains multiple activities required for BPV DNA replication. *Proc. Natl Acad. Sci. USA* 90, 702-706.

Sherr, C. J. (1995). D-type cyclins. *Trends Biochem Sci* **20**, 187-90

- Shoppe RE.** 1933. Infectious papillomatosis of rabbits. *J. Exp. Med.* 58: 607-627.
- Sima N, Wang W, Kong D, Deng Q Xu, Zhou G Xu, Meng L, Wang S and Ma D, 2008.** RNA interference against HPV16 E7 oncogene leads to viral E6 and E7 suppression in cervical cancer cells and apoptosis via upregulation of Rb and p53. *Apoptosis*, **13**:273-281
- Singer, A.** (1995). Cervical cancer screening: state of the art. *Baillieres Clin Obstet Gynaecol* **9**, 39-64.
- Smotkin, D. & Wettstein, F. O.** (1986). Transcription of human papillomavirus type 16 early genes in a cervical cancer and a cancer-derived cell line and identification of the E7 protein. *Proc Natl Acad Sci U S A* **83**, 4680-4.
- Snijders, P. J., Steenbergen, R. D., Heideman, D. A., and Meijer, C. J.** (2006). HPV-mediated cervical carcinogenesis: concepts and clinical implications. *J Pathol* **208**, 152-164.
- Sokolowski M, Furneaux H, Schwartz S.** 1999. The inhibitory activity of the AU-rich RNA element in the human papillomavirus type 1 late 3' untranslated region correlates with its affinity for the elav-like HuR protein. *J. Virol.* **73**: 1080-91.
- Solari, F. & Ahringer, J.** (2000). NURD-complex genes antagonise Ras-induced vulval development in *Caenorhabditis elegans*. *Curr Biol* **10**, 223-6.
- Stabel, S., P. Argos, and K. Philipson.** 1985. The release of growth arrest by microinjection of adenovirus E1A DNA. *EMBO J.* **4**:2329–2336.
- Storey, A., Pim, D., Murray, A., Osborn, K., Banks, L. & Crawford, L.** (1988). Comparison of the in vitro transforming activities of human papillomavirus types. *Embo J* **7**, 1815-20.
- Strauss MJ, Shaw EW, Bunting H, Melnick J.L.** 1949. “Cristalline” virus-like particles from skin papillomas characterized by intra-nuclear inclusion bodies. *Proc. Soc. Exp. Biol. Med.* **72**: 46-50.

- Strober**, B. E., Dunaief, J. L., Guha & Goff, S. P. (1996). Functional interactions between the hBRM/hBRG1 transcriptional activators and the pRB family of proteins. *Mol Cell Biol* **16**, 1576-83
- Takahashi**, Y., Rayman, J. B. & Dynlacht, B. D. (2000). Analysis of promoter binding by the E2F and pRB families in vivo: distinct E2F proteins mediate activation and repression. *Genes Dev* **14**, 804-16
- Tan W**, Felber BK, Zolotukhin AS, Pavlakis GN, Schwartz S. 1995. Efficient expression of the human papillomavirus type 16 L1 protein in epithelial cells by using Rev and the Rev-responsive element of human immunodeficiency virus or the cis-acting transactivation element of simian retrovirus type 1. *J Virol* 69: 5607-20.
- Tang *et al.* (2006)**. Short-term induction and long-term suppression of HPV16 oncogene silencing by RNA interference in cervical cancer cells. *Oncogene*, **25**, 2094-2104.
- Taubert**, S., Gorrini, C., Frank, S. R., Parisi, T., Fuchs, M., Chan, H. M., Livingston, D. M. & Amati, B. (2004). E2F-dependent histone acetylation and recruitment of the Tip60 acetyltransferase complex to chromatin in late G1. *Mol Cell Biol* **24**, 4546-56.
- Taylor-Harding**, B., Binne, U. K., Korenjak, M., Brehm, A. & Dyson, N. J. (2004). p55, the Drosophila ortholog of RbAp46/RbAp48, is required for the repression of dE2F2/RBF-regulated genes. *Mol Cell Biol* **24**, 9124-36
- Tedesco, D.**, Lukas, J., and Reed, S. I. (2002). The pRb-related protein p130 is regulated by phosphorylation-dependent proteolysis via the protein-ubiquitin ligase SCFSkp2. *Genes & Dev* **16**, 2946-2957.
- Thomas Farkas**, Klaus Hansen‡, Karin Holm, Jiri Lukas, and Jiri Bartek (2002). Distinct Phosphorylation Events Regulate p130- and p107-mediated Repression of E2F-4. *J. Biol. Chem* **277**, 30: 26741–26752

- Thomas**, D. M., Carty, S. A., Piscopo, D. M., Lee, J. S., Wang, W. F., Forrester, W. C. & Hinds, P. W. (2001). The retinoblastoma protein acts as a transcriptional coactivator required for osteogenic differentiation. *Mol Cell* **8**, 303-16.
- Thomas**, D. M., Carty, S. A., Piscopo, D. M., Lee, J. S., Wang, W. F., Forrester, W. C. & Hinds, P. W. (2001). The retinoblastoma protein acts as a transcriptional coactivator required for osteogenic differentiation. *Mol Cell* **8**, 303-16.
- Tommasino M and Crawford L., 1995.** Human Papillomavirus E6 and E7: proteins which deregulate the cell cycle. *Bioassays*, **17**:509-518
- Tommasino M.** 2001. The early genes of human papillomaviruses. In “Cancer Research-an Encyclopedic Reference”. (M. Schwab Editor), Springer, Heidelberg, Germany.
- Tschop K.**, Conery A.R., Lithovchick L., Decaprio J.A., Settleman J., Harlow E & Dyson N (2011). A kinase shRNA screen links LATS2 and the pRB tumour suppressor. *Genes Dev* **25**, 814-830.
- Vairo**, G., Livingston, D. M. & Ginsberg, D. (1995). Functional interaction between E2F-4 and p130: evidence for distinct mechanisms underlying growth suppression by different retinoblastoma protein family members. *Genes Dev* **9**, 869-81.
- Vandel**, L., Nicolas, E., Vaute, O., Ferreira, R., Ait-Si-Ali, S. & Trouche, D. (2001). Transcriptional repression by the retinoblastoma protein through the recruitment of a histone methyltransferase. *Mol Cell Biol* **21**, 6484-94.
- Watanabe**, H., Pan, Z. Q., Schreiber-Agus, N., DePinho, R. A., Hurwitz, J. & Xiong, Y. (1998). Suppression of cell transformation by the cyclin-dependent kinase inhibitor p57KIP2 requires binding to proliferating cell nuclear antigen. *Proc Natl Acad Sci U S A* **95**, 1392-7.
- Weinberg**, R. A. (1995). The retinoblastoma protein and cell cycle control. *Cell* **81**, 323-30.

- Weinmann**, A. S., Yan, P. S., Oberley, M. J., Huang, T. H. & Farnham, P. J. (2002). Isolating human transcription factor targets by coupling chromatin immunoprecipitation and CpG island microarray analysis. *Genes Dev* **16**, 235-44.
- Weintraub**, S. J., Chow, K. N., Luo, R. X., Zhang, S. H., He, S. & Dean, D. C. (1995). Mechanism of active transcriptional repression by the retinoblastoma protein. *Nature* **375**, 812-5.
- Weintraub**, S. J., Prater, C. A. & Dean, D. C. (1992). Retinoblastoma protein switches the E2F site from positive to negative element. *Nature* **358**, 259-61.
- Won**, K. A., Xiong, Y., Beach, D. & Gilman, M. Z. (1992). Growth-regulated expression of D-type cyclin genes in human diploid fibroblasts. *Proc Natl Acad Sci U S A* **89**, 9910-4.
- Woodworth**, C. D., Doniger, J. & DiPaolo, J. A. (1989). Immortalization of human foreskin keratinocytes by various human papillomavirus DNAs corresponds to their association with cervical carcinoma. *J Virol* **63**, 159-64.
- Yang**, L., Li, R., Mohr, I. J., Clark, R. & Botchan, M. R. 1991 BPV-1 replication *in vitro* is activated by the transcription factor E2. *Nature* **353**, 628-632.
- Yasumoto**, S., Burkhardt, A. L., Doniger, J. & DiPaolo, J. A. (1986). Human papillomavirus type 16 DNA-induced malignant transformation of NIH 3T3 cells. *J Virol* **57**, 572-7.
- Zamanian**, M. & La Thangue, N. B. (1993). Transcriptional repression by the Rb-related protein p107. *Mol Biol Cell* **4**, 389-96.
- Zerfass**, K., Schulze, A., Spitzkovsky, D., Friedman, V., Henglein, B. & Jansen-Durr, P. (1995). Sequential activation of cyclin E and cyclin A gene expression by human papillomavirus type 16 E7 through sequences necessary for transformation. *J Virol* **69**, 6389-99.
- Zerfass-Thome**, K., Zworschke, W., Mannhardt, B., Tindle, R., Botz, J. W. & Jansen-Durr, P. (1996). Inactivation of the cdk inhibitor p27KIP1 by the human papillomavirus type 16 E7 oncoprotein. *Oncogene* **13**, 2323-30.

- Zhang, B., Chen, W. & Roman, A.** (2006). The E7 proteins of low- and high-risk human papillomaviruses share the ability to target the pRB family member p130 for degradation. *Proc Natl Acad Sci U S A* **103**, 437-42.
- Zhao C, Tan W, Sokolowski M, Schwartz S.** 1996. Identification of nuclear and cytoplasmic proteins that interact specifically with an AU-rich, cis-acting inhibitory sequence in the 3' untranslated region of human papillomavirus type 1 late mRNAs. *Virology*. 70: 3659-67.
- Zhu, H., Nie, L. & Maki, C. G.** (2005). Cdk2-dependent Inhibition of p21 stability via a C-terminal cyclin-binding motif. *J Biol Chem* **280**, 29282-8
- zur Hausen H** (2000). Papillomaviruses causing cancer: evasion from host-cell control in early events in carcinogenesis. *J Natl Cancer Inst.* 3;92(9):690-8
- zur Hausen H.** 1976. Condylomata acuminata and human genital cancer. *Cancer Res.* 36(2 pt 2):794.
- zur Hausen, H.** (1996). Papillomavirus infections--a major cause of human cancers. *Biochim Biophys Acta* **1288**, F55-78.
- zur Hausen, H.** (2002). Papillomaviruses and cancer: from basic studies to clinical application. *Nat Rev Cancer* **2**, 342-350.