

LIST OF TABLES

| Table No. | Title | Page |
|-----------|---|------|
| 2.1 | Characteristics of the soil-transmitted helminth (STH) | 20 |
| 2.2 | Specific and general clinical manifestation of soil-transmitted helminth infection (STH) infections | 26 |
| 3.1 | Geographical distribution of the Orang Asli subgroups participated in this study stratified by location | 75 |
| 3.2 | Distribution demographic characteristics of survey population stratified by subgroup | 76 |
| 3.3 | Distribution socioeconomic, environmental sanitation and personal hygiene characteristics of survey population stratified by subgroup | 77 |
| 3.4 | Prevalence of soil-transmitted helminth (STH) infections according to population characteristic | 79 |
| 3.5 | Prevalence of single, double and triple infection of soil-transmitted helminth (STH) stratified by subgroup and age group | 83 |
| 3.6 | Odd ratios of the potential risk factor associated with soil-transmitted helminth (STH) infections in Temuan subgroup | 84 |
| 3.7 | Odd ratios of the potential risk factor associated with soil-transmitted helminth (STH) infections in Semelai subgroup | 85 |
| 3.8 | Odd ratios of the potential risk factor associated with soil-transmitted helminth (STH) infections in Jakun subgroup | 86 |
| 3.9 | Odd ratios of the potential risk factor associated with soil-transmitted helminth (STH) infections in Orang Kuala subgroup | 87 |
| 3.10 | Odd ratios of the potential risk factor associated with soil-transmitted helminth (STH) infections in Mah Meri subgroup | 88 |
| 4.1 | Overall prevalence of intestinal helminths in dogs and cats | 114 |
| 4.2 | Prevalence of intestinal helminths by species in dogs and cats | 114 |
| 4.3 | Prevalence of monoparasitism and poliparasitism in dogs and cats | 115 |
| 4.4 | Types of intestinal helminths in poliparasitism dogs and cats | 115 |
| 5.1 | Results achieved by real-time PCR coupled HRM analysis of ITS-2 amplicon from control genomic DNA for hookworm | 148 |

| | | |
|-----|---|-----|
| 5.2 | Hookworm infections detected by microscopy, conventional PCR and real-time PCR-HRM assay of human fecal samples in Peninsular Malaysia | 148 |
| 5.3 | <i>Necator americanus</i> and <i>Ancylostoma ceylanicum</i> infections in human fecal samples detected by real-time PCR-HRM assay according to villages in Peninsular Malaysia | 149 |
| 5.4 | Comparison between microscopy, conventional semi-nested PCR and real-time PCR-HRM assays | 149 |
| 5.5 | Hookworm species detected via both conventional semi-nested PCR and real-time PCR-HRM assays | 150 |
| 6.1 | Hookworm infections detected by microscopy and conventional PCR assay of dogs and cats fecal samples in Peninsular Malaysia | 174 |
| 6.2 | Hookworm species in dogs and cats fecal sample detected by conventional PCR assay according to locations in Peninsular Malaysia | 175 |
| 6.3 | Multiple alignments of <i>cox 1</i> gene of <i>A. ceylanicum</i> sequences from this study with the reference sequences obtained from GenBank, representing <i>A. caninum</i> and <i>A. duodenale</i> | 178 |
| 7.1 | Summary of descriptive database on the prevalence survey of soil-transmitted helminth (STH) infections between 1970 and 2012 in Peninsular Malaysia | 214 |
| 7.2 | Mean prevalence of individual and estimated combined STH infections between 1970 and 2012 | 215 |
| 7.3 | Regression coefficients used to estimate probability of person being infected in logistic regression model | 215 |
| 7.4 | Details of estimates of predicted prevalence, number being infected (total population and school-aged children) and number warranting mass treatment using 50% and 20% threshold at district level | 216 |
| H.1 | GenBank accession numbers of ITS-2 ribosomal RNA gene of <i>Necator americanus</i> isolated from human | 314 |
| H.2 | GenBank accession numbers of ITS-2 ribosomal RNA gene of <i>Ancylostoma ceylanicum</i> isolated from human, dogs and cats | 315 |

| | | |
|-----|---|-----|
| H.3 | GenBank accession numbers of ITS-2 ribosomal RNA gene of <i>Ancylostoma caninum</i> isolated from dogs and cats | 316 |
| H.4 | GenBank accession numbers of ITS-2 ribosomal RNA gene of <i>Ancylostoma braziliense</i> isolated from cat | 317 |
| H.5 | GenBank accession numbers of <i>cox 1</i> gene of <i>Ancylostoma ceylanicum</i> isolated from human, dogs and cats | 318 |
| I.1 | Summary of the Remote Sensing (RS) data and sources | 319 |
| K.1 | Regression coefficients used to estimate probability of infection being greater 50% or more | 321 |
| L.1 | Summary descriptive of the database on the prevalence of the STH infections in Malaysia | 323 |
| O.1 | Summary estimates of predicted prevalence, number being infected (total population and school-aged children) and number warranting mass treatment using 50% and 20% threshold at district level | 328 |
